



## REGIONAL ILEITIS



# REGIONAL ILEITIS

By

BURRILL B CROHN, M D

*Consulting Gastroenterologist Mount  
Sinai Hospital, New York*



GRUNE & STRATTON  
NEW YORK 1949

Copyright 1949  
GRUVE & STRATTON INC  
381 Fourth Avenue  
New York 16 N Y

Printed by L. V. P. nting Co. L Albany N Y  
Distributed by Moe & Co. Inc Baltimore Md  
Copyright by Gruve & Stratton Inc. All rights reserved.

## Preface

---

THE PROMPT ACCEPTANCE of the concept of regional ileitis by the profession at large has led, in the last few years to a wide and elaborate knowledge of the clinical aspects and surgical treatment of the malady. The literature on the subject extends over several hundred published references with a widespread distribution throughout the world.

To attempt to write the life history of a disease requires extended experience with its pathologic aspects, the intensive study of ample clinical samplings and a numerous and illustrative case material. Fifteen years is too short a period in the limited world of scientific and human experience to encompass more than a general concept of the life cycle of an entity. This is particularly true when the process of extension progresses most often slowly, and when intelligent recognition of the various stages becomes possible only after long and intimate association with the disease. However, by pooling the published observations of the many students in the field and by using such a vast data as a measuring standard with which to enlighten one's own more limited experiences, the task of organizing the progressive story of a disease can be attempted with greater confidence.

The etiology of regional ileitis at least as to a specific causative agency remains unknown. The clinical concept and diagnosis and the roentgenologic recognition is well advanced and accepted. The surgical cure or attempts at cure are still in a state of flux. Whatever skepticism

occurs is related to the field of surgery where frequent disappointment has been experienced and criticism has been levelled. Such scientific criticism is healthy and when constructive, can lead only to advance in thought and in practical knowledge.

Least understood is the field of ileo jejunitis, that affection of the higher segment of the small bowel which is rapidly being explored, its inflammatory and deficiency manifestations tabulated and organized. So too with the combined forms of ileitis and colitis which present a tangled skein of primary and secondary affects leaving much still to be studied.

This offering, begun as a brief monograph, has developed into a book, not from verbosity, but because the extensive literature and many worthy contributions from students throughout the world have made the extension necessary as a recognition of the wide spread interest of internists and surgeons in a subject which advances daily.

BURRILL B. CROHN M.D.

New York City  
February 1949

# Contents

PREFACE	v
CHAPTER	
1 GENERAL CONSIDERATIONS SCOPE OF STUDY	1
Historical Background 2 Regional Ileitis 3 Scope of the Present Study 7	
2 REGIONAL ILEITIS—ETIOLOGY (222 CASES)	9
Etiology 9 Specific Agencies 10 Protozoa 12 Bacterial Dysentery 13 Viruses 14 Sarcinosis 16 Trauma 17 Familial Incidence of Ileitis as a Possible Etiologic Factor 21 Theories Regarding the Pathologic Mechanism in the Creation of Ileitis 23 Vascular Change 24 The Theory of Lymphatic Block 25 Age in Regional Ileitis 28 Social Status Race Religion Nationality 30	
3 GROSS PATHOLOGY	34
Microscopic Pathology 41	
4 CLINICAL FEATURES	46
Onset 46 Physical Examination 54 Obstructive Phenomena 57	
5 FISTULA FORMATION IN REGIONAL ILEITIS	59
Internal Fistula 61 Perirectal Fistulas 67 Method of Anal Fistulation 67	
6 THE COURSE IN REGIONAL ILEITIS	70
Group 1 71 Group 2 Chronic Regional Ileitis without Progression 75 Group 3 Chronic Regional Ileitis with Progressive Extension 79	
7 COMPLICATIONS OF REGIONAL ILEITIS	82
Perforation of Intestine 82 Hemorrhage 85 General Complications 83 Gynecologic Manifestations 86	



occurs is related to the field of surgery where frequent disappointment has been experienced and criticism has been levelled. Such scientific criticism is healthy and when constructive, can lead only to advance in thought and in practical knowledge.

Least understood is the field of ileo jejunitis, that affection of the higher segment of the small bowel which is rapidly being explored, its inflammatory and deficiency manifestations tabulated and organized. So too with the combined forms of ileitis and colitis which present a tangled skein of primary and secondary affects leaving much still to be studied.

This offering, begun as a brief monograph, has developed into a book, not from verbosity, but because the extensive literature and many worthy contributions from students throughout the world have made the extension necessary as a recognition of the wide spread interest of internists and surgeons in a subject which advances daily.

BURRILL B. CROHN, M.D.

New York City  
February 1949

# Contents

PREFACE	v
CHAPTER	
1 GENERAL CONSIDERATIONS SCOPE OF STUDY	1
Historical Background 2 Regional Ileitis 5 Scope of the Present Study 7	
2 REGIONAL ILEITIS—ETIOLOGY (222 CASES)	9
Etiology 9 Specific Agencies 10 Protozoa 12 Bacterial Dysentery 13 Viruses 14 Sarcoidosis 16 Trauma, 17 Familial Incidence of Ileitis as a Possible Etiologic Factor 21 Theories Regarding the Pathologic Mechanism in the Creation of Ileitis 23 Vascular Change 24 The Theory of Lymphatic Block 25 Age in Regional Ileitis 28 Social Status Race Religion Nationality 30	
3 GROSS PATHOLOGY	34
Microscopic Pathology 41	
4 CLINICAL FEATURES	46
Onset 46 Physical Examination 54 Obstructive Phenomena 57	
5 FISTULA FORMATION IN REGIONAL ILEITIS	59
Internal Fistulas 61 Perirectal Fistulas 67 Method of Anal Fistulotomy 67	
6 THE COURSE IN REGIONAL ILEITIS	70
Group 1 71 Group 2 Chronic Regional Ileitis without Progression 75 Group 3 Chronic Regional Ileitis with Progressive Extension 79	
7 COMPLICATIONS OF REGIONAL ILEITIS	82
Perforation of Intestine 83 Hemorrhages 82 General Complications 83 Gynecologic Manifestation 86	

8	ROENTGENOGRAPHIC STUDY OF REGIONAL ILEITIS	88
	The Technique of Radiography 98	
9	THE DIAGNOSIS OF REGIONAL ILEITIS	103
	Differential Diagnosis 107	
10	CONSIDERATIONS REGARDING THERAPY IN REGIONAL ILEITIS	113
	Diet 115   Conservative Palliative Treatment of Regional Ileitis 116 (general directions 116   sample menu 116   diet 117   ambulation 119   various medications 120)	
11	SURGICAL TREATMENT OF CHRONIC REGIONAL ILEITIS	123
	Diagnosis and Treatment of Recurrences 134   The Treatment of Recurrent Ileitis 139	
12	ACUTE REGIONAL ILEITIS	142
	Distinct Entity or Part of Chronic Regional Ileitis? 149   The Eventual Outcome of Cases of Acute Ileitis 150   The Medical or Surgical Treatment of Acute Ileitis 152	
13	ILEO JEJUNITIS	156
	The Various Forms of Ileo Jejunitis 160   Etiology 166   Pathology 167   Symptomatology 169   Roentgenographic Recognition of Ileo Jejunitis 178   Differential Diagnosis 182	
14	ILEO JEJUNITIS—TREATMENT	186
	Course and Prognosis 186   Medical Treatment 188   Surgical Intervention 191	
15	ILEO COLITIS—COMBINED FORMS	200
	Etiology 204   Clinical Features 206   The Physical Signs 206   Prognosis 208   Medical Treatment 209   Surgical Approach 209	
	REFERENCES	213
	INDEX	225

# 1 General Considerations Scope of Study

---

THE TERM 'Regional Ileitis' was coined and created in 1932 to describe what was considered as a new pathologic and clinical entity, a clear cut disease affecting the terminal segment or segments of the small bowel characterized by a subacute or chronic necrotizing and ulcerating inflammation. While the term regional ileitis is still the most popular denomination of the disease many other descriptive names have been created by successive authors to cover new and expanding concepts of the anatomic and pathologic aspects of this malady. Terminal ileitis was the first term applied because in the original publication the first 14 cases were all restricted anatomically to the terminal 8 to 12 inches of the ileum. Regional ileitis however denoted a more widespread distribution. 'regional enteritis' was suggested by Brown, Baughn and Weber. 'chronic ulcerating enteritis' by Harris Bell and Brunn and by Cushman. 'chronic ulcerative enteritis' (Carr and Bocock). 'segmental enteritis' (Lewisohn) to denote both the tendency to scar formation and the more widespread anatomic distribution. 'ileo colitis' was suggested by Erb and Farmer and again by Colp to indicate the not infrequent involvement of the colon. To this concept must now be added 'ileo jejunitis' the same pathologic process extended to include anatomically all of the jejunum continuously or segmentally and finally 'ileo colitis' or 'colo ileitis' a term created to cover those combined cases where some

segments of the large as well as the small bowel are concerned.

As originally described, regional ileitis or enteritis is a nonspecific inflammatory disease, one of the larger group of intestinal granulomata, a process affecting mainly young adults and characterized by an acute, a subacute and a chronic phase of necrotizing ulcerating and cicatrizing inflammation. The disease is further marked by its tendency to fistula formation within adjoining loops of the small and large bowel, by external fistulas to the abdominal wall, by frequent rectal suppurative complications. A disproportionate increase of connective tissue may and does frequently lead to intestinal stenosis and obstruction. Clinically, the disease is featured by fever, diarrhea, loss of weight, a secondary anemia, and by the formation of an intra abdominal mass, constituted of and by approximated loops of inflamed intestine frequently joined together by internal fistulas. The process regularly begins or stops at the ileocecal valve, the exceptions to this localization being few in number. The disease advances upward by "skip lesions" separated by healthy "skipped areas." It is a progressive lesion in many instances, slowly or rapidly advancing; in others it will remain for twenty or more years strictly localized in the terminal ileum. Recurrences of the inflammatory process after palliative or radical surgical procedures further characterize the process as an active, persistent and recurrent disease.

#### HISTORICAL BACKGROUND

Nonspecific granulomata of the alimentary tract involving mainly the large bowel have been recognized for many years. The European medical literature has been replete with innumerable examples of such inflammatory masses located in large and small intestine some even in the stomach included in these descriptions undoubtedly are many instances of what we recognize now as typical ileitis or enteritis. In 1813, Combe and Saunders reported before

the Royal College of Physicians of London "A Singular Case of Stricture and Thicening of the Ileum" To quote in part "The lower part of the ileum, as far as the colon was contracted for the space of three feet to the size of a turkey quill" John Abercrombie discussed pathologic states of the ileum and reported cases very similar to regional ileitis His description published in 1828 is instructive "A girl aged 13, about a year before her death (1814) began to be afflicted with pain of the abdomen and frequent vomiting The cecum coli three inches along the ascending colon and the lower end of the ileum to the extent of about eighteen inches, was distended, thickened in the coats, externally of a reddish color and internally covered by numerous well defined ulcers varying in size from a diameter of a split pea to that of a sixpence" After a lapse of almost a century, the English literature again contained a case report by Movshian in 1907 entitled, "The Mimicry of Malignant Disease in the Large Intestine"

In the meantime, the German literature began rapidly to show an abundance of references to non-specific granulomata of inflammatory nature Braun in 1901 quotes 5 cases mostly of traumatic or hernial origin and later quotes 25 more cases from the current periodicals up to 1909 By 1920, Tietze had collated 281 references on this subject including at least 2 cases of typical regional ileitis (Lowen and also Willman) The literature now increased apace Dalziel contributing an excellent article The discussions in the German publications show an increasing interest Korte 1921 Landois 1923 Konjetzny 1932 report cases the whole subject crystallizing with a paper by Fischer and Lurmann This paper included the description of three cases of typical regional or terminal ileitis and recognized and granted priority to the original publication less than a year previous in this country [66] The discussion before the German Congress and thereafter was amplified by the experience of such noteworthy surgeons as von Hacker Gisbertz Peters Anschütz Fenster and others

segments of the large as well as the small bowel are concerned

As originally described, regional ileitis or enteritis is a nonspecific inflammatory disease, one of the larger group of intestinal granulomata, a process affecting mainly young adults and characterized by an acute, a subacute and a chronic phase of necrotizing ulcerating and cicatrizing inflammation. The disease is further marked by its tendency to fistula formation within adjoining loops of the small and large bowel, by external fistulas to the abdominal wall, by frequent rectal suppurative complications. A disproportionate increase of connective tissue may and does frequently lead to intestinal stenosis and obstruction. Clinically, the disease is featured by fever, diarrhea, loss of weight, a secondary anemia, and by the formation of an intra abdominal mass, constituted of and by approximated loops of inflamed intestine frequently joined together by internal fistulas. The process regularly begins or stops at the ileocecal valve, the exceptions to this localization being few in number. The disease advances upward by "skip lesions" separated by healthy "skipped areas." It is a progressive lesion in many instances, slowly or rapidly advancing, in others it will remain for twenty or more years strictly localized in the terminal ileum. Recurrences of the inflammatory process after palliative or radical surgical procedures further characterize the process as an active, persistent and recurrent disease.

#### HISTORICAL BACKGROUND

Nonspecific granulomata of the alimentary tract, involving mainly the large bowel have been recognized for many years. The European medical literature has been replete with innumerable examples of such inflammatory masses located in large and small intestine some even in the stomach included in these descriptions undoubtedly are many instances of what we recognize now as typical ileitis or enteritis. In 1813, Combe and Saunders reported before

recognized, tuberculosis, syphilis and actinomycosis were excluded by definition.

### REGIONAL ILEITIS [66]

From this large and general group of non-specific granulomata of the intestinal tract a subspecies which was termed regional ileitis was isolated, a distinction which was possible because this group was characterized by a consistent anatomic distribution, a characteristic clinical course, a progressive life history, typical complications of fistula formation and obstruction, thus group being amenable further to successful surgical interference. It is interesting to note historically the amplification of the original concept of terminal ileitis. The first report covered 14 cases of a typically characteristic disease involving only the terminal segment or segments of the ileum. Acute cases were included among those running the more easily recognized chronic inflammatory course. Shortly thereafter the involvement in some cases of the upper reaches of the ileum and all or part of the jejunum, as noted by Harris, Bell and Brunn led to the identification of ileojejunitis either as an extension or as a continuation of terminal ileitis or as a subgroup per se in which the involvement of the terminal segments of the small intestine was subordinated to the process higher up. The participation of the colon segmentally either by continuity from the terminal ileum or by scattered areas of coincidental disease was first noted by Colp and by Crohn and Rosenak and led to the creation of still another subgroup termed "ileocolitis" (Crohn and Bevan), a most confused and complicated pattern of involvement of ileum and various segments of the large bowel. In this latest subgroup the ileitis is regarded as the parent focus of infection the colonic involvement being secondary in time and extension.

Three years after the first designation of regional enteritis as an entity Brunn was able to collect 267 cases



By 1929, American writers had become cognizant of the subject of nonspecific granulomata of the intestinal tract. Coffen (1925) Horsley (including two fine case descriptions of ileitis), an excellent study by Moschcowitz and Wilensky (including an admirable photographic reproduction of ileitis) of 4 cases of granulomata, of which one was typical of ileitis as we now know it. Mock in 1931 summarized the subject of infective granulomata, attributing most of these pathologic masses to some interference with vascular supply to a given tissue, resulting in local necrosis followed by a proliferative reparative tissue reaction. He considered trauma as the most common etiologic agent. His cases, many of them due to penetrating foreign bodies, diverticulitis, strangulated hernia, silk and metallic ligatures, involved the sigmoid, stomach, omentum, splenic flexure and in one case participation of the terminal ileum. Golob added one case, and Erdmann and Burt 3 more cases, 2 of which had typical involvement of the terminal ileum with intestinal obstruction and of internal fistulas from ileum to cecum. Finally, in 1933 Ginzburg and Oppenheimer reported a group of 52 cases of nonspecific granulomata of the intestinal tract cases due to vascular disturbances, to herniation, to sealed off intestinal perforations and included several cases of the entity which had already been isolated and termed regional ileitis.

This large group of nonspecific inflammatory granulomata involved any or all parts of the alimentary tract and omentum. They formed masses often obstructive in nature, usually simulating in their clinical course the picture of a new growth. They were characterized by a piling up of granulation tissue with various stages of ulceration necrosis and fibroblastic proliferative secondary reactions. The histologic process consisted of round cell infiltration fibroblasts connective tissue stroma and new blood vessel formation with the occasional occurrence of giant cells of unusual magnitude (Moschcowitz and Wilensky). Etiologic agents apart from trauma and foreign bodies were not

and its subsequent course has been traced Above all, prognosis has been learned and finally the medical and surgical approaches to therapy have been tried, some discarded newer methods apprehended The whole problem of recurrences has come to light and been observed and a rational method of recognition and treatment has been attempted

### SCOPE OF THE PRESENT STUDY

This study is an analysis of instances of segmental enteritis, cases which have been under my own personal office and hospital care during all or part of the last sixteen to eighteen years They have all been private patients who have been observed carefully, treated and subjected to continuous analysis including follow up during this period Some of these patients were included in the original group that constituted the first fourteen cases published in 1932 as terminal ileitis The remainder have been registered during all of the succeeding years, the last additions to the list having terminated approximately six months ago to allow for some if not adequate follow up (October 1947) Ward cases in hospitals are not included—those that are listed have been accepted only because they solicited treatment and were registered as personal patients even though the previous surgical or medical treatment had been administered elsewhere For purposes of analysis and study this larger group may be subdivided into four essential categories

1 Regional ileitis—the chronic form	29 cases
2 Acute ileitis	16 cases
3 Ileo-jejunitis	38 cases
4 Combined cases—ileitis with involvement of some segment of the large bowel	22 cases
Total	105 cases

All four categories represent types of regional or circumscribed enteritis They are not distinct entities in fact, mixed forms are not unusual The identical pathogenesis and

from the literature, in 1939 Ravdin and Johnston catalogued 393 cases from current periodicals. Confirmatory reports from the English literature were not long in making their appearance (Jackman, Hadfield, Barbout and Stokes, Molesworth, Hurst and his co authors), as likewise from the Dutch literature (Groen and Pompen, Snapper, Pompen and Groen).

A dissident viewpoint was advanced by several writers who considered regional ileitis a clinical but not a pathologic entity because of the occurrence of the disease at various other areas of the intestines. Characteristic lesions involving the jejunum were shown in 2 cases (Harris, Bell and Brunn), of 18 cases reported from the Mayo Clinic in 1934 (Brown, Bergen and Weber) the ileum was involved in 9, the jejunum in 3, the terminal ileum and parts of the colon in 6 instances. Homans and Hass, while recognizing typical terminal ileitis, questioned the localization of the process and felt that the jejunum was more often involved than was then recognized. They thought that the appendix was possibly an important etiologic factor, they further remarked for the first time that the disease might have a predilection for Hebrews (Mixer). Meyer and Rosi emphasized the involvement of the mesentery particularly in fistula formation. Erb and Farmer emphasized the acute phase in children as it simulated appendicitis. Reichert and Mathes made the first suggestion regarding etiology namely, experimental lymphoedema by blocking lymphatics. Goldfarb and later Kantor gave the first veritable description of the roentgen appearance of the disease, the latter terming the appearance of the disease in the terminal ileum as the 'string sign'. Actually considerable knowledge has been added to the original concept of terminal or regional ileitis in the last sixteen years. The anatomic and pathologic states have been extended to include the upper reaches of the ileum and all of the jejunum. Opportunity has been afforded for studying the life history of the disease its progression and extensions. Acute ileitis has been analyzed

and its subsequent course has been traced. Above all, prognosis has been learned, and finally the medical and surgical approaches to therapy have been tried, some discarded newer methods apprehended. The whole problem of recurrences has come to light and been observed and a rational method of recognition and treatment has been attempted.

### SCOPE OF THE PRESENT STUDY

This study is an analysis of instances of segmental enteritis, cases which have been under my own personal office and hospital care during all or part of the last sixteen to eighteen years. These have all been private patients who have been observed carefully, treated and subjected to continuous analysis including follow up during this period. Some of these patients were included in the original group that constituted the first fourteen cases published in 1932 as terminal ileitis. The remainder have been registered during all of the succeeding years, the last additions to the list having terminated approximately six months ago to allow for some if not adequate follow up (October 1947). Ward cases in hospitals are not included—those that are listed have been accepted only because they solicited treatment and were registered as personal patients even though the previous surgical or medical treatment had been administered elsewhere. For purposes of analysis and study this larger group may be subdivided into four essential categories.

1 Regional ileitis—chronic form	22 cases
2 Acute ileitis	16 cases
3 Ileocecalitis	18 cases
4 Combined cases—ileitis with involvement of some segment of the large bowel	22 cases
Total	78 cases

All four categories represent types of regional or circumscribed enteritis. They are not distinct entities; in fact, mixed forms are not unusual. The identical pathogenesis and

from the literature, in 1939 Ravdin and Johnston catalogued 393 cases from current periodicals. Confirmatory reports from the English literature were not long in making their appearance (Jackman, Hadfield, Barbour and Stokes, Molesworth, Hurst and his co authors), as likewise from the Dutch literature (Groen and Pompen, Snapper, Pompen and Groen).

A dissident viewpoint was advanced by several writers who considered regional ileitis a clinical but not a pathologic entity because of the occurrence of the disease at various other areas of the intestines. Characteristic lesions involving the jejunum were shown in 2 cases (Harris, Bell and Brunn), of 18 cases reported from the Mayo Clinic in 1934 (Brown, Birgen and Weber) the ileum was involved in 9, the jejunum in 3, the terminal ileum and parts of the colon in 6 instances. Homans and Hass, while recognizing typical terminal ileitis, questioned the localization of the process and felt that the jejunum was more often involved than was then recognized. They thought that the appendix was possibly an important etiologic factor, they further remarked for the first time that the disease might have a predilection for Hebrews (Mixer). Meyer and Rosi emphasized the involvement of the mesentery particularly in fistula formation. Erb and Farmer emphasized the acute phase in children as it simulated appendicitis. Reichert and Mathes made the first suggestion regarding etiology, namely, experimental lymphoedema by blocking lymphatics. Goldfarb and later Kantor gave the first veritable description of the roentgen appearance of the disease, the latter terming the appearance of the disease in the terminal ileum as the 'string sign'. Actually considerable knowledge has been added to the original concept of terminal or regional ileitis in the last sixteen years. The anatomic and pathologic states have been extended to include the upper reaches of the ileum and all of the jejunum, opportunity has been afforded for studying the life history of the disease its progression and extensions. Acute ileitis has been analyzed

## 2 Regional Ileitis—Etiology (222 cases)

---

REGIONAL ILEITIS is a nonspecific chronic recurrent granulomatous disease affecting mainly young adults, and characterized by a necrotizing, ulcerating, inflammatory process, one in which necrotizing elements in the lumen, in cases is an important feature. The terminal ileum is apparently the seat and origin of the pathologic process though the whole ileum may be involved. Typically the process rests at the ileocecal valve, transcending that barrier only exceptionally. The disease clinically is marked by a low grade fever by abdominal pain most frequently by mild diarrhea and by progressive loss of weight and a secondary anemia. The occurrence of an intra-abdominal mass is frequently noted. The most striking clinical feature is the formation of fistulas of ileal origin, these fistulas leading from ileum to the abdominal wall or from ileum to other loops of bowel or other hollow viscera. Rectal complications of a suppurative nature constitute a striking manifestation of the disease. The pathologic process is a progressive and enduring one lasting over the course of many years. Eventually intestinal obstruction and hemorrhage may intervene. Spontaneous cures are minimal but may occur. Surgical intervention is still the therapeutic method of choice.

### ETIOLOGY

The true causation of ileitis remains unknown in spite of diligent efforts to a certain a bacterial or specific agency

pathologic histology is common to them all. But they differ sufficiently clinically and anatomically to merit their consideration as separate groups, each group has identifying characteristics, has its own life history and prognosis as well as its individual approach to therapy.

The largest group of 222 cases of ileitis comprises the bulk of the study and provides the greatest number of the classic variety of the disease in which the ileum, particularly the terminal segments of small bowel, are involved. The disease being one of low grade pathogenicity and chronicity requires that this group be regarded as chronic regional ileitis. In contradistinction, the acute cases have been separately considered because of their own identifying characteristics though some confusion will arise when it attempts are made to separate true primary acute ileitis from acute exacerbations of chronic recurrent ileitis.

The group of ileocolitis has been separately considered because the involvement of the whole or of most of the small bowel introduces new elements of clinical significance and misses many of the distinguishing characteristics of the terminal form. Its life history is different, its therapeutic approach is entirely singular, its differential diagnosis and prognosis call for a more novel viewpoint. The fourth category of combined ileitis and colitis is so confusing with so many cross elements and complications and is so difficult a problem in therapy particularly its successful surgical treatment, that it requires its own generic consideration.

## 2 Regional Ileitis—Etiology (222 cases)

---

REGIONAL ILEITIS is a nonspecific chronic recurrent granulomatous disease affecting mainly young adults, and characterized by a necrotizing ulcerating inflammatory process one in which caseating elements in the long enduring cases is an important feature. The terminal ileum is apparently the seat and origin of the pathologic process though the whole ileum may be involved. Typically, the process rests at the ileocecal valve, transascending that barrier only exceptionally. The disease clinically is marked by a low grade fever by abdominal pain, most frequently by mild diarrhea and by progressive loss of weight and a secondary anemia. The occurrence of intra-abdominal mass is frequently noted. The most striking clinical feature is the formation of fistulas of ileal origin these fistulas leading from ileum to the abdominal wall, or from ileum to other loops of bowel or other hollow viscera. Rectal complications of a suppurative nature constitute a striking manifestation of the disease. The pathologic process is a progressive and enduring one lasting over the course of many years. Eventually intestinal obstruction and hemorrhage may intervene. Spontaneous cures are minimal but may occur. Surgical intervention is still the therapeutic method of choice.

### ETIOLOGY

The true causation of ileitis remains unknown in spite of diligent efforts to ascertain a bacterial or specific agency.



as the factor in its initiation. It is of first importance to exclude tuberculosis as the specific infecting agency. Much confusion in the past evolved from the inclusion of cases of nonspecific ileitis within the category of hypertrophic ileocecal tuberculosis. All resected specimens of ileitis have been conscientiously subjected to careful histologic study, to careful staining methods for the recognition of the Koch bacillus in cut sections, to cultures and animal inoculation.

Material from the bowel wall of the ileum and from mesenteric lymph nodes in resected cases were inoculated into guinea pigs, rabbits and chickens and proved negative in 5 of the original cases. Subsequently, whenever a suspicion of tuberculosis has arisen, the most careful procedures have been followed to eliminate the Koch factor as an etiologic agency. When tubercle bacilli have been demonstrated in cut sections, or when guinea pig inoculations have been positive, the case has been considered as intestinal tuberculosis and has been excluded from the series. Singularly this has been necessary only once during the eighteen years of study, with only one case of primary intestinal tuberculosis simulating ileitis. Frazer and Haggan reported a case of regional ileitis resected, in which the mesenteric lymph nodes showed old caseating tuberculosis; the specimen of ileitis itself was free of any evident invasion by the tuberculous process and was nonspecific granulomatous in nature by histologic study.

Syphilis and actinomycosis, two specific infections that may involve the intestine have been excluded as an etiologic agency because of the fact that serologically syphilis and histologically actinomycosis are easily differentiated and excepted.

#### SPECIFIC AGENTS

Many authors have found various bacteria in cultures of tissues or in peritoneal exudates and have implied that these bacteria were the etiologic agents of the disease. This occurred many times in the earlier publications after the disease was first described. recent literature has made no

such claims and no such confutations. Erb and Farmer in one case found in stained sections of the involved gut gram negative bacilli apparently of the B. coli group. By culture they obtained a similar organism from the ileal mucosa, mesenteric lymph glands, liver and bile. In their comment on the finding Ravdin and Johnston point out that these tissue cultures were taken eight hours after death and that no specific agglutinations against the patient's blood had been essayed. Ross has found similar bacteria in one case. Mixer noted frequent mixed cultures in the chronic cases, but regained in one case an anaerobic streptococcus from the peritoneal fluid from the mesenteric lymph nodes and from the deep surface of the intestinal ulcerations. Peters found the usual enterococcus of intestinal type in the cultures of superficial tissue but no bacteria in the sections of deeper tissue. Jackman, in 2 cases of acute ileitis, was not able to demonstrate bacteria in the deeper layers of the resected specimens. Others have found gram positive cocci in the submucosa (Konjetzny), enterococci in stool cultures (Fischer and Lurmann), aerobic bacteria aerogenes (Hallgren and Hallgren). Maier found streptococcus viridans in two of his cases.

These various observations are of interest but fail to give consistent findings which would regularly implicate a single causative bacterial agency. Numerous attempts by staining and by aerobic and anaerobic cultural methods and by animal inoculations have similarly failed to discover a singular offending agency.

A most serious attempt to investigate the bacterial etiology of the disease was made by Pumphrey at the Mayo Clinic. Thirteen cases of granuloma of the bowel were studied 10 of which were typical regional ileitis. Cultures were taken by sterile technique in the operating room from resected specimens and regional lymph nodes. The specimens were macerated and planted in media such as blood agar, eosin methylene blue plates, dextrose brain broth, chick mash overlaid with paraffin and agar slants. Many

organisms both positive and negative were recovered, none of which could be said to be predominant throughout the series. In no instance was he able to isolate an organism of the dysentery group. None of these bacterial cultures were consistently agglutinated by the serum of the patient from whom the culture had been obtained nor by the serum of other patients suffering from the same disease. Cultures from the throats of 2 patients with the disease yielded streptococcus viridans, the injection of these cultures into rabbits failed to reproduce the disease. Exhaustive studies for tubercle bacilli and for spirochetes were similarly negative.

Johnes disease in cattle has many resemblances to human ileitis in its anatomic configuration except that the former is regularly demonstrated to be associated with a specific micro organism the *Mycobacterium johnes*. Williams injected johnes subcutaneously into patients with regional ileitis and also into normal controls without establishing any causal relationship between ileitis and Johnes disease.

### PROTOZOA

The entamoeba histolytica was implicated by Cori and Bocch on the basis of one case finding. However, the typical pathologic lesions of amebiasis were absent from other segments of the large bowel as is characteristic of amoebic infestation. No other authors in the literature have made similar claims. In other instances *Trichocephalus dispar*, *Oxyuris vermicularis*, *Giardia lamblia* and other parasites have been found in the stool examinations but never sufficiently consistently as to imply that their presence was the cause of ileitis.

Foreign bodies which play so large a role in granulomata of the intestinal tract such as fish bones, metal ligatures, food residues, pits, fragments of bone, are absent from cases of true regional ileitis or jejunitis. Large giant cells are regularly found in histologic sections of ileitis these giant cells are not tuberculous or syphilitic in origin. They

are best explained on the basis of foreign body inclusions but the extraneous substances that give rise to such giant cells are not gross or material in substance. They are probably in some instances due to lecopodium spores from dustings of surgical gloves at a previous surgical intervention or are due to minute fragments of food detritus collected and secondarily caught in the ulcerated pockets of the mucosa.

### BACILLARY DYSENTERY

Felsen claims a common source of origin for a large percentage of the cases of distal ileitis, chronic ulcerative colitis and nonspecific granulomata. This concomitant agent is in his belief the *Shigella* dysenteriae. In a follow up of 400 cases of acute bacillary dysentery he found 29 cases of acute distal ileitis, 22 cases of chronic distal ileitis, 18 cases of combined chronic ileocolitis and 84 cases of chronic ulcerative colitis. His study includes a survey of the late end results of the epidemic of bacillary dysentery which occurred at Jersey City in September 1933. He gives 84 cases with high serum agglutinations against *Shigella* dysenteriae and eight with positive stool cultures. He describes particularly an acute dysenteric appendicular form with involvement of the small bowel and various types of mixed forms involving terminal ileum and colon. His classification and acceptance of dysentery is based upon agglutination titers of 1:100 and over except for the Sonne-Duval type in which he accepts 1:50 as significant. Positive stool cultures were obtained in 10 of 70 cases examined.

The findings and implications of Felsen so often quoted in the literature may be variously criticized. The agglutination in the serum of patients against recovered organisms and against type laboratory cultures are today open to the criticism of nonspecificity. Too many cross and nonspecific agglutinations occur to allow definite laboratory and clinical conclusions. The small number of positive stool cultures (ten) recovered after an epidemic of acute dysentery is not significant for ileitis unless it can be cor-

organisms, gram positive and gram negative were recovered, none of which could be said to be predominant throughout the series. In no instance was he able to isolate an organism of the dysentery group. None of these bacterial cultures were consistently agglutinated by the serum of the patient from whom the culture had been obtained nor by the serum of other patients suffering from the same disease. Cultures from the throats of 2 patients with the disease yielded streptococcus viridans. The injection of these cultures into rabbits failed to reproduce the disease. Exhaustive studies for tubercle bacilli and for spirochetes were similarly negative.

Johns's disease in cattle has many resemblances to human ileitis in its anatomic configuration, except that the former is regularly demonstrated to be associated with a specific micro organism the *Mycobacterium johni*. Williams injected johni subcutaneously into patients with regional ileitis and also into normal controls without establishing any causal relationship between ileitis and Johns's disease.

### PROTOZOA

The entamoeba histolytica was implicated by Corr and Bocck on the basis of one case finding. However, the typical pathologic lesions of amebiasis were absent from other segments of the large bowel as is characteristic of amoebic infestation. No other authors in the literature have made similar claims. In other instances *Trichocephalus dispar*, *Oxyuris vermicularis*, *Giardia lamblia* and other parasites have been found in the stool examinations but never sufficiently consistently as to imply that their presence was the cause of ileitis.

Foreign bodies which play so large a role in granulomata of the intestinal tract, such as fish bones, metal ligatures, food residues, pits, fragments of bone, are absent from cases of true regional ileitis or jejunitis. Large giant cells are regularly found in histologic sections of ileitis, these giant cells are not tuberculous or syphilitic in origin. They

are best explained on the basis of foreign body inclusions but the extraneous substances that give rise to such giant cells are not gross or material in substance. They are probably in some instances due to lycopodium spores from dustings of surgical gloves at a previous surgical intervention or are due to minute fragments of food detritus collected and secondarily caught in the ulcerated pockets of the mucosa.

### BACILLARY DYSENTERY

Elsen claims a common source of origin for a large percentage of the cases of distal ileitis, chronic ulcerative colitis and nonspecific granuloma. This concomitant agent is in his belief the *Shigella dysenteriae*. In a follow up of 400 cases of acute bacillary dysentery he found 29 cases of acute distal ileitis, 22 cases of chronic distal ileitis, 18 cases of combined chronic ileocolitis and 84 cases of chronic ulcerative colitis. His study includes a survey of the late end results of the epidemic of bacillary dysentery which occurred at Jersey City in September 1933. He gives 84 cases with high serum agglutinations against *Shigella dysenteriae* and eight with positive stool cultures. He describes particularly an acute dysenteric appendicular form with involvement of the small bowel and various types of mixed forms involving terminal ileum and colon. His classification and acceptance of dysentery is based upon agglutination titers of 1:100 and over except for the Sonne-Daval type in which he accepts 1:50 as significant. Positive stool cultures were obtained in 10 of 70 cases examined.

The findings and implications of Elsen, so often quoted in the literature may be variously criticized. The agglutination in the serum of patients, against recovered organisms and against type laboratory cultures are today open to the criticism of nonspecificity. Too many cross and nonspecific agglutinations occur to allow definite laboratory and clinical conclusions. The small number of positive stool cultures (ten) recovered after an epidemic of acute dysentery is not significant for ileitis unless it can be cor-

robortated in cases of regional ileitis by successive investigations, in a sufficient number of cases, over a sufficient length of study. Bisgard and Henski recovered dysentery bacillus from the urine and right kidney in a single case of regional ileitis. The identity of the organisms was established by fermentation and agglutinations against stock sera.

In our own series of almost 500 cases, and in the experience with the additional ward cases at the Mount Sinai Hospital over eighteen years, I can recall only 2 cases with a transient positive culture for the Flexner type of dysentery and only an exceptional case with high agglutinations against similar bacteria. In only one instance an agglutination of 1:320 against Hiss was reached.

While one may accept the comment of Felsen that a type of disease similar to acute distal ileitis may occur during epidemics of bacillary dysentery and that occasional very exceptional instances of positive cultures or high titer serum agglutination may be observed, the imputation that regional ileitis or ileo jejunitis is regularly caused by the Shigella dysenteriae has not been substantiated and is not acceptable to most authors other than himself.

#### VIRUSES

The virus of lymphopathia venereum has been suggested as the cause of ileitis. Stafford in 10 cases of lymphopathia venereum, found 3 suffering from a form of segmental colitis, in no case was the small bowel involved. Because of the cicatrizing tendencies and the possible similarity to cicatrization found in the rectum in lymphopathia venereum, Koster and his associates carried out the Frei test in 4 cases of resected ileitis and in 2 others of ileitis not resected with negative results. Rodniche, Kusner and Palmer considered that the two diseases resembled each other in the primary lymphatic involvement and later development of severe stenosing lesions. However, in 4 cases of ileitis, the Frei test was negative with mouse and human

antigens. The serum was also found negative for neutralizing antibodies against the virus. Similar negative results were found by Friedl Meyer. However, Lombart and Mavern described a case of lymphopathia venereum with multiple strictures of the ileum as well as a rectal stricture. The terminal ileum and the cecum were marked by chronic like beading, the lesion extending down to the submucosa and even to the muscularis. Ocular manifestations were also present, confirmatory of the virus type of infection. Coutts and Oporo in a study of several cases of lymphopathia venereum were able to demonstrate only one instance with ileal involvement. In this person there was an extensive proctitis and an ileitis involving a large part of the terminal ileum.

Lakely and Lasa report in detail, in a Nicolas being the obvious perineal and vulval lesions of lymphogranuloma, an involvement of the small intestine, lower and mid ileum by a granulomatous infiltration having all the histologic characteristics of ileitis or of nonspecific granuloma. Five lesions were found in the upper and more proximal part of the ileum the lowermost being six inches from the ileocecal valve. The wall of the bowel was thickened, the lumen dilated, the mucosal surface was granular and showed numerous shallow ulcers.

However from this exceptional case of lymphopathia venereum with involvement of the ileum, no conclusions can be drawn regarding ileitis as a virus disease at least not as the identical virus of the Nicolas Favre disease. My own attempts at obtaining a positive Prier test in ileitis have been unsuccessful. The frequency of recurrences in ileitis even after extensive resections and the inability to demonstrate a specific bacterial agency allows one to maintain an open mind on the possibility of a virus with a predilection for the ileal mucosa. The fact that two observers did find involvement of the ileum in lymphopathia venereum suggests that a similar though not the identical virus may be considered as a possible true etiologic agency in ileitis.



Allergy has been suggested by Tallroth as the etiologic agent, he described two case histories of "ileitis allergica." The finding of many eosinophils in the histologic sections was repeatedly stressed and obviously constitutes the basis for his suggestion. The vascular injury was also typical of an allergic granulomatous process. He regards the pathologic process in ileitis as a local anaphylactic reaction in the nature of an Arthus phenomenon.

### SARCOIDOSIS

The histologic pathology of Boeck's sarcoid and of ileitis is almost identical. Homans and Hiss stressed the similarity in the lymph gland pathology in the two diseases. Longcope and Picison in their thorough review of sarcoidosis make no reference to intestinal lesions. Williams and Nickerson review 4 cases of sarcoidosis but without any ileal or intestinal or colonic involvement. Snapper reviews the pathology, the clinical incidence and the histology of the two diseases terming them pseudo tuberculosis in nature. He differentiates them sharply, however, in their respective clinical manifestations and considers them unrelated in all respects.

More recently, isolated sarcoidosis of the small intestine simulating nonspecific ileojejunitis has been described. This is the first instance where the real differential diagnosis of ileitis from sarcoidosis involving the small bowel has been verifiably noted though in the literature small intestine involvement had been occasionally suspected. In this paper by Watson and his associates two case histories are detailed which require analysis. The radiographic picture resembled ileitis or ileojejunitis. Resected specimens had many of the characteristics similar to those of ileitis except for evidence of polypoid. The histologic picture was almost identical. Hadfield in studying 20 cases of ileitis had found 13 in which epithelioid cell tubercles were present and cautiously suggested the identity of ileitis and sarcoidosis.

Morland has just published an article entitled, "A Case of Sarcoidosis of the Lung with Regional Ileitis." A filling defect of the cecum was observed radiographically. At operation, resection of the terminal ileum was performed. Typical changes of a "sarcoid" nature were histologically observed in the removed mesenteric lymph nodes. This case is not convincing as one of ileitis. It could very well however, be a good example of involvement of the ileum and colon in sarcoidosis.

The generalized picture of sarcoidosis in all its clinical manifestations, including skin and pulmonary lesions and the absence of fistulas, string sign and rectal complications have convinced most authors that the two diseases have little relationship. The most that can be said is that sarcoidosis may occasionally show involvement of the small bowel. Bernstein, Konzelman and Sidlick report a case of Bocck's sarcoid with true involvement of mucosa of the ileum marked by thickening of the wall with ulcerations measuring 2 x 4 cm. In the case of Lencartovicz and Rothfeld the gastrointestinal tract was diffusely involved. Homans and Hass stated that some observers have called ileitis "sarcoid of the small intestine."

#### TRAUMA

Moek (1931) in discussing infective granulomas states that "no case of granuloma the direct result of an accident such as from an automobile or a fall was ever reported." He makes an exception of course, for granulomata resulting from tears of the mesentery from direct violence to the abdomen, foreign bodies, splinters of wood, strangulated hernias and residual suture material. However as early as 1936 Reichert and Mithes cite the case of a man who suffered a severe blow to the abdomen from the steering wheel of his automobile. Two weeks later he developed severe abdominal pain with vomiting and diarrhea. At resection of the ileum revealed a granulomatous lesion twelve inches in length which was proximal to and not occupying the usual site for regional or terminal ileitis.

Two cases of regional enteritis following external trauma have been reported from the Mayo Clinic [176]. In one case, three months prior, while the patient had been horse back riding, his mount had fallen on him delivering a severe crushing blow to the abdomen. On exploratory operation some weeks later, an obstructing lesion was found about seven feet from the ileocecal valve, relief followed a short encircling procedure. Later the lesion was resected.

The second case involves a farmer who four months previously had been caught between two heavy farm implements, sustaining a severe crushing injury to the abdomen. At operation, several loops of ileum were bound together by inflammatory adhesions and were fixed to one loop of lower jejunum which was definitely obstructed. The usual edema of the mesentery and enlarged lymph nodes and fistulous tracts so characteristic of ileitis were absent. These cases do not resemble ileitis in anatomic distribution or in symptomatology. They represent true direct trauma to the small intestine with resultant granulomata and obstruction. Nominally they are instances of ileitis of traumatic origin.

Ken Kate in 1936 very early reported two cases of regional enteritis following severe trauma. In one case, a rupture of the ileum was found at operation, though nothing that resembled clinical ileitis. In another case, a characteristic lesion with a fistula from the ileum to the bladder was observed at operation.

Spellberg and Gray cite the case history of a soldier motor cyclist who ran into the end of a stalled truck. Abdominal pain and vomiting supervened within thirty days. X-ray revealed a typical jejunitis beginning just below the fossa of Treitz. At operation the jejunum resembled a garden hose being indurated and edematous, with accompanying enlarged lymph nodes. Resection of the lesion was eventually required.

In my own experience in spite of an initial skepticism, there seems to be sufficient basis for associating trauma and ileitis in the relationship of cause and effect (Crohn

[62, 63]) This conclusion is based upon experience with the following six cases briefly abstracted

*Case 1 S. I.* A man 39 years of age in an automobile accident was violently thrown forward against the steering wheel of his car sustaining severe abdominal trauma. Abdominal cramp and diarrhea supervened almost immediately though the man had previously always been well. Operation performed less than two weeks later because of the severity of the complaints revealed a typical terminal ileitis cured by resection. The patient received a large award from the courts. This was my first experience of that type as an expert I testified against the likelihood of trauma creating such a lesion as ileitis though unable to deny the probable aggravation of a previously latent lesion. My viewpoint was changed by the succeeding experiences.

*Case 2 J. 4.* A boy of 17 years of age always previously well was thrown violently against the wheel of a bus as it was involved in a collision striking his abdomen on the forward seat. He immediately developed hematemesis and symptoms of a traumatic duodenal ulcer. A gastric enterotomy was performed a few weeks later the stomach functioned badly. At reoperation a typical regional ileitis was noted and resected. It seems reasonable to deduce that both the duodenal ulcer and the ileitis which involved the terminal loops of the small bowel were traumatic in origin.

*Case 3 M. M.* A 20 year old man suffered at 3 years of age a profound trauma when struck by a moving truck. The child immediately developed an abscess in the right lower quadrant of the abdomen requiring several operations and characterized by an obstinate fecal fistula to the abdominal wall. When 8 years of age the previously closed fistula reopened spontaneously and remained open for several months. When 12 years of age the fistula again reopened but closed without surgical intervention. The same sequence of events took place four years ago. For the last ten months the patient has presented the typical symptom of regional enteritis with diarrhea, fever and abdominal pain. Two fecal fistulas present themselves in the anterior abdominal wall. Radiographically the lower ileum is distorted in its mucous membrane pattern the mucosa of the terminal ileum is normal.

*Case 4 H. 4.* A woman of 50 little a heavy man and immediately felt a pain in the right lower quadrant of the abdomen. She suffered no diarrhea or fever. Seven weeks later on physical examination a tender mass layed in the right lower quadrant of the abdomen. Radiographically a clear cut picture of ileitis was seen involving twelve to eighteen inches of the terminal ileum with several short fistulous tracts to the cecum and adjoining loop of small bowel. The patient was observed for two years during which time successive x-ray showed complete and spontaneous resolution. The mass in the deep abdomen is appeared under observation.

The lifting of the weight ruled out the cause of the ileitis was most probably a factor in bringing to light a latent and a symptomatic regional enteritis.

Two cases of regional enteritis following external trauma have been reported from the Mayo Clinic [176]. In one case, three months prior, while the patient had been horse back riding, his mount had fallen on him delivering a severe crushing blow to the abdomen. On exploratory operation some weeks later, an obstructing lesion was found about seven feet from the ileocecal valve, relief followed a short circuiting procedure. Later the lesion was resected.

The second case involves a farmer who four months previously had been caught between two heavy farm implements, sustaining a severe crushing injury to the abdomen. At operation, several loops of ileum were bound together by inflammatory adhesions and were fixed to one loop of lower jejunum which was definitely obstructed. The usual edema of the mesentery and enlarged lymph nodes and fistulous tracts so characteristic of ileitis were absent. These cases do not resemble ileitis in anatomic distribution or in symptomatology. They represent true direct trauma to the small intestine with resultant granulomata and obstruction. Nominally they are instances of ileitis of traumatic origin.

Ken Kate in 1936 very early reported two cases of regional enteritis following severe trauma. In one case, a rupture of the ileum was found at operation, though nothing that resembled clinical ileitis. In another case, a characteristic lesion with a fistula from the ileum to the bladder was observed at operation.

Spellberg and Gray cite the case history of a soldier motor cyclist who ran into the end of a stalled truck. Abdominal pain and vomiting supervened within thirty days. X-ray revealed a typical jejunitis beginning just below the fossa of Treitz. At operation the jejunum resembled a garden hose, being indurated and edematous, with accompanying enlarged lymph nodes. Resection of the lesion was eventually required.

In my own experience in spite of an initial skepticism, there seems to be sufficient basis for associating trauma and ileitis, in the relationship of cause and effect (Crohn

## FAMILIAL INCIDENCE OF ILEITIS AS A POSSIBLE ETIOLOGIC FACTOR

Ulcerative colitis non-specific, a sister disease rarely occurs in more than one member of a family. My own experience, and medical literature in general, indicate that fact. On the other hand, regional ileitis or enteritis occurs in multiple instances in intimately blood related members of a family sufficiently often to call for attention because herein may lie a clue, if not to etiology at least to familial predisposition. My notes on eleven such striking examples is of such interest as to warrant a closer analysis.

In 1934 the first instance of ileitis occurring in siblings was published. A boy of 14 (1 rohn [38]) who represented historically a very early clinical history of ileitis was resected, and the diagnosis firmly established. Two years later, his sister many years older (32 years of age) complained of abdominal pain and vomiting and developed signs of intestinal obstruction. At operation a typical ileitis involving the terminal eight inches of small intestine was resected. The anatomic situation and pathologic attributes of the lesions were identical in both siblings.

In another pair of siblings sister and brother the same identical observations were made. In the sister the diagnosis being confirmed by operation. We have notes of two brothers both suffering with ileitis, one in the acute form slowly subsiding to spontaneous recovery, one going on to the chronic form requiring resection.

In two brothers, one had ileitis resected the second brother was a case of typical ulcerative colitis. Mothers and daughters constitute a group which was seen three times, all the cases being confirmed. In another instance the mother had ileitis the daughter ulcerative colitis. In a fourth instance the child had ileitis and the mother and two maternal aunt of the child had ulcerative colitis. We have

*Case 5 M I* A man (f 2) was while in the Armed Service severely stunned by the explosion of a bomb on board a U S destroyer. Upon recovery he began to complain of abdominal pain and diarrhea. The radiographic picture upon his discharge was typical of ileitis. Interestingly he was able to continue in the Armed Services and obtained an honorable discharge. Under observation the symptom and radiographic evidence of ileitis has subsided and his case is considered one of spontaneous cure.

*Case 6* A man in an auto accident was thrown violently against the steering wheel of his car with severe bruises to his abdominal wall. Two months after the injury he developed diarrhea abdominal pain and lost thirty pounds in one year. A resection of the terminal ileum revealed a typical regional ileitis. He was well after the operation for three years when he again suffered an accident this time being hit by a street car and thrown violently backward. Immediately the diarrhea recurred. A recurrence of the ileitis in the loop of ileum proximal to the previous anastomosis was suspected though not proven.

Ileitis is a disease of great chronicity, often latent for years. In the analysis of case histories in general it is not unusual to read of mild almost negligible occasional complaints for five, ten or twenty years, before the eventual diagnosis is established. Nor is diarrhea a constant clinical symptom in ileitis or enteritis. The thought seems therefore justifiable that in these instances where trauma has been related to ileitis actually one is dealing with an aggravation of a pre-existing latent ileitis. However the cases cited from the Mayo Clinic and the first three cases in my personal experience seemed sufficiently well checked and proven to allow the inference that the trauma preceded and caused the intestinal lesion. In the other cases the aggravation of a lesion by the trauma seems justifiable.

Traumatic enteritis in its clinical course is identical with ileitis as we know it in its well recognized forms. Abdominal mass, fistula formation diarrhea fever are all present. Segments other than those of the terminal ileum are more likely to be affected by the indiscriminate application of the trauma.

Spontaneous resolution was twice observed in incidence that would seem higher than in the control nontraumatic cases.

## FAMILIAL INCIDENCE OF ILEITIS AS A POSSIBLE ETIOLOGIC FACTOR

Ulcerative colitis, nonspecific, a sister disease rarely occurs in more than one member of a family. My own experience and medical literature in general bears out that fact. On the other hand regional ileitis or enteritis occurs in multiple instances in intimately blood related members of a family sufficiently often to call for attention because herein may lie a clue, if not to etiology, at least to familial predisposition. My notes on eleven such striking examples is of such interest as to warrant a closer analysis.

In 1934 the first instance of ileitis occurring in siblings was published. A boy of 14 (Cohn [38]) who represented historically a very early clinical history of ileitis was resected, and the diagnosis firmly established. Two years later, his sister many years older (32 years of age) complained of abdominal pain and vomiting and developed signs of intestinal obstruction. At operation a typical ileitis involving the terminal eight inches of small intestine was resected. The anatomic situation and pathologic attributes of the lesions were identical in both siblings.

In another pair of siblings, sister and brother, the same identical observations were made; in the sister the diagnosis being confirmed by operation. We have notes of two brothers both suffering with ileitis, one in the acute form slowly subsiding to spontaneous recovery, one coming on to the chronic form requiring resection.

In two brothers, one had ileitis resected, the second brother was a case of typical ulcerative colitis. Mothers and daughters constitute a group which was seen three times, all the cases being confirmed. In another instance the mother had ileitis, the daughter ulcerative colitis. In a fourth instance the child had ileitis and the mother and two maternal uncles of the child had ulcerative colitis. We have



again noted an uncle suffering with ileitis, his niece with segmental or right sided colitis




Fig 1—Familial Incidence in Four Siblings (A) Ulcerative colitis (B) segmental colitis (right sided) (C) regional ileitis (D) duodenal ulcer

Further, three family groups afford a striking picture. In the first, a young woman, her brother and her half brother, have ileitis; the father is suffering from ulcerative colitis. In the second family group we note ileitis, seg-

mental colitis, and ulcerative colitis in three adult siblings (fig. 1). In the third family group a child and her two aunts by consanguinity, have all had ileitis confirmed by operation. Bockus refers to two cousins suffering with regional ileitis with extension into the colon, ileocolitis both resected and well. They had previously lived together in the same house. Brown and Schueffly have reported regional enteritis affecting three siblings, two sisters and one brother. In two of the siblings the colon was involved as well as the terminal ileum. Very recently a case of regional ileitis in a father and son has also been described [135].

The significance of these findings remains conjectural; they are too interesting, however, to be dismissed without consideration. If a proven bacterial or viral agency could be incriminated as the cause of ileitis or of ulcerative colitis one could understand contact infection, such as occurs in epidemics of bacillary dysentery or in amoebiasis, by finger transmission or by insects particularly flies. Or one could conceive of a common vector. Since such a specific etiologic agent is not known, it seems more logical to fall back upon at least a familial predisposition to the unrecognized infective agency such a factor predisposing to ileitis to ulcerative colitis, or to both diseases. The mixed cases of ileocolitis seen in this series only again draw attention to the common liability of small and large intestine to the offending agency.

#### THEORIES REGARDING THE PATHOLOGIC MECHANISM IN THE CREATION OF ILEITIS

Most of the hypothetic considerations regarding the actual location of the lesion, and the possible direct physical factors in initiating the granuloma have centered about the terminal ileum and the ileocecal valve. This viewpoint disregards the fact that the lesion may occur at any level of the small intestine and also that recurrences of ileitis

occur far from the original site of inflammation and from the valve of Bauhin.

Naturally the first suspicion was directed at the appendix as a causative factor. Many writers have suggested that a fibrinoplastic process beginning in the appendix extended to the ileum, thus constituting terminal ileitis (Lowen). Fischer made the same claim but this time through the mediation of the mesenteric lymph nodes. Ravdin and Rhoads again pointed out the similar pathologic changes in fibrioplastic appendicitis, and the chronic stage of regional ileitis. Homans and Hays, and Erb and Farmer, suggested primary appendiceal disease as an etiologic factor.

In refutation of this idea, it should be noted that the appendix had previously been removed in over 25 per cent of cases of ileitis. The life course of the disease is in no way altered or stayed. The resected appendix, apart from showing slight chronic changes of the usual type seen in adult life is free of typical granulomatous changes such as characterize granulomatous ileitis. I know of only two cases where acute phlegmonous appendicitis occurred some years subsequent to a short excising operation for ileitis. Nor is there reason to expect the appendix to be involved in ileitis since the appendix originates from the base of the cecum, beyond the ileocecal valve and is separated anatomically from the ileum by the ileocecal valve, which is such an efficient barrier in preventing the spread of the ileitis to the cecum. Nor does acute appendicitis occur more frequently in the combined cases with cecal involvement. Histologically the appendix in the resected specimens of combined ileocolitis does not participate in the peculiar and characteristic granulomatous inflammation.

#### VASCULAR CHANGES

The analogy of ileitis with intestinal granulomatosis might well suggest that the similar pathologic picture in both diseases result from an identical impairment of the blood supply. Those authors who have written on the subject of

granulomata of the intestinal tract have emphasized volvulus, intussusception, strangulated hernia, and traumatic rents as direct causative agents leading to granulomatous mass formations. Ginzburg and Oppenheimer particularly stress the reduced vascularization that results in mucosal necrosis ulceration and secondary hyperplastic granulomatous changes. Barbour and Stokes mentioned the possibility of a chronic recurrent intussusception. Bockus and Lee discussed the peculiar blood supply of the terminal ileum and its mesentery, suggesting that the terminal branch of the ileocolic artery might be compromised by undergoing rotation. Bell, however, in experimental studies could not reproduce granulomata by interference with the blood supply.

#### THE THEORY OF LYMPHATIC BLOCK

Of much greater interest is the question of the relationship of the mesenteric lymph glands and the lymphatics of the intestinal wall to granulomatous ileitis. The constant finding of enlarged lymph nodes in the mesentery of the affected loops of intestine, while easily explained on the basis of secondary infection, has caused some writers to advance the hypothesis that the primary lesion may be resident in the lymphatic tissues, the organic changes in the wall of the gut being secondary thereto. Holman, in discussing a paper by Bell in 1934, asks whether the lymphatic hyperplasia is an expression of a chronic infection originating in the bowel or is it possibly the primary lesion with subsequent fibrosis and secondary constriction of the intestinal canal? Two years later, Reichert and Mathes published a paper on 'Experimental Lymphedema of the Intestinal Tract and its Relation to Regional Enteritis'. The dominant features of the intestinal lesion in ileitis resembled a low grade infection with a concomitant chronic lymphatic obstruction and edema. They observed the pathologic resemblance to elephantiasis as seen in the extremities of the body. In animal experiments, using a fine needle, sclerosing substances were injected into the

occur far from the original site of inflammation and from the valve of Bauhin.

Naturally the first suspicion was directed at the appendix as a causative factor. Many writers have suggested that a fibrinoplastic process beginning in the appendix extended to the ileum, thus constituting terminal ileitis (Lowen). Fischer made the same claim but this time through the mediation of the mesenteric lymph nodes. Radin and Rhoads again pointed out the similar pathologic changes in fibrioplastic appendicitis, and the chronic stage of regional ileitis. Homans and Hass, and Eib and Farmer, suggested primary appendiceal disease as an etiologic factor.

In refutation of this idea, it should be noted that the appendix had previously been removed in over 25 per cent of cases of ileitis. The life course of the disease is in no way altered or stayed. The resected appendix, apart from showing slight chronic changes of the usual type seen in adult life, is free of typical granulomatous changes such as characterize granulomata and ileitis. I know of only two cases where acute phlegmonous appendicitis occurred some years subsequent to a short enteric operation for ileitis. Nor is there reason to expect the appendix to be involved in ileitis since the appendix originates from the base of the cecum beyond the ileocecal valve and is separated anatomically from the ileum by the ileocecal valve, which is such an efficient barrier in preventing the spread of the ileitis to the cecum. Nor does acute appendicitis occur more frequently in the combined cases with cecal involvement. Histologically the appendix in the resected specimens of combined ileocolitis does not participate in the peculiar and characteristic granulomatous inflammation.

#### VASCULAR CHANGES

The analogy of ileitis with intestinal granulomata might well suggest that the similar pathologic picture in both diseases result from an identical impairment of the blood supply. Those authors who have written on the subject of

terial absorption. Studies on the ileo cecal junction (ileo cecus) by Barger and his associates demonstrated longitudinal muscle fibers entering the labia of the valve for a considerable distance and meeting at the crest. Gelatin solutions injected at the cecal mucosa were blocked at the tip of the labia, similar injections made on the ileal side in 15 cadavers were also blocked at the valve. Obviously there is a lymphatic blockage at the tip of the labia of the ileo cecal junction.

*The possible relationship of acute mesenteric lymphadenitis to acute and chronic regional enteritis* presents an interesting thought. Jackson suggested the kinship of the two diseases. However, in 15 cases he found only a mild hyperemia of the terminal 3 to 4 inches of the terminal ileum, he has no record of any case of ileitis originating in, or from acute mesenteric lymphadenitis. Rockey noted thickening of the ileum in children with acute mesenteric lymphadenitis in 4 cases. All were valid cases of acute ileitis with marked lymphadenopathy, and this in September 1932 before the first publication on ileitis.

On the other hand, Eliason and Johnson state that they know of no case of acute ileitis originating in acute mesenteric lymphadenitis. In the 15 cases of acute ileitis observed by them, and operated upon, they found no intraperitoneal picture that resembled acute mesenteric lymphadenitis either as regards abundant free peritoneal fluid, or massive disseminated lymph gland enlargement.

In 1932 Strombeck published his observations on 40 cases of acute nontuberculous mesenteric lymphadenitis controlled by operation. In 22 instances, he noted the occurrence of acute terminal ileitis which he presumes was the point of origin of the infection. Ten to 20 to 50 cm of terminal ileum proximal to the ileo cecal valve were involved, the intensity of the affection being at the valve and diminishing upward. The Peyer's patches were large and 'grumpy boat shaped swellings'. The greatest frequency occurred between the ages of 5 and 15 years corresponding to the

mesenteric lymphatics and serosal lymphatics of the wall of the intestine

The materials used for sclerosing purposes included crystalline silica mesh, bismuth oxychloride, indelible lead or sodium morrhuate. In some animals, a 24 hour broth culture of *B. coli* was given intravenously one to three hours before the lymphatic injection, those bacterial injections occasionally being repeated at intervals subsequent to the acute experiment. When the animal was sacrificed, the treated segment of bowel felt thickened, some of the mesenteric lymphatics were dilated, some sclerosed. Free peritoneal fluid was not present. The thickened bowel wall showed inflammation and thickening of the serosa, edema of the circular and longitudinal muscle fibers, the submucosa was swollen and infiltrated with leukocytes, the mucosa swollen but not ulcerated. Chronic pathologic changes persisted without subsidence for months after even a single sclerosing injection. The changes were seen in all specimens (12) examined, but were most marked in those experiments in which bacterial suspensions were intravenously injected coincidentally with the lymphatic block. The intestinal lymphedema was found to persist as long as ten months without any evidence of subsidence.

It is interesting to note that Poppe, using similar method of lymphatic sclerosing injections, claims to have produced ulcerative colitis in dogs by obliteration of intestinal lymphatics. However, Sinaiko and Necheles, repeating their work, not only failed to produce ulcerating lesions in the colon of dogs, but point out, in contradistinction to the results of Reichert and Mathes that the lesion in the colon showed a tendency to rapid disappearance and no tendency to chronicity.

It had already been suggested (Probstein and Gruenfeld) that normally the flow of the intestinal content is retarded at the ileocecal valve, the lymphatic tissue in the mucosa and submucosa is here more abundant than in the segments of the bowel, a factor which enhances the likelihood of bac

is well represented. Childhood and old age are scantily represented. The oldest case was 71 years at observation, the youngest case was 9 months in one instance, and 2½ years in another instance. The 9 months old case was confirmed by operation and by x-ray studies.

These figures agree well with those of other authors who have collected large series of cases. Thus Strombeck collected from the literature 91 cases, the greatest incidence (40 cases) fell between 20-30 years of age, namely 44 per cent, agreeing almost identically with the figures in this series. Similarly, Ravdin and Johnston in a collection of 393 cases from the literature give the age incidence as shown in table 2.

TABLE 2.—Age Incidence in Regional Hentis (Ravdin and Johnston)

Age Group	Number of Cases	Percentage
1-10	18	4.7
10-20	68	17.8
20-30	140	36.6
30-40	76	19.9
40-50	40	10.4
50-60	4	6
60-70	13	3.4
70-80	3	0.8
Total	38	

### SEX

Males predominate over females in their susceptibility to this disease. In this present series, there were 124 males and 98 females involved representing a proportion of 56 per cent males to 44 per cent females. These figures again agree with the mass compilation of cases from the literature in which males constituted 56.6 per cent of the cases (Ravdin and Johnston). The figure given in earlier years based on a smaller group of cases (Crohn [59]) of males predominating over females in a proportion of 2:1 is not borne out when large group statistics are invoked.

Schiff garnered the statistics from the literature on 48 cases of hentis in childhood. The ages varied from 1½ years to 16 years as shown in table 3.



greatest frequency of acute ileitis Strombeck distinctly states that in the follow up covering at least four years, he has never seen any such case develop into the picture of a chronic regional ileitis Of 31 cases of nontuberculous mesenteric lymphadenitis observed by Erskine 10 showed some involvement of the ileum in an inflammatory process associated with large hypertrophied lymph nodes in the mesentery He does not relate the disease to true regional ileitis In his opinion the lymph nodes are first involved and then the appendix or the ileum or both by an extension of the process

The observations of Strombeck more or less dispose of the hypothesis that chronic regional enteritis begins as an acute mesenteric lymphadenitis The largest percentage of cases of ileitis in this series (222 out of 238 cases) were chronic in nature throughout their course The abdominal findings at operation in acute mesenteric lymphadenitis is generalized enlargement of mesenteric lymph nodes, much clear fluid, occasional involvement of terminal ileum In acute ileitis, one observes scant free fluid, edematous engorgement of the terminal ileum and only localized lymph node involvement Perforation of the acutely inflamed ileum with localized walled off peritonitis is not unusual

#### AGE IN REGIONAL ILEITIS

In our series the age distribution at onset of symptoms may be noted in table 1

TABLE 1—*Age Distribution at Onset of Regional Ileitis*

Years	Number of Cases	Incidence (%)
1-10	6	2.7
10-20	34	24.4
20-30	98	44.0
30-40	49	20.0
40-50	9	4.0
50-60	4	1.4
60-70		0
Total	192	

The average age at onset was 27.8 years The maximum incidence falls between 20-30 years, though adolescence

both psychic strain and physical disease as the background for the etiology of regional ileitis is seen in the following case

*H B* Serial No 20 male age 29 years Patient enlisted in the U S Army in 1939 serving in the Philippine Islands He was a prisoner of war to the Japanese from 1942 to 1945 one of the many who participated in the Death March from Corregidor He suffered from bacillary dysentery amebic dysentery, undulant fever malaria pellagra scurvy beriberi and severe malnutrition During all this time he had severe diarrhea In July 1945 the diagnosis of regional ileitis was made by x ray examination the granulomatous process involving the whole of the ileum When seen in 1946 he had recovered weight and strength and was well except for the diarrhea X ray examination demonstrated a normal colon but a severe diffuse regional ileitis The sigmoidoscopy was negative (fig 2)



Fig 2—Regional Ileitis in a Man who Survived the Death March and Three Years in a Japanese Prisoners Concentration Camp Favorable clinical course under medical therapy

All classes of society are affected equally by ileitis, private hospitals and public wards seem equally represented

TABLE 3—*Age Incidence of Ileitis in Childhood (Schiff)*

<i>Age</i>	<i>Number of Cases</i>
1½ years	1 case
2-3 years	4 cases
4-6 years	6 cases
7-9 years	18 cases
10-12 years	2 cases
13-14 years	7 cases
15-16 years	7 cases
Total	43 cases

He emphasized particularly the acute phase of the disease in children and the similarity of the clinical picture to that of acute mesenteric lymphadenitis.

#### SOCIAL STATUS, RACE, RELIGION, NATIONALITY

It is unlikely that any of these social factors play a role in the etiologic incidence of the disease. Boclus says of the personal pattern that very few of these cases are phlegmatic and calm, they are usually sensitive, emotional and excitable. Four of 19 cases in his series were severely psychoneurotic, in 2 of them the disease having been treated as a neurosis. My own observations do not impress me with the fact that the ileitis case is much different than a control group of the population. True, many of them are high strung and anxious, in several instances the patients had been institutionalized for psychoneurosis or anxiety state, the diarrhea having been completely overlooked. The secondary nervous manifestations resulting from prolonged illness, diarrhea, inanition, loss of weight and anemia may simulate a psychoneurosis but only to the uninitiated (Blackburn).

The personality of the ileitis case is more stable than that of the patient suffering from ulcerative colitis or of peptic ulcer. He has less of the psychosomatic factors so evident in both these latter groups. However, the number of cases of ileitis that have been rescued from institutions for the treatment of mental diseases emphasizes not the personality but the end results of the drain of the disease upon the psychic constitution of the sufferer. An interesting instance of

Hindu from British India. Strombeck of Stockholm with 26 cases reported and with an extremely careful analysis of etiology, fails to mention race or social position. In England (Hadfield), 6 out of 22 cases occurred in Jews.

in proportion to population Bockus seems impressed with the failure to encounter regional ileitis in families who have been in the upper economic brackets for one or two generations All races and climes seem represented in this disease since the literature from all over the world contains publications of cases (Africa, India, Sweden, Puerto Rico [8], East Africa [130], South America [26-28]) Bockus again mentions the fact that no cases had been reported from Latin America or from Cuba This deficiency has since been supplied in an article based on experiences in Puerto Rico He also observed paucity of cases from the southern sections of South America In 1944, there were 134 admissions from ileitis in the total U S Army in 1945 there were 145 similar admissions for the entire Army The corresponding rate per 1000 strength for each year is .02 (personal communication) Kiefer found that regional ileitis occurred seventy times in 100,000 registrants at the Lahey Clinic

As regards race, a greater incidence of the disease among the Jewish race was first suggested by Homans and Hass In the series by Mixer, 8 of his 11 patients were of Jewish origin But as these patients were reported from the Beth Israel Hospital at Boston as were mine from the Mount Sinai Hospital in New York, both being institutions with predominant Jewish admissions, this race incidence was not particularly noteworthy In the experience of Bockus, 13 of 21 cases were Jewish *thrice the normal expectancy* of the disease for the sample from which the cases were drawn However in Marshall's series, at the Lahey Clinic, only 3 of 29 instances were Jews Ashley at the Harper Hospital in Detroit found only 8 of 31 cases in his series to be Jews, Brown and Donald 26 per cent Jews (Mayo Clinic) The study comprising 22 cases of ileitis in the records of the Binghamton City Hospital and other hospitals in Broome County New York State include no cases involving Jewish patients (Sneirson and Ryan) Negroes, Orientals Latin Americans are all represented and reported in the literature one of my own cases was a

Hindu from British India Strombeck of Stockholm with 26 cases reported and with an extremely careful analysis of etiology, fails to mention race or social position In England (Hadfield), 6 out of 22 cases occurred in Jews

REGIONAL ILEITIS is a nonspecific granulomatous inflammation of the ileum, involving most frequently the terminal segment or segments of the small bowel. In gross appearance it has definitely characteristics that distinguish it from granulomata elsewhere in the alimentary tract, these specific features being most pathognomonic when the terminal ileum alone is involved. The histologic microcopy of the disease is nonspecific and not to be differentiated from other granulomata, from sarcoid and often only with difficulty from tuberculosis.

In our series of 222 cases, the extent of ileal invasion was noted in our records in 156 instances. The inability to state the amount of anatomic involvement in the remaining cases was due to the fact that they had been previously operated elsewhere or that no surgical corroboration of the radiographic picture had been obtained.

The anatomic distribution in this series may be seen in table 4.

TABLE 4 — *Anatomic Distribution in Regional Ileitis (Fig. 3)*

<i>Distal (inches)</i>	<i>Cases</i>	
4	8	
6	18	Terminal 1 <sup>st</sup> inches 93 cases
8	16	
10	14	
12	5	
15	17	Terminal 2 <sup>d</sup> inches 47 cases
18	15	
24	15	
30	8	
36	8	Extension to the remainder of ileum 6 cases
48	3	
60	4	
84	3	
Total	156	

In one remaining case, the terminal ileum and the upper most ileum were discontinuously involved. The predominance of the lesion limited to the terminal 12 inches of ileum is noteworthy, the next longest group involves not only the terminal segment but extends continuously to cover the distal 24 inches of ileum, the smallest group involves the terminal and all of the upper reaches of the ileum which we arbitrarily indicate as being about seven feet




Fig 3—Regional Ileitis. Anatomical Distribution of Pathologic Lesion (108 Cases). Table represents area from ileo-cecal valve to general region of transition from ileum to jejunum. Majority of cases fall between 8 and 12 inches from the ileo-cecal valve.

in length from the ileo-cecal valve. In the series of cases reported by Blackburn, Hadfield and Hunt, of 22 cases the terminal ileum was involved in 19 instances, the distal ileum in 3 instances. In 6 cases the cecum was involved in an analogous extension of the process. The length of ileum involved varied between 4 and 60 cm with an average of 19 cm ( $7\frac{1}{2}$  inches).

Why the terminal ileum is so often the sole seat of the disease and apparently the oldest and parent lesion in



the process is difficult to state. Most diseases of the small bowel are located at or near the terminal ileum just proximal to the ileo cecal valve. These include benign and malignant new growths, foreign body granulomata, primary and secondary hyperplastic tuberculosis, sarcoid, carcinoid and endometriosis. The normal narrowing of the lumen of the small bowel as it progresses distally is maximum just proximal to the valve of Bauhin, the valve itself constitutes a mechanical device for slowing up the progress of the column of chyme, the lymphatics of the intestinal wall, particularly the mesenteric lymph nodes are in their greatest density at or about the ileo cecal angle. The natural blockage of the lymphatic drainage at the labia of the valves, as shown by Barger et al, localizes an infection proximal to the valve and discourages or prevents in the largest percentage of cases, the transition or extension of the disease to the cecum. Occasional cases are cited where the lesion involves a small segment of ileum proximal to the valve without involving the actual terminal segment of the ileum, but these cases are exceedingly rare.

The gross appearance of ileitis as seen at the operating table or as studied in the resected specimen is that of a brilliantly red congested edematous mass, solid and firm, resembling a rubber hose, or as Dalziel has described it, of "an eel in a state of rigor mortis." An equally apt description is that of Jackson who likened the diseased segment of ileum to "a dead water soaked night crawler." The serosal surface is greatly injected, usually covered by a fibrinous exudate the lumen of the gut may be dilated, or in the end stages reduced to a narrowed cordlike tube. The extent of involvement is manifest by the varying degrees of hyperemia, the engorgement of the vessels, the serosal exudate and by the gross enlargement of the accompanying mesenteric lymph nodes. The hyperplasia of the mesenteric lymph nodes parallels the extent and degree of the inflammatory lesion. Where the lesion ends, usually more or less abruptly the serosa of the proximal bowel

appears normally shiny and smooth, and the obviously enlarged mesenteric nodes are no longer visible.

The mesentery of the involved bowel is boggy, swollen and visibly thickened. Usually one or more loops of small intestine are matted together forming a palpable inflammatory mass. On attempting to separate the loops it will frequently be noted that they are adherent because of fistulous communications, fistulous tracts originating




FIG. 4—Characteristic Lesion in Typical Terminal Ileitis (Regional). Note the diminished lumen which is the basis for the string sign.

in the terminal ileum extending through the mesentery to the adjoining loops of small or of large bowel. Occasionally a slow walled off perforation of the terminal loop of ileum will be observed covered over by a plastic exudate and sealed off by neighboring viscera usually another loop of ileum or the sigmoid. Bockus emphasizes the point that there is a close relationship between the destructive changes in the mucosa and the extent of covering of the ulcers with mesenteric fat. The termination of the mesenteric fat

the process is difficult to state. Most diseases of the small bowel are located at or near the terminal ileum just proximal to the ileocecal valve. These include benign and malignant new growths, foreign body granulomata, primary and secondary hyperplastic tuberculosis, sarcoid, carcinoid and endometriosis. The normal narrowing of the lumen of the small bowel as it progresses distally is maximum just proximal to the valve of Bauhin, the valve itself constitutes a mechanical device for slowing up the progress of the column of chyme, the lymphatics of the intestinal wall, particularly the mesenteric lymph nodes are in their greatest density at or about the ileocecal angle. The natural blockage of the lymphatic drainage at the labia of the valves, as shown by Barger et al, localizes an infection proximal to the valve and discourages or prevents in the largest percentage of cases, the transition or extension of the disease to the cecum. Occasional cases are cited where the lesion involves a small segment of ileum proximal to the valve without involving the actual terminal segment of the ileum, but these cases are exceedingly rare.

The gross appearance of ileitis as seen at the operating table or as studied in the resected specimen is that of a brilliantly red congested edematous mass, solid and firm, resembling a rubber hose, or as Dalziel has described it, of "an eel in a state of rigor mortis." An equally apt description is that of Jackson who likened the diseased segment of ileum to "a dead water soaked night crawler." The serosal surface is greatly injected usually covered by a fibrinous exudate the lumen of the gut may be dilated, or in the end stages reduced to a narrowed cordlike tube. The extent of involvement is manifest by the varying degrees of hyperemia, the engorgement of the vessels, the serosal exudate and by the gross enlargement of the accompanying mesenteric lymph nodes. The hyperplasia of the mesenteric lymph nodes parallels the extent and degree of the inflammatory lesion. Where the lesion ends, usually more or less abruptly, the serosa of the proximal bowel

lumen of the bowel itself is reduced and compromised by the swelling of the walls until the actual lumen is reduced to a cord like stenotic opening. This reduction of the lumen of the bowel in old chronic distal ileitis accounts for the characteristic "string sign" as seen in radiographic films. The process may continue to true stenosis with intestinal obstruction and proximal dilatation of the bowel loops, the stenosis being due to circumferential contraction of old inflammatory scar tissue at the site of the ulcerating lesions.

The mucous membrane has a cobble stone appearance due to the fact that the transverse folds of the mucosa are in part destroyed and are intersected by deeply grooved longitudinal ulcerations that extend through the length of the specimen. These longitudinal ulcerations are deepest along the mesenteric border or attachment. It is in this area along the mesentery that the sinus and fistula tracts emanate. The surface of the mucosa is superficially desquamated the markings are gone. Deep ulcers 0.5 cm. to 1 cm. in diameter are interspersed between areas of proliferated almost polypoid hyperplastic mucosa. True polypoid formation in late cases as is seen in ulcerative colitis is rarely observed. The submucosa is markedly thickened and edematous and congested. The muscularis is thickened and hyperplastic. The serosa its sheen gone is bright red many times its normal thickness in depth and covered by caudite.

The gross lesion in the bowel wall terminates rather abruptly. A sharp line of demarcation is lacking, but the mucosa within one or two inches of the proximal area of involvement will suddenly lose its edema and thickening and shortly resolve into normal mucosa. Unfortunately the inflammatory process has a tendency to extend proximally by kangaroo like jumps. 'skip lesions' of involvement are common proximal to the distal involved segment, these 'skip lesions' being separated by 'skip areas' of normal

on the antimesenteric portion of the serosa corresponds to the line of demarcation between diseased and normal mucosa




Fig 5—Characteristic Arrangement of Lesion in Terminal Ileitis. The most terminal segment of the ileum proximal to the ileo-cecal valve shows the characteristic stenosis of the lumen as the basis for the formation of the string sign

Upon opening the resected specimen one notes the edema and thickening of all the coats of the bowel (figs 4, 5). The

stitutes to a large degree the disappointment and results of surgical intervention

### MICROSCOPIC PATHOLOGY

There are no specific features that characterize or are pathognomonic of ileitis in contrast to granulomata or so called pseudo tuberculous diseases in general. The stained sections show varying degrees of acute subacute and chronic inflammation, predominantly marked by infiltration of polymorphonuclear leukocytes, plasma and round cells and fibroplastic elements. The earliest lesion is apparent in the submucosa where the intensity of the lesion is the greatest (Hadfield), (Schepers). The mucosa is desquamated the architectural structure of the glandular tissue is destroyed and replaced by plasma and round cell infiltration. In the late stages of the disease the inflammatory reaction is more focal in character covered on the mucosal side by nodular thickening reminiscent of tubercles. The enlarged congested skeins of inflamed lymphatics are often apparent even through the thickened serous peritoneal covering.

Microscopically the presence of giant cells is very striking. In the vicinity of the giant cells one often sees groups of pale cellular structures resembling vegetable cells in nature. The giant cells are not specific to ileitis they had been noticed previously by Mosechowitz and Wilensky in their study of intestinal granulomas and were thought by them, and by subsequent writers, to be due to foreign body inclusions vegetable matter or leopodium spores. Homans and Hass thought the giant cells to be a reaction to lipid material. Reichert and Mithes to possible silica or talc inclusions. This vegetable material becomes entrapped in the deep mucosal ulcerations, enters the lymphatics and becomes encapsulated in the process of healing. The attempts of earlier authors to include ileitis as a form of tuberculosis was probably due to the findings of giant cells of the Langerhans type.

mucosa. The extent of the skip may vary from a few inches, or in multiple "skip lesions" to several inches, up to 2½ feet between the parent and the "skip lesion" (fig. 6).

In older phases of the disease, the exudative or inflammatory reaction is replaced by a fibrostenotic process, the mucosa appears atrophic with occasional superficial erosions and islands of papillary or polypoid hyperplasia. The lumen of the gut may be so reduced as barely to admit a large probe.




FIG. 6—Varieties of Skip Lesions and Skipped Area in Ileitis. Figures represent inches.

Pemberton and Brown have made an important contribution in differentiating the lesion in ileitis into two categories: (1) Involvement of a shorter or localized segment of the ileum, a single lesion. (2) a similar process which involves larger segments, usually multiple with a greater tendency to progressive extension. Clinically, this differentiation on a pathologic basis seems amply verifiable. The first or localized group may and does progress to stenotic fibrosis, remains stationary and may lead to intestinal obstruction. They lend themselves favorably to surgery. The second group is scattered, progressive, in the nature of a mucosal or submucosal lesion less well defined anatomically with less involvement of regional lymph nodes, and less amenable to accurate definition during exploratory operation. Thus, the "mucosal" or submucosal type con-

stitutes to a large degree the disappointing end results of surgical intervention

### MICROSCOPIC PATHOLOGY

There are no specific features that characterize or are pathognomonic of ileitis in contrast to granulomata or so called pseudo tuberculous diseases in general. The stained sections show varying degrees of acute, subacute and chronic inflammation, predominantly marked by infiltration of polymorphonuclear leukocytes plasma and round cells and fibroplastic elements. The earliest lesion is apparent in the submucosa where the intensity of the lesion is the greatest (Hadfield) (Schepers). The mucosa is desquamated the architectural structure of the glandular tissue is destroyed and replaced by plasma and round cell infiltration. In the late stages of the disease the inflammatory reaction is more focal in character covered on the mucosal side by nodular thickening, reminiscent of tubercles. The enlarged congested sheins of inflamed lymphatics are often apparent even through the thickened serous peritoneal covering.

Microscopically the presence of giant cells is very striking. In the vicinity of the giant cells one often sees groups of pale cellular structures resembling vegetable cells in nature. The giant-cells are not specific to ileitis they had been noticed previously by Moschcowitz and Wilenski in their study of intestinal granulomas and were thought by them and by subsequent writers to be due to foreign body inclusions vegetable matter or lecopodium spores. Hornans and Hass thought the giant cells to be a reaction to lipid material Reichert and Mathes to possible silica or talc inclusions. This vegetable material becomes entrapped in the deep mucosal ulcerations enters the lymphatics and becomes encapsulated in the process of healing. The attempts of earlier authors to include ileitis as a form of tuberculosis was probably due to the findings of giant cells of the Langerhans type.





Fig 7—Pseudo tubercle with Epithelioid Cells Arranged in a Whirl around Giant Cell in Submucosa (Regional Ileitis)



Fig 8—Lymph Node Showing Pseudo tubercle Formation with Giant Cell Systems

Hadfield has made an intensive histologic study of twenty resected specimens at St. Bartholomew's Hospital in London. He particularly emphasized the early and intense involvement of the submucosa considering that the mucosa was probably secondarily involved in the process by extension and by necrosis due to vascular infiltration. The changes early in the submucosa consist of hyperplasia of the lymphatic tissue and obstructive lymphadenitis. The submucosal




Fig. 9—Longitudinal Section of Skip Lesion in Regional Ileitis. Showing Marked Submucosal Thickening due to Granulomatous Infiltration. Epithelium Completely Altered and Surface Covered with a Purulent Exudate.

hyperplasia and early and extensive involvement of the lymphatics occurs in his experience in 93 per cent of the cases. These early areas of lymphadenoid tissue show clear evidence of cellular activity with a central zone of proliferating reticulum cells surrounded by a narrow zone of lymphocytes (figs. 7 and 8). A specific response occurs. The affected germinal centers become replaced by proliferat-

ing endothelial cells. A Langerhans giant cell can usually be found at the center of the nodule. The follicle now becomes a "giant cell system." Hadfield states that in the several hundred sections he has examined caseation has never been observed. The typical specific reaction, with giant cell inclusions, characterized the sections of lymph nodes found in the mesentery and accompanying the specific lesion in



Fig. 10—Cross Section of Chronic Terminal Ileitis Showing Stenosis of the Lumen Caused by Extensive Granulomatous Infiltration of Submucosa

the gut (figs. 9 and 10). Actually, Hadfield found giant cells in 13 out of 20 cases examined. Necrosis within the granulomatous areas in the ileum or in the lymph nodes rarely occurs; caseation never. In the series of Warren and Sommers, necrosis was observed in 8 of 61 cases (13 per cent). Reference has been made to the case of Frazer and Haggan in which caseating tuberculous mesenteric lymph nodes were observed in the resected specimen though the ileitis proper was nonspecific in origin.

Endoenteritis obliterans and perivascular infiltration with collections of plasma cells were emphasized by Binney. Early involvement of nerve tissue particularly Auerbach's plexus was noted by Barbour and Stokes, though these changes are not specific. They even suggested that a chronic recurring intussusception might initiate the process by impairing the circulation and vitality of the gut wall. Schepers divides the lesion into two components: a primary phase characterized by (a) a stage of edema of the submucosa and the serosa with dilatation of submucosal lymphatics and hyperemia of juxta-muscular adventitial blood vessels, (b) plasma cell infiltration of submucosa and serosa, (c) diffuse fibrosis with disappearance of plasma cells, (d) heralding a secondary phase characterized by ulceration of the mucosa and submucosa which is superimposed on the primary phase with eventual tendencies to late or early perforation, fistulation and granuloma formation.

## 4 Clinical Features

### ONSET

THE USUAL CASE of regional ileitis begins not as an acute and sudden manifestation, its onset is usually hidden in the deep past, with such intervening symptoms as an occasional spell of diarrhea, a progressive malaise, or as nervousness often verging on true psychoneurosis. A rectal suppurative complication may long precede the manifest initiations of diarrhea. Gradually the diarrhea increases in severity and is now accompanied by abdominal pain, occasionally low grade temperature is noted with loss of weight. In the prodromal stages the patient loses little efficiency or time from his work or occupation. The progressive symptoms build up over the course of years to the finally fully developed picture of regional ileitis. Such is the most common picture of the onset. Some unusual forms of onset have been noted in this series such as initial intestinal obstruction, 2 cases, onset with hematemesis and melaena, 2 cases, onset with dysuria, one case, onset with sealed off perforation, one case.

The duration of symptoms before the actual recognition and diagnosis of the disease, in this series may be seen in table 5.

TABLE 5—Duration of Symptoms before Recognition of Regional Ileitis

0-3 months	8 cases
3-6 months	14 cases
6-12 months	15 cases
1-3 years	20 cases
3-5 years	7 cases
5-10 years	13 cases
10-20 years	6 cases
20 years	1 case
Total	183 cases

It will be seen that the largest group of cases falls between one and five years as the duration of the symptoms before the initiation of therapy the next largest group falls between five and twenty years before recognition of the disease. The one with a fifty year unrecognized diagnosis was a Hindu from British India. Blackburn states that the average duration of symptoms before operation was  $31\frac{1}{2}$  months. Clark and Dixon found that the average duration of symptoms before coming under observation was 41 years, the longest duration of symptoms was 25 years.

If one were to speak of prodromata one could mention only the suppurative rectal complications which so often precede by years the onset of manifest symptoms. These rectal or peri rectal abscesses and fistulas occurred in 44 of the 222 cases the real significance lies in the fact that they so often precede the onset of diarrhea. If they do not actually antedate the diarrhea, they are so obvious and the diarrhea so minimal that to most uninitiated observers only the fistula or peri rectal abscess exists. On one occasion a patient who was admitted to the wards for routine hemorrhoidectomy was observed to have a small rectal fistula. An alert resident physician detected a vague history of diarrhea, and asked for radiographic study which disclosed a terminal ileitis.

The one other phenomenon which stigmatizes the onset of ileitis is the scar of a previous appendectomy. This scar was evident in 68 or 30.6 per cent, of our series of cases and was a signal of previous abdominal pain localized in the lower right quadrant of the abdomen. Ravdin and Johnston found in a review of currently published cases in the literature that a previous appendectomy had been performed in 27.1 per cent of the cases. The diarrhea which characterizes most of the cases of ileitis was undoubtedly overlooked the abdomen was opened, an innocent appendix removed and the ileitis not observed.

The onset of the true symptoms of ileitis or enteritis

is characterized, when typical, by the initiation of diarrhea, abdominal pain, low grade fever progressive anemia and loss of weight and eventually by the ability of the clinician to feel an intra abdominal mass. *Diarrhea* was present in 165 of the 222 cases of regional ileitis. The diarrhea is an outstanding feature of the disease, though in number of movements and in intensity it rarely approaches that of ulcerative colitis. The rule that "frequency of fecal movements in an inflammatory intestinal process is in inverse proportion to the distance from the anal sphincter," holds good. Ulcerative proctitis gives usually 10 to 20 stools a day with urgency, tenesmus and the appearance of fresh blood in the stools. Ulcerative colitis in the distal colon may average 5-10 stools a day, right sided or segmental colitis is notable for lessened frequency of defecation, perhaps 3-6 per day. Regional ileitis with all its pathologic process proximal to the ileocecal valve is noted for milder diarrhea, 2 to 4 or 5 movements in twenty four hours.

Each individual movement is usually preceded by a spasm of abdominal pain. The characteristic sequence of events is similar to the classic description of Moynihan regarding duodenal ulcer, where it is "pain food, relief, pain." In ileitis, the sequence is abdominal pain or colic, desire to defecate, the feeling of sudden rumbling of gas, followed by defecation with relief of all subjective abdominal symptoms, so called predefecation pain. This sequence may repeat itself every few hours of the day less at night, it is often initiated by the taking of a meal. The defecations are not urgent are not spurting in nature. The stool is mushy, semi formed or liquid contains some excess of fluid mucus never evident pus and rarely blood (11 instances out of 222). In fact occult bleeding in the stool on examination with guaiac is usually negative. Hurst, however, reports 5 cases of regional ileitis and 2 additional cases previously noted in all of which occult blood was regularly found in the stool. The stool may be slightly acid or alkali

linc or it may be neutral the reaction here has no significance

Either a normal number of daily movements (41 cases) or constipation (18 cases) mark the remainder of the cases in this series. Not infrequently even in a mature case of ileitis no history of diarrhea is obtainable and in some few cases (18 in all) actual constipation was noted. The presence of constipation does not necessarily denote obstruction since only in the late phases of intestinal obstruction is the constipation acute and obdurate. In those cases in which constipation has been present, the remaining symptoms of abdominal pain, colic, a mass and a progressive course have also been negligible, the radiographic study establishes the diagnosis.

*Abdominal pain* was noted in 126 instances in this series. This figure is surprising for pain in the abdomen seems so constant a complaint in the anamnesis that one would expect its presence in a much higher percentage of the cases. The pain is usually mildly colicky in nature, most often situated in the lower abdomen, preferring the right lower quadrant or the suprapubic region in its site. Occasionally the pain is solely in the left lower quadrant and in a few cases covers the whole lower and upper abdominal segments. The pain precedes and is relieved by defecation. When the inflammatory mass becomes adherent to the sigmoid loop of colon and pinned down to that viscus by internal fistulas the pain and the mass is likely to be localized in the left lower rather than the right lower quadrant of the abdomen.

The origin of the pain is in the mildly obstructive narrowed loop of the ileum that segment which radiographically corresponds to the 'string sign'. The attempt of the intestinal fluid column to pass through this congested, edematous and narrowed stretch of intestinal lumen gives rise to the colicky sensation of pain. The absence of pain occurs in those cases in which the inflammatory lesion is diffuse the 'mucosal type' in which the beginning eiec



tricular process is still young, the mucosal pattern is altered by the stenosing string sign, the quill like narrowing has not yet supervened. The higher the process in the ileum, the less likely the stenosis and hence the less likely predefecatory pain. In the late long standing cases where true stenosis supervenes, the pain is of course severe and sharply colicky. Occasionally primary intestinal obstruction characterizes the onset of regional ileitis, or at least its symptomatic phase. Such was the unusual onset in the case quoted by Dickinson and Zimmerman.

*Nausea and vomiting* are rare manifestations and occur only in the later stenotic phase of the disease. Appetite is usually well preserved, this is in contrast to ulcerative colitis, where loss of appetite, absolute anorexia and even vomiting are common manifestations and parallel the intensity of the toxic process. Ulcerative colitis is a much more "toxic" disease than regional ileitis and the interference with appetite and with nourishment is in contrast to the milder manifestations and retained desire for food which is present in ileitis.

*Fever* occurred in 69 cases of the 222 cases in this series. The fever is of the low grade type, the temperature usually normal in the morning, rises to 100.5, 101 or rarely 102 F in the late afternoon. It is irregular in type, though often persisting, it is not accompanied by chills or sweats. Many, if not most of the cases are afebrile throughout. The fever characterizes the acute or inflammatory phases of the process, the intervening stages of remission are afebrile, the late cicatrizing phase of the disease is again afebrile.

The continuous low grade fever of ileitis like that of segmental or right sided colitis is frequently mistaken, where the diarrhea is minimal and unnoticed, for many of the other diseases which are also characterized by low grade fever and low leukocyte counts in the circulating blood. Undulant fever, brucellosis, subacute bacterial endocarditis, lupus erythematosus disseminatus are frequently the connotations given to the low grade febrile type of

ileitis without any noticeable diarrhea or abdominal pain.

The leukocyte count in the circulating blood in ileitis is not usually materially elevated. The range may be between 9,000 and 12,000 white blood cells, rarely higher. A shift to the left of the polymorphonuclear leukocytes and toxic granulations is rarely seen in contrast to severe ulcerative colitis where both latter conditions are frequent. Eosinophilia, as in ulcerative colitis is common, the rate rising to 4 to 8 or 12 per cent in a small number of cases. The explanation of the eosinophilia in both conditions is not apparent. The attempt to explain the eosinophilia on an allergic basis is not satisfactory, nor are foreign body inclusions a reasonable answer, the eosinophilia is part of the general systemic reaction to the disease. The leukocyte counts given by Koster are considerably higher than observed in this series. He noted frequent rises to 16,000 or 20,000 or even above this figure. This is mainly to be explained by the fact that as a surgeon he saw a greater percentage of cases of acute ileitis. It is true that in acute ileitis the total white count is higher and a greater percentage of polymorphonuclear leukocytes is present in the circulating blood a fact that only intensifies the confusion in differentiating acute ileitis from acute appendicitis. An increased sedimentation rate of the red blood cells is usually observed, according to Snapper, Pompen and Groen.

A moderate often progressive anemia characterizes regional ileitis. In advancing cases the hemoglobin reading may range from between 60 per cent and 70 per cent on the Sahli scale. Lower readings are not unusual even in the absence of gross intestinal hemorrhage. In one case in this series the hemoglobin fell to 32 per cent Sahli without evidence of melena. In another case twice operated upon for recurrent ileitis the severe anemia of 16 per cent hemoglobin was explained when the true picture of a primary pernicious anemia became apparent in the blood smears. An achylia gastrici was present, reticulocyte response to liver extract was absent in this individual. The

pernicious anemia in this case was incidental and probably not related to the regional and recurrent ileitis

Gastric secretory changes in ileitis are unusual. One regularly notes a normal type of curve in the fractional test meal, the secretory curve may, however, be low (hyp acidity). Apart from the above cited case complicated by pernicious anemia, only once was achylia gastrica seen in this series.

The *general symptoms* of ileitis are those of weakness, loss of strength and loss of weight. A material weight loss was noted in 57 cases in this series, in one instance the loss amounting to 50 pounds. In the advanced cases, those with intestinal obstruction, and following debilitating operations, the loss of flesh may be much greater and extreme, in terminal cases amounting to emaciation. This extreme is, however, unusual. The average case may lose 10, 15 or 20 pounds, the patient finds it difficult to regain weight because of the persistent diarrhea and abdominal pain. Following successful operative procedure the regain of weight is rapid and ample. The loss of strength is not extreme, many patients with mild ileitis are able to continue at their work with little loss of efficiency. In one instance, a man in the Armed Forces during the recent war was able to continue an active service for three years after the onset of diarrhea which signalized the initiation of the ileitis, the diagnosis then being unrecognizable or unrecognized. In a man 71 years of age who complained of anorexia, weakness, moderate diarrhea and a loss of 25 pounds of weight simulating cachexia, the entire clinical picture was one of *masked carcinomatosis*. Repeated x ray examinations of the gastrointestinal tract were reported as negative. Eventually exploratory operation revealed a regional ileitis involving 60 cm (24 inches) of terminal ileum.

Retarded growth and delayed secondary sexual maturation was noted only once in this series of regional ileitis. Where the higher reaches of the ileum and the jejunum

are involved the interference with growth and maturity will be observed to be much higher

Menstruation is usually regular and maintained in contrast to ulcerative colitis, where amenorrhea characterizes the more severely toxic nature of that disease

Joint pains were present in 4 of this series, the larger joints such as ankle, knee and elbow being usually involved, as is so commonly seen in segmental colitis. The small joints of the hands and feet are spared in ileitis. Clubbing of the fingers of the hands occurred four times in this series

The nervous, or rather, psychic manifestations of this disease are sometimes so manifest that they overwhelm the true somatic manifestations. This occurred in 10 of this series of cases, probably much more often. In 6 of the cases, the onset of symptoms occurred while in active service in the Armed Forces. Most of the cases were regarded as psychoneurotic following exposure to battle, or due to bomb shock. Some were classed as nervous diarrhea, or as food poisoning. In 2 of the cases, the diagnosis was properly made by the Army. During the war I had ample opportunity to visit neighboring camps in the vicinity of New York City and was shown many correctly diagnosed cases of ileitis. In civilian life, 4 cases were treated as depressions or as neurasthenia, 2 of them in institutions for mental care. The evident psychic and nervous symptoms that accompany ileitis often lead to delayed recognition of the true cause for the symptoms. Of Blackburn's 22 cases, 1 suffered from a true psychosis. 4 were referred to psychiatrists for mental treatment when, as he states, "a routine clinical, pathological and radiological investigation would undoubtedly have shown organic disease of the ileum."

However, ileitis can hardly be classed as a psychosomatic disease, occurrences or recurrences do not follow upon psychic shock as evidently as they can be observed in ulcerative colitis with obvious relationship of cause to effect.

pernicious anemia in this case was incidental and probably not related to the regional and recurrent ileitis.

Gastric secretory changes in ileitis are unusual. One regularly notes a normal type of curve in the fractional test meal, the secretory curve may, however, be low (hyp acidity). Apart from the above cited case complicated by pernicious anemia, only once was achylia gastrica seen in this series.

The *general symptoms* of ileitis are those of weakness, loss of strength and loss of weight. A material weight loss was noted in 57 cases in this series, in one instance the loss amounting to 50 pounds. In the advanced cases, those with intestinal obstruction and following debilitating operations, the loss of flesh may be much greater and extreme, in terminal cases amounting to emaciation. This extreme is, however, unusual. The average case may lose 10, 15 or 20 pounds. The patient finds it difficult to regain weight because of the persistent diarrhea and abdominal pain. Following successful operative procedure the regain of weight is rapid and ample. The loss of strength is not extreme. Many patients with mild ileitis are able to continue at their work with little loss of efficiency. In one instance a man in the Armed Forces during the recent war was able to continue an active service for three years after the onset of diarrhea which signalized the initiation of the ileitis, the diagnosis then being unrecognizable or unrecognized. In a man 71 years of age who complained of anorexia, weakness, moderate diarrhea and a loss of 25 pounds of weight simulating cachexia, the entire clinical picture was one of masked carcinomatosis. Repeated x-ray examinations of the gastrointestinal tract were reported as negative. Eventually exploratory operation revealed a regional ileitis involving 60 cm (24 inches) of terminal ileum.

Retarded growth and delayed secondary sexual maturation was noted only *once in this series of regional ileitis*. Where the higher reaches of the ileum and the jejunum

are involved the interference with growth and maturity will be observed to be much higher

Menstruation is usually regular and maintained in contrast to ulcerative colitis, where amenorrhea characterizes the more severely toxic nature of that disease

Joint pains were present in 4 of this series, the larger joints such as ankle, knee and elbow being usually involved, as is so commonly seen in segmental colitis. The small joints of the hands and feet are spared in ileitis. Clubbing of the fingers of the hands occurred four times in this series

The nervous, or rather psychic manifestations of this disease are sometimes so manifest that they overwhelm the true somatic manifestations. This occurred in 10 of this series of cases probably much more often. In 6 of the cases, the onset of symptoms occurred while in active service in the Armed Forces. Most of the cases were regarded as psychoneurotic following exposure to battle or due to bomb shock some were classed as nervous diarrhea, or as food poisoning. In 2 of the cases, the diagnosis was properly made by the Army. During the war I had ample opportunity to visit neighboring camps in the vicinity of New York City and was shown many correctly diagnosed cases of ileitis. In civilian life, 4 cases were treated as depressions or as neurasthenia. 2 of them in institutions for mental care. The evident psychic and nervous symptoms that accompany ileitis often lead to delayed recognition of the true cause for the symptoms. Of Blackburn's 22 cases 1 suffered from a true psychosis. 4 were referred to psychiatrists for mental treatment when as he states, "a routine clinical, pathological and radiological investigation would undoubtedly have shown organic disease of the ileum."

However ileitis can hardly be classed as a psychosomatic disease. occurrences or recurrences do not follow upon psychic shock as evidently as they can be observed in ulcerative colitis with obvious relationship of cause to effect

## PHYSICAL EXAMINATION

The two outstanding features in the physical examination of the cases of regional ileitis are (1) an abdominal mass and (2) evidence of fistula formation external and internal

The abdominal mass (observed in 76 cases in this series) is usually found in the right lower quadrant of the abdomen, in an anatomic area corresponding to the inner surface of the cecum. The mass varies in size from that of a finger, to that of a lemon, a large orange, occasionally even larger. The mass is usually fixed to the posterior peritoneal wall, to the cecum, or to the anterior abdominal wall, particularly when an external fistula is present leading anteriorly from the mass. Tenderness on palpation is constantly present, when the mass is small and in its typical site it is with difficulty differentiated from a spastic cecum. The mass consists materially of the inflamed terminal loop of ileum and of adherent loops of nearby ileum or of cecum, ascending colon or sigmoid.

On separating the segments of the mass at the operating table, it will usually be seen that the intestinal loops are adherent because of single or multiple intercommunicating fistulous tracts. The mass, particularly when adherent to the dome of the urinary bladder, may be apparent in the lower mid abdomen, in one instance simulating a urachus cyst. In another instance the loops of ileum were adherent to and drawn over by the sigmoid colon, the mass being palpable to the left of the midline of the abdomen.

The most common seat for the mass is low in the right lower quadrant of the abdomen, palpable frequently by rectal or vaginal pelvic examination or by the bimanual technic. This fact is easily explainable: the position of the terminal ileum is dependent from the cecum. Testut places the cecum below the pelvic inlet in 15 per cent of males and in 30 per cent of female bodies. In the upright position, the cecum and loops of terminal ileum fall into the pouch of Douglas in the male and between the uterus and the rectum in the female. Very often these loops lie directly on the pelvic floor (fig 11) supported by the Levator Ani

muscle. Monks found the terminal ileum to be in the pelvis in more than 50 per cent of his studies. In our own




Fig. 11 —Anatomic Position of the Normal Terminal Ileum (1) Usual position of cecum and ileum (2) high cecum ileum to pelvis (3) low cecum high medial implantation of ileum (4) complete descent of cecum and ileum in pelvis (5) exceptional position of terminal ileum abrupt at cecum. The majority of cases fall within groups 1 and 3. Reprinted from Radiology 33:323-330 1939



radiographic studies, made to determine the position of the normal ileum, it was found (Crohn and Yarnis) that in 150 control normal cases descent of the cecum and terminal ileum into the lower pelvis occurred in 28 per




Fig 12—Anatomic Position of the Diseased Terminal Ileum as Classified from a Study of 25 Cases of Terminal Ileitis. The weight of the diseased loop carries it so as to lie on the pelvic floor. Reprinted from *Radiology* 33: 325-330 1939

cent of the observed cases in that series. In the pathologic ileum (25 cases of terminal ileitis) the terminal affected loop of ileum was found in the pelvis in all cases, in fact on the floor of the pelvic cavity (fig 12). It is apparent that the soggy weighted diseased ileum descends low down in the right lower abdominal quadrant and being even heavier than normal in weight occupies almost uniformly the pelvic inlet. The mass of ileitis is then found and easily palpated

by bimanual technic in the fossa of Douglas. Only when adherence to other organs takes place is the mass felt other than in the right lower abdominal cavity. When fistulous tracts burrow into and through the apposed base of the mesentery, the necrotic process may cause a diffuse suppurative mesenteritis which participates in and increases the size of the inflammatory mass.

Apart from the abdominal mass and the demonstration of characteristic fistulas the physical examination of the cases of ileitis is likely to be negative. Enlargement of the spleen was noted four times, clubbing of the fingers four times not necessarily in the same patient. Both manifestations indicate the degree of general body affection as part of the localized intestinal process.

Erythema nodosum on the anterior aspect of the lower extremities was noted once in acute ileitis and twice in chronic regional ileitis. This phenomenon, which is so much more prevalent in ulcerative colitis possibly links the two diseases in a common etiological factor.

#### OBSTRUCTIVE PHENOMENA

Intestinal obstruction occurred in 10 instances in this series of 222 cases usually late in the course of the disease. Twice in this series it was an early, in fact the presenting symptom at the onset; in the other cases it indicated a long standing, chronic inflammatory process eventuating in chronic cicatricial stenosis of the bowel at the site of the lesion. Dickinson and Zimmerman report a single case with intestinal obstruction as the initial presenting symptom. Sherill and Hall indicate 2 cases with obstruction as the initial phenomenon though both were old chronic cases. Jackman reported 2 cases of hypertrophic enteritis as a cause of intestinal obstruction both cases of the acute type in both of which the terminal ileum was perforated during the operative manipulations.

Occasionally the obstruction may be due to angulation and kinking of a loop of bowel by peritoneal adherence or by ileo ileal fistulation. Loops of distended intestines

radiographic studies, made to determine the position of the normal ileum, it was found (Crohn and Yarnis) that in 150 control normal cases descent of the cecum and terminal ileum into the lower pelvis occurred in 28 per



Fig. 12—Anatomic Position of the Diseased Terminal Ileum as Classified from a Study of 25 Cases of Terminal Ileitis. The weight of the diseased loop carries it so as to lie on the pelvic floor. Reprinted from *Radiology* 33: 325-330, 1939.

cent of the observed cases in that series. In the pathologic ileum (21 cases of terminal ileitis), the terminal affected loop of ileum was found in the pelvis in all cases, in fact on the floor of the pelvic cavity (fig. 12). It is apparent that the soggy weighted diseased ileum descends low down in the right lower abdominal quadrant and being even heavier than normal in weight, occupies almost uniformly the pelvic inlet. The mass of ileitis is then found and easily palpated

by bimanual technique in the fossa of Douglas. Only when adherence to other organs takes place is the mass felt other than in the right lower abdominal cavity. When fistulous tracts burrow into and through the apposed base of the mesentery the necrotic process may cause a diffuse suppurative mesenteritis which participates in and increases the size of the inflammatory mass.

Apart from the abdominal mass and the demonstration of characteristic fistulas the physical examination of the cases of ileitis is likely to be negative. Enlargement of the spleen was noted four times, clubbing of the fingers four times not necessarily in the same patient. Both manifestations indicate the degree of general body affection as part of the localized intestinal process.

Erythema nodosum on the anterior aspect of the lower extremities was noted once in acute ileitis and twice in chronic regional ileitis. This phenomenon, which is so much more prevalent in ulcerative colitis possibly links the two diseases in a common etiological factor.

#### OBSTRUCTIVE PHENOMENA

Intestinal obstruction occurred in 10 instances in this series of 222 cases, usually late in the course of the disease. Twice in this series it was an early, in fact the presenting symptom at the onset; in the other cases it indicated a long standing chronic inflammatory process eventuating in chronic cicatricial stenosis of the bowel at the site of the lesion. Dickinson and Zimmerman report a single case with intestinal obstruction as the initial presenting symptom. Sherill and Hall indicate 2 cases with obstruction as the initial phenomenon though both were old chronic cases. Jackman reported 2 cases of hypertrophic enteritis as a cause of intestinal obstruction both cases of the acute type in both of which the terminal ileum was perforated during the operative manipulations.

Occasionally the obstruction may be due to angulation and kinking of a loop of bowel by peritoneal adherence or by ileo ileal fistulation. Loops of distended intestines

radiographic studies, made to determine the position of the normal ileum, it was found (Crohn and Yarnis) that in 150 control normal cases descent of the cecum and terminal ileum into the lower pelvis occurred in 28 per




Fig 12—Anatomic Position of the Diseased Terminal Ileum as Classified from a Study of 25 Cases of Terminal Ileitis. The weight of the diseased loop carries it so as to lie on the pelvic floor. Reprinted from Radiology 33 323-330 1939

cent of the observed cases in that series. In the pathologic ileum (25 cases of terminal ileitis) the terminal affected loop of ileum was found in the pelvis in all cases, in fact on the floor of the pelvic cavity (fig 12). It is apparent that the soggy weighted diseased ileum descends low down in the right lower abdominal quadrant and being even heavier than normal in weight, occupies almost uniformly the pelvic inlet. The mass of ileitis is then found and easily palpated

## 5 Fistula Formation in Regional Ileitis

---

FISTULA FORMATION is the most constant clinical and pathologic phenomenon in ileitis. These intestinal fistulas are of three types:

1. Fistulas to the anterior abdominal wall.
2. Internal fistulas between loops of bowel and adherent hollow viscera.
3. Rectal and perirectal fistulas.

Fistulas to the anterior abdominal wall are among the most common types of fistulation and occur exclusively in the site of the scar of a previous laparotomy. The antecedent operation in most of the cases (30.6 per cent) was an appendectomy. We encountered 36 instances of anterior wall fistulas, all of them in the appendectomy scar or in the scar of a previous exploratory laparotomy. Fistulas to the abdominal wall are infrequent after simple appendectomy for acute appendicitis (Ginzburg). According to Colp, in 2940 cases of acute appendicitis abdominal wall fistulas were found only 33 times, an incidence of 1.1 per cent; only 3 required reoperation. Clark and Dixon in 44 cases of regional enteritis found 9 external fecal fistulas, one of which passed feces and gas through the urethra. Four of these 9 patients also had intraintestinal fistulas, one a fistula to the urinary bladder.

In some of the instances of ileitis not a single but multiple fistulas were present. In one case 4 anterior wall fistulas in several scars of eight previous laparotomies (Crohn) [58]. These fistulas occur usually within a few weeks of the apparent healing of the operative scar. The scar

may be seen through the thin abdominal wall, visible peristalsis is not uncommon and is accompanied by severe abdominal cramps and borborygmi. A scout film of the abdomen will show puddling and fluid levels. Vomiting may be present in the more severe stenotic phases. The obstruction can usually be temporarily alleviated by the use of the Miller Abbott tube.

Postoperative intestinal obstruction, such as characterizes many cases of intestinal resection or short circuiting operations is unfortunately only too frequent but fortunately does not occur as frequently as in reconstructive surgery for ulcerative colitis.

rise to some speculation as to their nature. They are apparently lytic rather than infectious in type. They traverse large areas of mesentery and distal fascial planes without giving rise to infection terminating their course by finding an exit point somewhere on the surface of the body. They are lytic or proteolytic the activated intestinal or pancreatic enzymes of the fecal chyme slowly digesting their way through planes of fascia and muscle to exit on




Fig. 13—Fistulation Terminal Ileum to any Adjacent Hollow Viscera. Note termination in such unusual sites as right fallopian tube, left ureter, body of uterus.

the body wall. Such fistulas become infected and suppurate only as a result of skin bacterial contamination at the point of exit from the abdominal cavity.

#### INTERNAL FISTULAS (FIG. 13)

The mechanism of formation of internal intestinal fistulas is identical with that of external abdominal wall fistulas. These tracts originate in the sodden leaking ulcerated terminal ileum usually on the side of mesenteric attachment. They traverse the mesentery of the small bowel and terminate in an adjacent loop of small intestine,



becomes reddened, and inflamed at a localized area, a low grade fever becomes apparent with regional pain and tenderness. An abscess forms and either breaks spontaneously or is incised, following which fecal drainage occurs and a chronic fecal fistula is established. In some instances the abdominal wall fistula occurs only months or years later, two years, four years and in one instance eight years after a presumably successful resection for terminal ileitis. The eight intervening years were marked by excellent health until the appearance of the fistula signaled a recurrence of the ileitis.

These external wall fistulas are apparently very facile of creation, in two instances just knocking the diseased bowel wall accidentally during operation resulted in immediate fistula formation in the healing incision, even though great surgical care was immediately exercised in suturing the cut in the wall of the ileum.

Spontaneous fistulas to the abdominal wall without a previous laparotomy have not been observed. But unusual exit sites for external fistulas are occasionally noted. Multiple external fistulas to the inguinal region have been noted twice, at least four fistulas in one instance, there being no previous operative scar. Hurst has noted a fistulous tract originating in the ileum and pointing downward and infiltrating the extraperitoneal tissues in the right groin. Brown and Donald reported 47 cases with fistula formation out of a total of 178 cases an incidence of 26.5 per cent. This is a higher incidence of fistulation than encountered in this present or in most of the current literature. They further state that fistula formation occurs invariably only after surgical procedures.

Fistulas to the lumbar region have been seen three times twice by the author and once mentioned and reported by Snapper, where a series of external fistulas occupying the right lumbar area were observed expelling vegetable matter and fecal content. The fistulas were traced to a terminal ileitis. The linear extent of these intestinal fistulas gives

rise to some speculation as to their nature. They are apparently lytic rather than infectious in type. They traverse large areas of mesentery and distal fascial plane without giving rise to infection, terminating their course by finding an exit point somewhere on the surface of the body. They are lytic or proteolytic, the activated intestinal or pancreatic enzymes of the fecal chyme slowly digesting their way through planes of fascia and muscle to exit on




Fig. 13—Fistulation. Terminal Ileum to any Abdominal Hollow Viscera. Note termination in such unusual sites as right fallopian tube, left ureter, body of uterus.

the body wall. Such fistulas become infected and suppurate only as a result of skin bacterial contamination at the point of exit from the abdominal cavity.

#### INTERNAL FISTULAS (FIG. 13)

The mechanism of formation of internal intestinal fistulas is identical with that of external abdominal wall fistulas. These tracts originate in the sodden, leaking, ulcerated terminal ileum usually on the side of mesenteric attachment. They traverse the mesentery of the small bowel and terminate in an adjacent loop of small intestine,

in any segment of the large bowel, or in any adjoining hollow intra abdominal viscus, such as ureter, urinary bladder, uterus or fallopian tube

The internal fistulas were noted in 23 of our series (approximately 10 per cent) though the incidence of such internal fistulas would appear in practice to be much higher



Fig. 14.—Diagrammatic Sketch. Fistulation Terminal Ileum to Small Intestine and Cecum

Variations of these internal fistulas may be listed as in table 6

TABLE 6—Variations of Internal Fistulas	
Terminal ileum to	
sigmoid	6 instances
cecum	3 instances
ascending colon	3 instances
other loops of ileum	1 instance
duodenum	1 instance
urinary bladder	4 instances

Other variations are ileum to right ureter, to left ureter to urinary bladder, to the intra uterine cavity to the right fallopian tube

On two occasions the opening of the fistula into the sigmoid colon was seen by sigmoidoscopy. In another instance the fistula which broke into the right fallopian tube gave rise to a salpingitis discovered inadvertently by a gynecologist operating for supposed adnexal inflammation. The discharge of the proteolytic content of the fis-



Fig. 13.—Combined Ileitis (Terminal) with Involvement of the Ascending Colon and Cecum. Note fistulous tracts between ileum and base of cecum. Meckel's diverticulum not involved (Involved segment to the left; Meckel's diverticulum inverted).

tulous tract into any hollow viscus usually terminates its course.

Probably the most common of all these internal fistulas are the ileocolic variety; they are readily demonstrated at the operating table (figs. 14 to 16). Many or most of the fistulous tracts are notable and visible in the preoperative radiographic films provided they are diligently sought after (figs. 17 and 18). The fine lines seen with a magnifying lens emanating from the diseased ileum indicate potential fistulas

(fig 18) These are frequently seen overlying the dome of the urinary bladder and creating urinary symptoms, actual gross fistula exit into the urinary bladder is rare in ileitis,




Fig 16—Terminal Ileitis Multiple Fistulas Regional Skip Areas throughout Ileum and Jejunum A S 21 years male Regional colitis Picture after three month of medical treatment

much less common than are such similar fecal fistulas which arise in a sigmoid diverticulitis Forbes and Duncan report an interesting case of ileitis in which a large fistula existed between the terminal ileum and the urinary bladder, the patient passed gas and fecal material per urethra Ileo

transverse colostomy without transection of the ileum was performed but the intestinal symptoms and those of the ileo vesical fistula persisted. At a second operation, the ileum distal to the anastomosis was transected as the sole procedure except for a temporary ileostomy of the distal ileal loop. Within four months the patient gained 25



Fig 17—Terminal Ileitis. Fistula, Ileum to Sigmoid

pounds in weight was free of symptoms and the dysuria and passage of gas and feces per urethra had disappeared. Unfortunately a severe recurrence of the ileitis occurred after the date of publication of their article. In another case cited by these authors an ileo vesical fistula persisted for some time in spite of an ileo colostomy. This fistulous tract was resected broke down again with the secondary formation of an additional fecal fistula but eventually both fistulous tracts healed with subsidence of all symptom.

Brown and Donald culled from a review of 178 cases of regional enteritis with complications of fistulas the variations in the site of the disease as shown in table 7

TABLE 7—*Site of Regional Enteritis with Fistula (Brown and Donald)*

Site of disease	Fistulas	
	Internal	External
Jejunum—0 cases	0	0
Upper ileum—8 cases	0	1
Lower ileum—99 cases	4	18
Ileum and Colon—66 cases	4	14



Fig. 18—Regional (Terminal) Ileitis with Vascular Symptom—Multiple Minute Formation of Fistula

The exemption of the jejunum to fistula formation is thus seen, the lower ileum is the site of most of the internal and

external fistula formation Pugh describes fistulation in six of his cases the fistulas being identified as in table 8

TABLE 8—*Fistula from Ileum (Pugh)*

Ileum to	
sigmoid	—
cecum and jejunum to cecum	1
sigmoid to urinary bladder	1
cecum	1
transverse colon	1

### PERIRECTAL FISTULAS

In 1934, Bassell published a case report of terminal ileitis in which multiple rectal fistulas occurred as a complication of the intestinal disease. He felt that the presence of the inflammation of the ileum in the cul de sac could easily have given rise to a rectal abscess and persistent rectal fistula. In 1938 the diagnostic significance of perianal, rectal and rectovaginal fistulas was emphasized [190]. In that publication 20 cases of perirectal fistulation were noted in a group of 110 cases of ileitis an incidence of 18.2 per cent. This was almost equal to the number of external abdominal wall fistulas (12 cases) and internal intestinal fistulas (11 cases) combined as shown in that series.

In the present series of 222 cases of regional ileitis, perirectal abscesses and fistulas were noted 39 times in incidence of 17.5 per cent. This does not include 3 instances of rectovaginal fistulation. The total number is somewhat below the combined number of external and internal intestinal fistulas in this present series (59 cases 26.5 per cent).

Jackman and Smith in 1943 recorded at the Mayo Clinic 36 cases of anal abscess or fistula in a group of 114 cases (31.6 per cent) a figure high above those in this series. They included those cases having had a history of operation for anal fistula within a three year period prior to their initial visit. Eight of the series of 114 cases came to the Clinic primarily because of anal fistula the intestinal disturbance being so mild as to have been overlooked.

### METHOD OF ANAL FISTULATION

Either of two methods may explain anal suppuration (1) Most of these fistulas occur as a result of infection in



the rectal crypts of Morgagni above the anal sphincter, the contaminated fecal infectious material being transported thereto by the propulsion of the fecal column. The infection of the crypts gives rise to a perirectal abscess or an ischio rectal abscess which opening simultaneously on the surface of the buttock and within the anal canal forms a permanent fistula. When the abscess localizes anteriorly in the female, it may break simultaneously through the




Fig. 19—Gravity Fistula. Route of Direct Fistulation from Terminal Ileum to Perirectal Spaces. This type is comparatively rare in comparison with the type which originates in transported infection with lodgement in the crypts of Morgagni.

perineal body into the vagina and posteriorly into the rectum, forming a recto vaginal fistula.

This type of local rectal infection in ileitis is almost identical with that seen in ulcerative colitis. In both, the internal opening of the fistulous tract is just within the anal margin, in the recto vaginal fistulas the anterior opening is just within and above the fourchette, constituting a small opening which emits a thin fecal leakage.

(2) The second method of fistulation to and about the anus and perineum is more problematic. This fistulous tract is considerably more circuitous proteolytic material escapes from the porous terminal loop of ileum lying heavily on the pelvic floor. The material seeps downward through the pelvic peritoneal reflection and the retroperitoneal fat and burrows through the pelvic fascia following the planes of the levator ani muscle (fig. 19). The material makes its exit into the rectum above the sphincters, or, piercing the slinglike attachments of the fibers of the levator ani makes its exit at the perianal margins. The ischio rectal fossa may be contaminated forming an abscess which will reveal itself as a perirectal fistula communicating internally with the rectum. Attempts to follow such long fistulous tracts are difficult. Lipiodol injected into the tract from below will ascend for several inches but will be lost somewhere in the pelvis. A streak of barium following a radiographic study may be seen pointing downward in the same direction. A direct and continuous path has not been unequivocally demonstrated except in one case in the literature where at autopsy the continuity of the path was clearly dissected (Patricelli).

The diagnostic significance of perianal fistulas has not been sufficiently appreciated. They may occur one to fourteen years before the onset of active diarrhea and abdominal pain constituting the early prodromal manifestations of the disease.

It is a fair statement to make as a generalization that suppurative perianal fistulas in the presence of diarrhea indicate a pathologic inflammatory process somewhere in the intestinal tract. These fistulas occur with great frequency in ileitis, ulcerative colitis and in intestinal tuberculosis. They do not occur in nervous gastrogenous, pancreatic allergic or thyrogenic diarrheas. Exceptions to this general rule have not been noted however for the sake of scientific accuracy it would be better to keep an open mind though the general truth of the statement will probably continue to operate.

## 6 The Course in Regional Ileitis

---

REGIONAL ILEITIS may begin as an acute phase of the disease, or it may be first observed as an acute exacerbation of a clinical process which has been overlooked, in the vast majority of cases it comes under observation as a disease of great chronicity. The acute case, simulating acute appendicitis (except for the presence of diarrhea as a symptom) resolves in 25 per cent of the cases without further evidence of recrudescence. The remainder of the acute cases subsides spontaneously, usually in spite of the laparotomy and customary appendectomy and lapse into the low grade typical chronic course of the disease.

The course of long standing, chronic regional ileitis is one of great chronicity. Spontaneous remissions that endure for years may occur, during which remissions the diarrhea will abate or be absent, the temperature remain normal, appetite be preserved and nutrition be only slightly impaired. Except for a slight and slowly progressive loss of weight, mild persistent anemia and occasional short bouts of diarrhea and abdominal pain, the patient may follow a normal career of physical well being and school or occupational activity. As the disease progresses in the intestinal tract the periods of activity will be greater and last longer, the remissions become shorter. In time either the continuous diarrhea or the loss of weight and strength will indicate the progression into a more severe phase of the disease. Eventually the onset of obstruction (10 cases) will supervene with abdominal pains, vomiting, and the inability to feel an abdominal mass on palpating the abdomen.

At any time in the course of the disease a peri rectal abscess or a fistula in the scar of the abdominal wall may supervene.

In the continued study of chronic regional ileitis with the assistance of radiographic observation, it will be noted that three variations in the course must be considered. In the first group, spontaneous healing of the lesion with disappearance of all radiographic evidence will supervene. This occurred in 13 of this series of 222 cases of regional ileitis. The second group is constituted of cases in which anatomically the lesion remains static, usually localized to the terminal ileum; progression upward of the pathologic process cannot be radiographically demonstrated. In the third group the lesion is obviously a progressive one and continues to spread mildly either continuously or by skip lesions until the upper reaches of the ileum or the jejunum become involved.

### Group 1

Chronic long standing regional ileitis with spontaneous healing (13 cases). It was surprising to note the fact that spontaneous healing could take place in regional ileitis; this even in cases where internal fistulation had been radiographically demonstrated. In 5 of these cases the diagnosis had been first observed and confirmed at the time of previous appendectomy. Some of the details of these 13 cases are of great interest particularly since spontaneous resolution of the inflammatory process is so unusual.

*Case 1. H. S.* Male. Age 27 years. Six years of pain and diarrhea. At exploratory operation in 1940 3 feet of terminal ileum were observed to be involved; the operation was restricted to the exploration. Four years later the patient was well with one or two firm stools and a regain of weight (x rays were completely negative).

*Case 2. J. B.* (Serial No. 19) Male 43 years of age. For two years previously the patient had suffered with loose and frequent bowel movements, fever up to 10 F, predefecation cramps and a loss of 25 pounds of weight. At the radiologic examination a typical terminal ileitis involving the last 18 inches of ileum was easily discernible (1943). The course of the disease for four years was benign, practically nonsymptomatic. An acute incident resembling a subacute

intestinal obstruction supervened (1947) but subsided without operation under medical treatment. The patient has remained well to date.

**Case 3 A H** (Serial No 70) Male, aged 42 years. Fifteen years previously the patient underwent an appendectomy. The wound suppurated for four months. Two years later the wound broke open establishing a fecal fistula which later closed spontaneously. He remained well for thirteen years except for slight attacks of abdominal pain and mild diarrhea. When seen in 1942 he was in good health; the radiologic examination disclosed a string sign involving the terminal 12 inches of the ileum. On physical examination a suggestion of a mass was palpable under the lower right rectus muscle. The follow up two years later reported the man as in perfect health and free of all symptoms.

**Case 4 J C** Female age 27 years. Appendectomy 1938. For six years following this operation continued pain, regular bowel movements. Marriage and birth of two healthy children without complications or difficulties. Subsequent onset of diarrhea. Subsidence of all symptoms but persistence of radiographic evidence of disease involving the distal 2 feet of the terminal ileum. Regarded as well in view of subsidence of all clinical symptoms for the last two years (1946).

**Note** August 1948—exacerbation of pain and diarrhea, suspicion of low grade obstruction. The reversal of the prognosis in the case from that of apparent cure to active symptoms suggesting surgical intervention is most disappointing.

**Case 5 H 4** Female. Terminal ileitis, fistula ileum to cecum. No operation. Three years later mass had disappeared, internal fistula no longer visible, patient well. Radiographic examination negative.

**Case 6 S F** (Serial No 50) Female age 47 years, widow. At 18 years of age appendectomy. Intestine seen to be inflamed. Twelve years ago cholecystectomy for multiple biliary calculi. Two years abdominal cramps, diarrhea, four to five watery movements daily. Physical examination negative when first seen in 1944. Radiographic examination marked inflammatory involvement of terminal two loops of ileum. Follow up three years later well, asymptomatic. Persistent evidence of fibrotic lesion in terminal ileum was demonstrated by recent radiographic examination.

**Case 7 I C** (Serial No 29) Male age 16 years. Had been operated upon when 8 weeks of age for congenital pyloric stenosis. Three years ago had been operated for supposed intestinal obstruction due to adhesions(?) For the last two years following upon this operation (1944) had suffered bout of diarrhea and abdominal pain with occasional febrile exacerbations. The radiographic examination at that time was reported as negative. Radiographic study in 1946 distinct irregularity of terminal ileum. Subsequent and objective improvement under sulfathiazine therapy. When last seen October 1947 well and symptom free.

**Case 8 E A** (Serial No 209) Female age 39 years, married, three healthy children. For five years has suffered with recurrent diarrhea.

and periumbilical pains loss of 9 pounds of weight One year ago small bowel barium series showed a mild irregularity of the terminal ileum Repeat series (1945) by both barium enema and barium meal one year later showed definite and distinct lesion involving several loops of the terminal ileum A third radiographic study a year later (1946) demonstrated a definite involvement of the terminal twenty inches of the small intestine Marked clinical improvement regain of weight cessation of diarrhea Radiography almost two years later December 1947 revealed practically a normal terminal ileum

*Case 9 M I* (Serial No 213) Male 21 years of age Served in U S Navy during the war from 1943 to 1946 Onset of symptoms following the explosion of a bomb on the deck of a destroyer thrown some distance across the deck Developed abdominal pains nervousness fear of going below decks anorexia Sharp cramplike abdominal pains diarrhea three to four loose daily movements Hospitalized in Naval Hospital as case of psychoneurosis no intestinal radiographic studies made Radiographic studies 1946 strongly indicative of terminal ileitis Repeat radiographic studies by barium meal and barium enema showed involvement of terminal 12 inches of ileum Follow up 1947 and February 1948 patient well and free of symptoms

*Case 10 S C* Male 23 years of age single Appendectomy abdominal wall fistula spontaneous closure of fistula Radiographic evidence of terminal ileitis 1941 Well (1947)

*Case 11 A G* Female 23 years of age Appendectomy abdominal wall fistula Spontaneous closure of fistula Barium meal demonstrates inflammatory lesion of terminal ileum 1944 Follow up 1947 well and free of symptoms

*Case 12 H F* (Serial No 35) Male age 62 years Seen in 1939 for first time Ten year history of occasional attacks of watery diarrhea associated with mild abdominal pain Physical examination revealed vague mass under lower right rectus muscle No fever loss of 10 pounds of weight Radiographic examination revealed typical string sign involving terminal 16 inches of ileum The patient continued spontaneously to improve is now free of symptoms and continued so to this date nine years later (December 1947)

*Case 13 E G* (Serial No 63) Female age 19 years single Ten month history 5-6 feet of ileum involved in x ray Active diarrhea and fever Three years later only 1-1½ feet of ileum involved by radiographic study Three years later patient well radiographic examination negative

It will thus be observed that in a minority of fortunate individuals an apparent spontaneous cure is possible fistulas internal and external are under exceptionally favor

Since this list was compiled this patient has been operated for recurrence of his supposedly healed lesion

intestinal obstruction supervened (1947) but subsided without operation under medical treatment. The patient has remained well to date.

*Case 3 A H* (Serial No 70) Male aged 42 years. Fifteen years previously the patient underwent an appendectomy. The wound suppurated for four months. Two years later the wound broke open, establishing a fecal fistula which later closed spontaneously. He remained well for thirteen years except for slight attacks of abdominal pain and mild diarrhea. When seen in 1942 he was in good health, the radiologic examination disclosed a string sign involving the terminal 19 inches of the ileum. On physical examination a suggestion of a mass was palpable under the lower right rectus muscle. The follow up two years later reported the man as in perfect health and free of all symptoms.

*Case 4 J C* Female age 27 years. Appendectomy 1938. For six years following this operation continued pain regular bowel movements. Marriage and birth of two healthy children without complications or difficulties. Subsequent onset of diarrhea. Subsidence of all symptoms but persistence of radiographic evidence of disease involving the distal 2 feet of the terminal ileum. Regarded as well in view of subsidence of all clinical symptoms for the last two years (1946).

*Note* August 1948—exacerbation of pain and diarrhea, suspicion of low grade obstruction. The reversal of the prognosis in the case from that of apparent cure to active symptoms suggesting surgical intervention is most disappointing.

*Case 5 H 4* Female. Terminal ileitis fistula ileum to caecum. No operation. Three years later mass had disappeared internal fistula no longer visible patient well. Radiographic examination negative.

*Case 6 S F* (Serial No 50) Female age 47 years widow. At 18 years of age appendectomy. Intestine seen to be inflamed. Twelve years ago cholecystectomy for multiple biliary calculi. Two years abdominal cramps diarrhea four to five watery movements daily. Physical examination negative when first seen in 1944. Radiographic examination marked inflammatory involvement of terminal two loop of ileum. Follow up three years later well asymptomatic. Persistent evidence of fibrotic lesion in terminal ileum was demonstrated by recent radiographic examination.

*Case 7 I C* (Serial No 29) Male age 16 years. Had been operated upon when 8 weeks of age for congenital pyloric stenosis. Three years ago had been operated for supposed intestinal obstruction due to adhesions(?) For the last two years following upon this operation (1944) had suffered bout of diarrhea and abdominal pain with occasional febrile exacerbations. The radiographic examination at that time was reported as negative. Radiographic study in 1946, distinct irregularity of terminal ileum. Subsequent and objective improvement under sulfathiazole therapy. When last seen October 1947 well and symptom free.

*Case 8, F A* (Serial No 209) Female age 39 years, married three healthy children. For five years has suffered with recurrent diarrhea.

apparent health and in spite of apparent surgical cures. The final word on the life history of the disease as pertains to self healing may require future elaboration.



Fig. 21—Seven Months after Onset

#### GROUP 2. CHRONIC REGIONAL ILEITIS WITHOUT PROGRESSION

The exact percentage of cases in this group cannot be stated since comparative radiographic studies over the course of years is not often capable of fulfillment (figs 20-22). A sufficient number of cases have, however, been observed to allow the inference to be drawn that in many cases observed over the course of years radiographically the lesion remains stationary. This applies even in cases where skip lesions are already demonstrable. As for example

**CASE 115** (Serial No 216) Female age 32 married. First seen in November 1947. Nine years ago incision and drainage of a recto-vaginal fistula. Four years ago appendectomy. For nine years patient had been suffering with diarrhea and abdominal cramps four to five watery stools daily. Radiographic studies mid three years ago showed



able conditions, capable of self healing. Restoration to normal health and well being seems possible. Wilensky has published such a case of spontaneous healing in non specific granuloma of the terminal ileum, although his conclusions are based purely on radiographic readings, diag-



Fig. 20, 21 and 22—Terminal Ileitis. Conservative Medical Therapy for a Period of a Year and Two Months. Note absence of progression of lesion. Retraction of process to terminal 12 inches of ileum.

Fig. 20—At Onset

the patient had never been present in his case and there were no corroborative surgical exploratory findings to confirm the statements in this one case. These observations on self healing are made with the reservation that relapses have been noted even after as much as nine to sixteen years of

because of mild but persistent symptoms a barium meal was undertaken. This radiographic examination revealed involvement of the terminal 8 inches of ileum (1944). Two years ago (1946) obstructive phenomena now became apparent for the first time. One year later the patient underwent an exploratory laparotomy because of persistent abdominal pain and diarrhea. An ileo transverse colostomy with transection of the ileum was performed for a cicatrizing and contracting ileitis involving the terminal 8 inches of the ileum. In other words



Figs 23 24 25 and 26—Series of Radiographs Illustrating Progress of Lesion in a Case of Regional Ileitis. Observations extend for little more than one year.

Fig 23—Only the Terminal Ileum Involved

during four years of observation the lesion remained limited to that anatomic area which was present when first properly observed, namely the terminal 8 inches of ileum.

Case 3 F F (Serial No 202) Female single age 34 years. First seen in 1946. Fifteen years ago had been operated for a rectal fistula. Exploratory operation revealed the terminal 8 inches to be involved in what was then regarded as intestinal Crohn's disease. A localized resection was undertaken. Mild intestinal diarrhea continued for the

involvement of 5 cm (2 inches) of the terminal ileum. Similar radiographic studies carried out two years ago showed identical involvement of the last two inches of terminal ileum. Three months ago a barium meal demonstrated involvement of the last 2-3 inches of terminal ileum with the finding of a significant 'string sign'. A tender pipe stem like mass was now palpable in the right lower quadrant of the abdomen.



Fig. 22—After One Year and Two Months of Medical Treatment

This case is typical of the stationary or nonprogressive form of terminal ileitis.

*Case - D 4* (Serial No 10) Male age 15 years. First seen in April 1943. For the previous ten months had suffered with recurrent attacks of diarrhea and abdominal cramps the pains occurring just before defecation and relieved by that act. He had lost 20 pounds of weight in that period. A mild secondary anemia existed (hemoglobin 72 percent). The original x ray examinations failed to visualize the terminal ileum and the correct diagnosis was missed. One year later

These cases without extension of the disease over the course of years seem to lend themselves best to surgical intervention. While late intestinal obstruction is often characteristic of this chronic type, recurrences after operation are less likely to occur. This group should be the most satisfactory for surgical cure. In one case the lesion was recognized at exploratory operation eighteen years ago.



Fig. 16. —Two Month Later More Extensive Involvement of Whole Lower Bowel

the true nature was not recognized and no further surgical intervention was attempted. The disease lay dormant and presumably asymptomatic. Recurrence of diarrhea and a positive x-ray picture led to operation and resection of the persistent and unextended lesion in the terminal ileum.

### GROUP 3 CHRONIC REGIONAL ILEITIS WITH PROGRESSIVE EXTENSION

These are the cases in which the study of the life history of the disease over a course of years reveals active progression of the lesion in an oral direction. Again there

last fifteen years. At exploratory operation performed in 1946 the identical section of ileum namely 8 to 10 inches was involved a short circuiting operation ileo-sigmoidostomy with transection of the ileum above the diseased area was carried out.

*Case 4 I B* (Serial No 197) Male age 39 years. First seen in 1946. For ten years had had intermittent attacks of diarrhea and abdominal cramps. Eight years ago had been operated upon for a rectal fistula. For the last two years had severe recurrent attacks of abdominal cramps with fever up to 101.2 F the cramps at times being so severe as to require hypodermatic administration of morphine. Two



Fig. 24—Six Month Later Extension of Progress to Involve Approximately Two Terminal Loops of Ileum

years ago radiographic examinations supposedly demonstrated only a spastic colon. A progressive loss of weight amounting to 50 pounds had been observed during the last ten years. A radiographic examination performed one and a half years ago showed suggested involvement of the terminal loop three to four feet of the small bowel. Six months ago the patient underwent an exploratory laparotomy because of persistent symptoms of fever and diarrhea. At the operation the identical involvement of the last 3 to 4 feet of ileum was observed with the additional finding of a fistulous tract between terminal ileum and sigmoid. In spite of the mucosal type of ileitis here demonstrated and the lack of cicatrization or the identification of a typical string sign, the disease when finally explored by operation had not extended beyond the original site of involvement. A short circuiting operation was performed with apparent excellent results.

It is likely that this is the so called mucosal or submucosal type of case that gives the disappointing results after surgical therapy, the rapid progression probably indicates that much more of the upper bowel is involved than is demonstrable on the x ray plates

In the chronic case of regional ileitis remission of long periods may be observed such periods covering months and many years have been noted up to five to ten years between recurrences of diarrhea and fever. In one case the onset of symptoms dated eight years after a previous appendectomy for supposed "chronic appendicitis". In the later stages of the disease, the symptoms become increasingly severe, the recurrences more frequent, and of longer duration, until continuity of severe complaints causes complete disability

are insufficient personal observations to allow of any percentages to be offered, but an occasional observed case is illuminating. As examples

*Case 1* M H Male age 42 In 1935 exploratory operation for terminal ileitis fistula in ano Eight to 10 inches then involved no resection In 1943, eight years later whole ileum involved up to jejunum Hypoproteinemia marked emaciation




Fig. 26—Five Months Later Involvement of the Whole Ileum. This is the mucosal or submucosal type of ileitis with progressive extension.

*Case 2* (Figs 23-26) M W (Serial No 222) Female married 31 years of age First seen June 1947 For the last 13 months suffered with mild diarrhea and attacks of bloating and of abdominal cramps she has been constantly uncomfortable after eating She has sustained a loss of 30 pound in the last six months The dates and the results of the various radiographic examinations were as follows October 1946—8 to 10 inches of terminal ileum involved March 1947—10-12 inches of terminal ileum involved June 1947—18 inches of terminal ileum involved December 1947—30 inches of terminal ileum involved

In this last case, in which the successive radiographs were present for comparison, the oral progress of the lesion was capable of clear demonstration (figs 23-26)

the form of melæna, twice hematemesis was noted. Ellis quotes a case with repeated profuse and massive hemorrhage passed by the bowel (melæna). Two previous x-ray examinations had been negative. Eight transfusions were required to control shock. Eventually, at operation, 27 inches of ileum were resected with cure. Galambos and Mittelman cite a case of atypical terminal ileitis in which gastric hemorrhage and melæna dominated the clinical picture and led to the presumptive diagnosis of a peptic ulcer. Lunich and Crohn reported the case of a 53 year old male in which the onset of symptoms was ushered in by two massive intestinal hemorrhages in the form of melæna producing syncope and collapse. The second hemorrhage was almost fatal.

It is difficult to explain hematemesis when it occurs. In the 2 cases in which such a phenomenon was seen in this series, the diagnosis of regional ileitis was confirmed by exploratory operation. These both were cases in which the terminal ileum was involved, no higher segment of ileum or jejunum being implicated.

#### GENERAL COMPLICATIONS

In this series of cases of ileitis, duodenal ulcer was noted five times, cholelithiasis three times and acute cholecystitis was encountered once during an operation for terminal ileitis. Regional enteritis involving a Meckel's diverticulum has been observed (Horn and Rhoads) in one case with perforation at the base of the sac. In my own series I have once seen Meckel's diverticulum involved in a skip lesion. In another case a Meckel's diverticulum causing low intestinal obstruction by torsion was clinically preoperatively confused as a case of ileitis. Patricelli also mentions a case of regional ileitis with Meckel's diverticulitis with fatal outcome.

Some general complications which occur are actually incidental and not related to the disease. Thus this series includes 4 cases with chronic rheumatic heart disease, one



## 7 Complications of Regional Ileitis

---

### PERFORATION OF INTESTINE

ACUTE FRANK PERFORATION with generalized peritonitis is extremely rare, but can occur. Slow walled off perforation is manifest, particularly in the acute instances. In the chronic cases, on laparotomy, it is not uncommon to find a walled off abscess about the inner surface of the terminal ileum, limited by the neighboring loops of ileum and the mesial wall of the cecum, the whole being agglomerated in one confused mass. Upon separating the loops of involved ileum there may be a gush of a small quantity of pus. The neighboring loops are covered with fibrinous shaggy peritoneal deposit, a small quantity of inflammatory cloudy fluid may be present in the peritoneal cavity (300-400 cc). The granulomatous low grade type of infection with concomitant fibroblastic reaction precludes the occurrence of free open perforation with only rare exceptions. Attempts to resect terminal ileitis in the presence of a walled off abscess with localized peritonitis is fraught with a high operative mortality. Arnheim reports acute perforation of five days' standing, in a chronic ileitis, death occurred before operative interference. Two perforations of the terminal ileum were observed with localized abscess and peritonitis.

### HEMORRHAGES

Gross hemorrhage was noted in 11 instances in this series usually in the course of the disease occasionally (twice) as an initial symptom. The hemorrhage occurs usually in

the form of melena twice hematemesia was noted Pollis quotes a case with repeated profuse and massive hemorrhage passed by the bowel (melena). Two previous examinations had been negative. Eight transfusions were required to control shock. Eventually, at operation, 27 inches of ileum were resected with cure. Galambos and Mittelman cite a case of atypical terminal ileitis in which gastric hemorrhage and melena dominated the clinical picture and led to the presumptive diagnosis of a peptic ulcer. Yumich and Crohn reported the case of a 55 year old male in which the onset of symptoms was ushered in by two massive intestinal hemorrhages in the form of melena producing syncope and collapse. The second hemorrhage was almost fatal.

It is difficult to explain hematemesia when it occurs. In the 2 cases in which such a phenomenon was seen in this series, the diagnosis of regional ileitis was confirmed by exploratory operation. These both were cases in which the terminal ileum was involved no higher segment of ileum or jejunum being implicated.

#### GENERAL COMPLICATIONS

In this series of cases of ileitis duodenal ulcer was noted five times, cholelithiasis three times and acute cholecystitis was encountered once during an operation for terminal ileitis. Regional enteritis involving a Meckel's diverticulum has been observed (Horn and Rhoads) in one case with perforation at the base of the sac. In my own series, I have once seen Meckel's diverticulum involved in a skip lesion. In another case a Meckel's diverticulum causing low intestinal obstruction by torsion was clinically preoperatively confused as a case of ileitis. Patricelli also mentions a case of regional ileitis with Meckel's diverticulitis with fatal outcome.

Some general complications which occur are actually incidental and not related to the disease. Thus this series includes 4 cases with chronic rheumatic heart disease, one

with a decompensated right heart on the basis of generalized atherosclerosis. Occasional mention is made of joint pains and arthritis in a case showing spondylitis involving the cervical spine. In an unusual case, active pulmonary tuberculosis was present, though the resected ileitis was characteristically of the nonspecific granulomatous type, in this case, too, diffuse involvement of the large joints was noted. Infantilism and delayed sexual maturity has already been mentioned. One case observed had had a large retroperitoneal or intra abdominal abscess with consequent obstruction of the inferior vena cava.

Tetany in regional ileitis is uncommon, occurring more frequently in those cases of diffuse disease involving the upper ileum and the jejunum. One case following two resections of the small bowel for recurrent ileitis developed typical carpopedal spasm. Blood calcium in this case was 6.7 milligrams per cent. Immediate relief was noted upon injection of calcium gluconate and control of the diarrhea. In another case, typical tetanic convulsions were noted with a blood calcium of 3.4 milligrams per cent.

Amyloidosis does not occur in any of the cases in this series in spite of the long duration and chronicity of the disease. This is due probably to the nonpyogenic nature of the discharge from surface fistulas or in the intestinal flux. However, I am privileged to mention one case of generalized amyloidosis which occurred as a sequel to or associated with regional ileitis observed and studied at autopsy (Mount Sinai Hospital New York City, courtesy of Dr. Hilliard Cohen) \*.

Primary amyloidosis of the alimentary tract has been reported by Golden in a Negro observed for 16 years who presenting symptom was hematemesis, dyspnea and ortho-

In the recent article by Olman and Sussman 3 cases of amyloidosis associated with or secondary to regional ileitis are recorded. The amyloid involves various viscera throughout the body and is a prominent feature only in the one case which is identical with that about to be published by Cohen and Fishman (personal communication).

pnea. At autopsy, the stomach, duodenum and terminal ileum were all involved, the liver, spleen, kidneys and adrenal were free of amyloid. This however, is different from true primary regional ileitis, in which the systemic amyloidosis is a sequel or a complication. In primary systemic amyloidosis, the gastrointestinal tract may be involved as may any other organ. Involvement of the arterioles and of the stroma occurs frequently in the intestine. However, systemic amyloidosis secondary to ileitis or non-specific ulcerative colitis is extremely rare (Koletsky and Stricker, Moscheowitz).

Gaucher's disease was seen in a case of ileitis as an incidental finding. Pernicious anemia was seen in one case in this series (serial No. 73) in a man 54 years of age who had undergone a short circuiting operation for regional ileitis followed promptly by a recurrence of diarrheal symptoms. At a second operation the original site of the disease, the terminal ileum and the ascending colon were resected but no recurrence of the disease process in the new proximal ileum was noted. An unusually severe anemia and emaciation followed this operation. Two years later the blood count revealed a true primary pernicious anemia. Hemoglobin 26 per cent, RBC 1,400,000, WBC 2,650 cells, myelocytes one normoblast, marked anisocytosis and poikilocytosis, hematocrit 10.8 per cent, total proteins 4.6 per cent. Sternal marrow puncture 38 per cent myeloblasts. The administration of liver extract was followed by a dramatic response of reticulocytes to 30 per cent. On discharge the hemoglobin was 60 per cent, total proteins 6.1 Gm per cent.

In spite of the severe suppurative intra-abdominal complications which accompany walled off perforations in ileitis, pyelophlebitis and multiple liver abscesses are rarely seen. The report of Snively is the only one to be found in the literature. A colored male suffered with diarrhea, high fever, ascites was probably present. On one occasion only, cysts of *Amoeba histolytica* had been found. At examination, regional ileitis involving 5 feet of ileum was noted, a

fistulous communication between terminal ileum and the sigmoid was present, as also were pyelephlebitis and multiple small liver abscesses. No such complication of ileitis is seen in this series probably for the reason that almost all cases are instances of granulomatous rather than of suppurative disease. Warren and Sommers quote an additional case in a Negress in which autopsy disclosed thrombosis of the inferior vena cava and external iliac veins following upon intestinal obstruction and complicated by numerous internal intestinal fistulas. Intussusception is not mentioned in the literature as a complication of ileitis.

### GYNÉCOLOGIC MANIFESTATIONS

There is little disturbance of endocrine function in the female as a result of ileitis. Menstruation is not affected even in the severe cases. This is in contrast to ulcerative colitis in which amenorrhea is a frequent manifestation of the severe and toxic phase of the disease, the return of the menses usually indicates the improved prognosis and the beginning of a remission.

Fertility is unaffected by ileitis, pregnancy having been noted frequently in this series particularly during the remissions. The relationship of pregnancy to ileitis may be represented in table 9.

TABLE 9—*Relationship of Ileitis and Pregnancy*

Onset of Ileitis during Pregnancy	1 case
Onset of Ileitis after childbirth	
(a) immediately	1 case
(b) two months	2 cases
(c) eight months	1 case
(d) three years	1 case
Ileitis pregnancy still births (11 & 3rd week)	2 cases
Ileitis successful surgical cure subsequent successful pregnancy and normal childbirth	1 case

It is interesting to observe that one of these patients had two healthy babies after a successful short circuiting operation for ileitis (ileo transverse colostomy). It is noteworthy that a recto vaginal fistula which had been previously surgically resected in this case remained closed in spite of the two deliveries.

The ileitis case seems to tolerate pregnancy much better than do those mothers suffering from ulcerative colitis. In colitis, the pregnancy itself is uneventful and the diarrhea and active symptoms of the disease are likely to undergo a remission. But a most active and severe recurrence of all the symptoms is likely to recur within a few weeks after delivery. In contrast, we note in ileitis one case in which the disease was observed in its initial phase during pregnancy followed by normal delivery, 3 cases with successful delivery, the mothers remaining well thereafter (table 9). Five cases are further seen in which the ileitis recurred after the birth of the child in some of them the recurrence occurred so late thereafter that there is little reason to attribute the recurrence of the disease to the past pregnancy or parturition. Bibson cited a case of terminal ileitis with obstruction and with an abscess complicating the pregnancy and resulting in abortion. Quite recently Raffensberger described the case of a 22 year old woman upon whom previously a short circuiting procedure had been performed for a regional ileitis. She remained well and free of symptoms even during an uneventful pregnancy which unfortunately, terminated in a premature delivery of a stillborn child. A few days later, severe diarrhea and abdominal pain occurred, caused as demonstrated radiographically by an acute recurrence of the ileitis in the loop proximal to the previous anastomosis. Wilson Gruman and Ashburn mention profuse vaginal discharge in one case after resection of the terminal ileum the vaginal discharge ceased almost at once possibly a fistula to the uterine cavity or vagina had existed. In one case observed a right salpingitis occurred as a result of the fistulous tract leading from the terminal ileum to the right fallopian tube. In another case the fistulous tract from the terminal ileum ended in the vault of the vagina constituting a profuse suppurative open fistula successfully repaired by surgical intervention (personal communication Dr. Ralph Colp).

## 8 Roentgenographic Study of Regional Ileitis

---

THE LITERATURE on the radiographic recognition of terminal ileitis more or less recognizes the classic description by Kantor of the "string sign" as the characteristic manifestation of the disease. Not only was it the first radiologic study of ileitis, but, by its descriptive accuracy and terse and apt nomenclature, it has gained general and permanent acceptance. Crane had previously used the word "string sign" to designate the attenuated thinned out segments of colon identified in spastic or irritable bowel. The word and the description were so apt that Kantor borrowed the name to apply it to the similar appearance which occurred in the ileum in terminal ileitis.

The string sign is a "thin irregular linear shadow suggesting a cotton string in appearance, and extending more or less continuously from the region of the last visualized loop of ileum through the entire extent of the filling defect and ending at the ileo cecal valves" (figs 27-30).

The explanation of the "string sign" is to be found in the study of an opened specimen of old chronic terminal ileitis. The thickening of the walls of the gut by inflammation and scar tissue infiltration encroach upon the lumen to narrow it. The parallel process of cicatrization contracts or shrinks down the walls further to diminish the circumference of the intestine. If the normal proximal lumen is 6 cm in diameter that of the diseased terminal ileum may be 0.7 to 1.0 or 1.2 cm in diameter throughout the area

indicated by the string sign. Reconstruct such a specimen, as has been done, fill it with barium and take a roentgen film and the string sign is fully reproduced (figs 31-34).

Besides the "string sign" one observes in radiography



Figs 27-29-30—Various Types of Terminal Ileitis with Characteristic String Sign Formations

Fig 27

of ileitis an indentation on the inner wall of the cecum at the insertion of the ileocecal valve. This indentation is regarded as due to the edema and inflammatory swelling of the terminal segment of ileum just at its approach to or abutment upon the inner surface of the valve of Bauhin.



The cecum itself may be consistently spastic and irregular and frequently will be seen to be conical or persistently pointed in its shape. This appearance often suggests an amebic granuloma of the cecum, a frequent site for ame-



Fig. 28—(See Fig. 27)

bias implantation. Strombeck in acute ileitis, noted pronounced swelling of the valve of Bauhin, the two lips of which may be conical and may bulge into the cecum. He further claims that he was able to identify elongated ulcers 1-2 cm. in the long axis and somewhat rounded ulcers along the attachment of the mesentery; these ulcers occurring exactly at the point of passage of the intestinal vessels

through the muscularis. A soft shadow in the mucosal relief can often be observed at the site of an abscess.

In the upper ileum proximal to the diseased terminal segment ileitis produces alternate areas of constriction and dilatations corresponding to pathologic "skip lesions" and to 'skipped areas,' of normal mucosa. The dilatation



Fig. 9—(See Pl. 97.)

represents areas of normal mucosa lying between the constricted diseased 'skip lesions' of inflammatory disease. This disturbance of pattern may be followed upward through the ileum in advanced cases and even into the jejunum.

It must not be presumed that the 'string sign' of ileitis is present in all cases. In many instances, only a distorted mucosal pattern of the terminal ileum may be noted (Last) (fig. 35). The normal folds of ileum run

The cecum itself may be consistently spastic and irregular and frequently will be seen to be conical or persistently pointed in its shape. This appearance often suggests an amebic granuloma of the cecum, a frequent site for amo-




Fig. 28.—(See Fig. 27.)

biasis implantation. Strombeck in acute ileitis, noted pronounced swelling of the valve of Bauhin, the two lips of which may be conical and may bulge into the cecum. He further claims that he was able to identify elongated ulcers 1-2 cm. in the long axis and somewhat rounded ulcers along the attachment of the mesentery; these ulcers occurring exactly at the point of passage of the intestinal vessels

students disarrangements of the mucosal relief that were very similar to those produced by mild disease or deficiency states. Since this dislocation is often difficult to read and to translate, the purely radiologic diagnosis must rely upon



Fig. 31 (31-31)—Various Types of Regional Ileitis Involving one and two Loops of Distal Ileum. Disturbed mucosal pattern, tendency to polypoid form, an indication of more extensive terminal sign.

Fig. 31

concomitant clinical facts. Kirklin in a discussion of a paper by Brown, emphasized this point: he says that shortening and rigidity of the segment and the string sign are late manifestations of the disease. The earlier diag-

longitudinally and parallel crossed by the markings of the transverse folds of the intestine, thus, a fine network of mucosal folds are normally visible. When the area is in early stage of disease, when edema and active inflammation



Fig. 30—(See Fig. 27.)

is taking place, but before cicatricial narrowing has intervened, the so called 'mucosal type,' the distortion of the mucosal pattern is the only radiographic sign that may be interpretable and acceptable (fig. 36). Ruffin et al warn against interpreting slight changes in the mucosal arrangement as being due to disease or to deficiency patterns. They were able to note, in control observations upon normal

above and below the lesion show relative hypomotility and delay, the mucosal changes may be minimal. Weber further warns against misinterpretation of the x rays because of wide variability of the normal mucosal pattern



Fig. 33—(See Fig. 32)

The examination demands close scrutiny of each individual segment by multiple roentgenoscopic observations of the descending opaque meal and of the terminal ileum after it has been distended by the reflux of the opaque enema.

In many cases radiologic evidence of ileitis is completely missing and yet the clinical data are so convincing that the diagnosis must be adhered to in the absence of confirmatory

nosis must be based upon inner changes in the relief pattern of the mucosa, a sense of diminished pliability and motility and local or general intestinal hypermotility. The greater the deformity the older and more advanced the lesion.



Fig. 32—(See Fig. 31.)

Weber describes the radiologic picture of the early relatively unpronounced degree of submucosal infiltration as a flattening of the normally high mucosal relief pattern. If ulceration has taken place the relief pattern is jagged and irregular, not smooth and flat. In the advanced stages, the affected segment of intestine is shortened, its lumen narrowed, motility and pliability are lost. The affected segment shows relative hypermotility; the unaffected loops

above and below the lesion show relative hypomotility and delay the sigmoid changes may be minimal. Weber further warns against misinterpretation of the x rays because of wide variability of the normal mucosal pattern.



Fig. 33 — (See Fig. 32.)

The examination demands close scrutiny of each individual segment by multiple roentgenoscopic observations of the descending opaque meal and of the terminal ileum after it has been distended by the reflux of the opaque enema.

In many cases radiologic evidence of ileitis is completely missing and yet the clinical data are so convincing that the diagnosis must be adhered to in the absence of confirmatory



nosis must be based upon inner changes in the relief pattern of the mucosa, a sense of diminished pliability and motility and local or general intestinal hypermotility. The greater the deformity the older and more advanced the lesion.



Fig. 32—(See Fig. 31.)

Weber describes the radiologic picture of the early relatively unpronounced degree of submucosal infiltration as a flattening of the normally high mucosal relief pattern. If ulceration has taken place the relief pattern is jagged and irregular, not smooth and flat. In the advanced stages, the affected segment of intestine is shortened its lumen narrowed motility and pliability are lost. The affected segment shows relative hypermotility, the unaffected loops

originating from the wall of the ileum, frequently trending downward toward the pelvis or toward the urinary bladder (fig 18). These appear like small spikes, only an occasional one traverses large distance to form a true elongated fistula.

The upper "skip lesions" of disease and skipped areas of healthy bowel are usually easily picked out on the films.



Fig. 33.—Characteristic Disturbance of Mucosal Pattern in Terminal Ileitis. Courtesy of Dr. Franz Lust.

It is remarkable how accurate reading of such films corresponds to the locality of the resected specimen. One can ascertain almost to the inch the extent of the lesion as expressed in the x-ray films and as confirmed at the operating table.

Involvement of the cecum or of the other segments of colon is similarly to be sought after since combined instances of ileitis and colitis are not uncommon.

roentgen signs (Yumich and Crohn) The "string sign" is almost pathognomonic, is always highly suggestive, on the other hand, the absence of all roentgenographic evidence does not in any manner preclude a positive diagnosis of ileitis



Fig. 34—(See Fig. 31)

In a very careful scrutiny of films it is often possible to discern evidences of fistula formation. These fistulas appear as fine lines, attenuated streaks traversing, from terminal ileum, to cecum or to transverse colon to sigmoid or to pelvic colon. These lines may be seen only on fine study but should always be sought. With a magnifying glass it is often possible to see several small potential fistulas

of filling the proximal part of the intestine with a substance which will pass through the partially obstructed lumen with difficulty." The radiography of the colon being negative, insufficiency of the ileocecal valve occurs in a sufficient majority of instances (probably 80 per cent) to allow regurgitation of the barium clyster into the terminal loop of ileum. By this means the string sign or irregularity of the mucosal pattern in the terminal ileum is usually visible. Unfortunately, in a fair percentage of cases, redundant loops of colon or of a pelvic colon with a long mesentery may overlie the terminal ileum to becloud or eclipse the visualization of that segment. For that reason the barium enema must often be supplemented by the barium meal. The barium meal offers the additional opportunity of observing the upper ileum and jejunum and allows for a better visualization of sinus and fistula tracts before the overloaded and redundant colon fills and obstructs the reading of the finer structures. The barium meal, consisting of 4 ounces of barium in a flavored water vehicle, must be followed hour by hour in its course through the jejunum and the ileum, particular attention being given to that phase when the terminal ileum is filling or is filled with the opaque mixture. The films are taken at the optimum time as determined by fluoroscopy. The mistakes and the omissions of diagnosis are often due to the arbitrary standards for plate taking and timing as set in some laboratories. Those techniques that call arbitrarily for 3, 6 and 9 hour plates are likely to miss the optimum time for visualization of the terminal ileum. The terminal ileum may fill best in 3 hours or in 5 hours or 8 or 12 hours depending on conditions of individual intestinal motility, in cases with obstruction the delay may be long but the ultimate satisfaction in visualizing the disease in the terminal ileum will be rewarding. The barium meal further identifies the higher skip lesions and skipped areas the dilated proximal segments of small bowel indicating more distal obstruction of the lumen of the bowel. It affords the basis for the

## THE TECHNIC OF RADIOGRAPHY

The string sign can usually be demonstrated equally well by the barium enema and by the barium meal. The first preference in working up a case is for the barium enema since this method delimits the colon, indicates whether universal or segmental colitis is present or absent, it is




Fig. 36—Terminal Ileitis, Mucosal Type. Note absence of the string sign, irregular serrated margin of ileum, disturbance of mucosal pattern.

most important in a case of diarrhea primarily to eliminate the probability of ulcerative colitis, diverticulitis, or of neoplasm. Weber shows a distinct preference for the barium enema as the initial procedure in the radiologic diagnosis of ileitis. With the contrast enema the coils of the ileum are well elevated out of the pelvis, are distended and are readily manipulated under roentgenoscopic control by palpation through the abdominal wall. Furthermore, one must not overlook the danger

The string sign so characteristic of ileitis may be simulated by other diseases. Hyperplastic ileo cecal tuberculosis or ileal tuberculosis, primary or secondary, may imitate the string sign as observed by Stierlin and Chroul and more recently accentuated by Taylor. Connell has noted the



Fig. 37—Demonstration of Barium Iod Saline Methyl for Early Visualization of the Intestinal Tract. (Courtesy of Weintraub and Williams)

string sign to be present consistently in ileo cecal tuberculosis. He notes irregularity of the shadow in the terminal ileum and conical contraction of the cecum. The ileo cecal valve is visible to an exaggerated degree. The lips are gaping through the valve and occasional obstruction at this point may occur. The additional involvement of the cecum more likely invites the diagnosis of tuberculosis.

differential diagnosis of regional ileitis and jejunitis from the somewhat similar and easily confused picture of sprue, of vitamin deficiency manifestations, of intestinal Hodgkin's disease or of lymphosarcoma. It allows further for the minute study of distal fistula formation from loop to loop of small or of large bowel.

The novel suggestion of Weintraub and Williams of administering the barium meal followed soon thereafter by the drinking of an iced saline solution affords a rapid and simplified means of quickly filling the distal loops of the small bowel. The barium meal is very promptly propelled by peristaltic activity so that within a half hour almost all of the loops of jejunum and ileum are fully visualized. Even the mucosal pattern of the distal loops are well defined. Of 235 normal control cases, in 190 the head of the barium saline meal had reached the cecum in one half hour or less. The mucosal pattern was in all cases as good as that obtained on routine small bowel examination, especially in the terminal ileum. If the method of Weintraub is proven to be free of faults, and artefacts are not created by the rapid propulsion of the barium the newer technique may materially simplify and accelerate the radiographic recognition of diseased distal areas of intestinal involvement (fig. 37).

Oral intubation of the small intestine by means of a duodenal tube or a Miller Abbott tube, has been employed by some authors. The tube or balloon is passed to the required depth as determined by fluoroscopy, then barium suspension is injected through the tube to the probable site of the disease. This allows for intimate and exact study of a localized segment of intestine free from the overlapping shadows of extraneous loops of bowel. The method is good and rewarding though its use is not very universal. Actually the barium meal is so successful when properly administered and interpreted that the more devious and difficult and time consuming method of intestinal intubation is rarely called upon or utilized.

The string sign so characteristic of ileitis may be simulated by other diseases. Hyperplastic ileocecal tuberculosis or ileal tuberculosis, primary or secondary, may imitate the string sign as observed by Stierlin and Chroul and more recently accentuated by Taylor. Connell has noted the



FIG. 37.—Demonstration of Barium Iod Saline Method for Early Visualization of the Intestinal Tract. Courtesy of Weintraub and Williams.

"string sign" to be present consistently in ileocecal tuberculosis. He notes irregularity of the shadow in the terminal ileum and conical contraction of the cecum. The ileocecal valve is visible to an exaggerated degree. The lips are gaping through the valve and occasional obstruction at this point may occur. The additional involvement of the cecum more likely invites the diagnosis of tuberculosis.



differential diagnosis of regional ileitis and jejunitis from the somewhat similar and easily confused picture of sprue, of vitamin deficiency manifestations, of intestinal Hodgkin's disease or of lymphosarcoma. It allows further for the minute study of distal fistula formation from loop to loop of small or of large bowel.

The novel suggestion of Weintraub and Williams of administering the barium meal followed soon thereafter by the drinking of an iced saline solution affords a rapid and simplified means of quickly filling the distal loops of the small bowel. The barium meal is very promptly propelled by peristaltic activity so that within a half hour almost all of the loops of jejunum and ileum are fully visualized. Even the mucosal pattern of the distal loops are well defined. Of 235 normal control cases, in 190 the head of the barium saline meal had reached the cecum in one half hour or less. The mucosal pattern was in all cases as good as that obtained on routine small bowel examination, especially in the terminal ileum. If the method of Weintraub is proven to be free of faults and artifacts are not created by the rapid propulsion of the barium the newer technic may materially simplify and accelerate the radiographic recognition of diseased distal areas of intestinal involvement (fig. 37).

Oral intubation of the small intestine by means of a duodenal tube or a Miller Abbott tube, has been employed by some authors. The tube or balloon is passed to the required depth as determined by fluoroscopy, then barium suspension is injected through the tube to the probable site of the disease. This allows for intimate and exact study of a localized segment of intestine free from the overlying shadows of extraneous loops of bowel. The method is good and rewarding though its use is not very universal. Actually the barium meal is so successful when properly administered and interpreted that the more devious and difficult and time consuming method of intestinal intubation is rarely called upon or utilized.

## 9 The Diagnosis of Regional Ileitis

---

THE DIAGNOSIS of regional ileitis is essentially a clinical one, based upon the characteristic subjective symptoms and the physical findings. The x ray diagnosis is valuable as confirmatory evidence and allows for anatomic interpretation of the lesion. It is a grave error to think of ileitis as an obscure disease difficult or impossible of diagnosis. Ileitis has clear cut subjective and physical features, which when combined spell out a succinct clinical diagnosis. The combination of diarrhea, fever, anemia and fistulas, internal or external with or without an abdominal mass with or without radiographic confirmation means a possible or a probable diagnosis of ileitis. The negative sigmoidoscopy rules out the usual case of ulcerative colitis 90 per cent of which cases give a distinct pathologic sigmoidoscopic picture. The remaining possibility is a right sided segmental colitis to be differentiated from regional ileitis. Segmental colitis may have and does have anal complications and rectal fistulas but has no fistulas to the abdominal wall and no internal fistulas or intra abdominal masses.

To recapitulate the diagnostic significance of the symptoms and signs of ileitis:

The diarrhea is the most common feature, it may be absent in 10 per cent of the cases.

Fever is frequent and significant but the majority of the long standing low grade cases are afebrile.

Anemia may be so moderate as to be insignificant, a hemoglobin reading below 65-70 per cent is of value

but not necessarily so since combined ileitis and colitis or ileo colitis of a nonspecific granulomatous type must be also considered

Carcinoid tumors of the terminal ileum, endometriosis with implantations on the serosal surface of the ileum and occasionally benign new growths within the lumen of the ileum may stimulate the string sign of ileitis particularly when obstructive phenomena are also present. In rare instances, small round transparencies within the barium filled small intestine indicate the presence of polyps, these are the accompanying manifestations of ileitis, just as they indicate in ulcerative colitis an advanced stage of inflammatory reaction with secondary polyposis. In cases of ileitis with mass formation the loops of upper ileum may be displaced laterally by the mass leaving a void area corresponding to the inflammatory tumor formation.

Rosenberg mentions enterogenous cysts at the ileo cecal junction as simulating the roentgen picture of terminal ileitis.

The positive roentgenographic evidences of ileitis are usually present in a well defined clinical case. However, it must always be remembered that all confirmatory roentgenologic evidence may be missing in a subsequently clinically proven case. The roentgen limitations in atypical or early ileitis become self evident with increasing experience (Yumich and Crohn).

culous pathogenicity. In a recent survey of anal fistulas, summarizing 600 cases, Jackman and Buie found 87.8 per cent negative for tuberculosis. In the true tuberculous fistulas, primary affects elsewhere in the body, usually the lung, were always demonstrable. This reversal in the view point of the relationship of tuberculosis to anal fistulas is due in large part to the increasing role which ileitis and colitis play as etiologic factors in the incidence of such perirectal complications.

It is particularly important to have sufficient mental fortitude to sustain the diagnosis of at least probable ileitis, even though many significant clinical features are missing and the x-ray diagnosis is negative. On several occasions one has been saved from a blatant error by including in the provisional negative diagnosis at least the possibility of ileitis. The following experience is illustrative.

The case was reported as nervous diarrhea, irritable colon, but the proviso was added of a possible ileitis. Five years later the symptoms flared up and now exploratory laparotomy showed regional ileitis with an ileo sigmoid fistula. Not that every case of diarrhea should be classed as a possible ileitis, but the association of diarrhea with a fistula, a mass, fever, or anemia in any such combination calls for diagnostic caution and acuity.

The low grade fever of regional ileitis is often overlooked by general internists who are likely to pay little attention to mild diarrhea, or to the complaints of occasional front abdominal pain. The patient may not consider two or three loose bowel movements as diarrhea, the abdominal pain may be minimal and no positive physical signs may be discoverable. These cases of ileitis with low grade fever are usually confused with brucellosis or undulant fever particularly when a low titer agglutination against the *Melitensis* organism has been reported. Many a case of ileitis has been overlooked because of a positive agglutination of 1:40 or 1:80 against brucella. One such case was

Loss of weight is significant but may not be extensive. An abdominal mass may be present, but most frequently is absent.

Fistulas, internal, external and perirectal are the most nearly pathognomonic manifestations of the disease, particularly when associated with diarrhea and with the other positive signs and findings of ileitis.

The radiographic study is exceedingly important when positive, a negative finding does not necessarily nullify a clinical tentative diagnosis of ileitis. Any one symptom, especially any combination of two or more of these signs and symptoms should suggest the consideration of ileitis. Diarrhea alone, diarrhea plus fever, diarrhea with fistulas, an abdominal mass with fever, an intestinal or perirectal fistula, any combination of such symptoms and physical findings is suggestive. An illustrative case in point is worth quoting because of its significance.

*P B* (Serial No 197) Male age 39 years. For many years he had suffered with mild diarrhea. Eight years ago he had been operated upon for a rectal fistula. Subsequently he had lost 50 pounds of weight, complained of abdominal cramps and persistent diarrheal movement. The physical examination was negative except for slight tenderness over the cecal area of the abdomen and suggestive clubbing of the fingers. The rectal and sigmoidoscopic examination were negative. The radiographic examination—barium enema and barium meal—were reported as negative except for some evidence of a spastic colon. The terminal ileum was well visualized, appears somewhat irritable in its distal portion but fills well and is distensible. The report to the referring physician stated that although the diarrhea, fever and the perirectal fistula strongly suggested ileitis, the absence of confirmatory radiographic evidence made such a positive diagnosis impossible. It was urged that the patient be carefully observed for future development. Six months later the patient developed a fistula from the ileum to the sigmoid. At operation a regional ileitis was discovered involving 5 feet of ileum proximal to the ileocecal valve. An ileotransverse colectomy with transection of the ileum resulted in cure.

The significance of the perianal fistulas cannot be overlooked. In former years such fistulas were regularly regarded as either being tuberculous in nature or were suspected of a tuberculous origin. Mummery in 1927 regarded at least 15-20 per cent of perirectal fistulas as of tuber-

culous pathogenicity. In a recent survey of anal fistulas, summarizing 600 cases, Jackman and Buie found 87.8 per cent negative for tuberculosis. In the true tuberculous fistulas, primary affects elsewhere in the body, usually the lungs, were always demonstrable. This reversal in the view point of the relationship of tuberculosis to anal fistulas is due in large part to the increasing role which ileitis and colitis play as etiologic factors in the incidence of such perirectal complications.

It is particularly important to have sufficient mental fortitude to sustain the diagnosis of at least probable ileitis, even though many significant clinical features are missing and the x-ray diagnosis is negative. On several occasions one has been saved from a blatant error by including in the provisional negative diagnosis at least the possibility of ileitis. The following experience is illustrative.

The case was reported as nervous diarrhea irritable colon, but the proviso was added of a possible ileitis. Five years later the symptoms flared up and now exploratory laparotomy showed regional ileitis with an ileosigmoid fistula. Not that every case of diarrhea should be classed as a possible ileitis, but the association of diarrhea with a fistula, a mass fever, or anemia, any such combination calls for diagnostic caution and acuity.

The low grade fever of regional ileitis is often overlooked by general internists who are likely to pay little attention to mild diarrhea or to the complaints of occasional faint abdominal pain. The patient may not consider two or three loose bowel movements as diarrhea; the abdominal pain may be minimal and no positive physical signs may be discoverable. These cases of ileitis with low grade fever are usually confused with brucellosis or undulant fever particularly when a low titer agglutination against the *Melitensis* organism has been reported. Many a case of ileitis has been overlooked because of a positive agglutination of 1:40 or 1:80 against brucella. One such case was

even published in medical literature in this country (Sproull) as a case of undulant fever with unusual intestinal changes, this, before the first description of ileitis as a clinical entity. Years later, exploratory operation and resection of the ileum revealed the true nature of the disease as one of ileitis with low grade fever, the doubtful low grade agglutination titer against brucella having only confused the diagnostic issue.

Ileitis which presents itself as a case of continuous low grade fever, with low leukocyte count, with occasional joint involvement, clubbing of fingers or eye complications, particularly when a heart murmur is present, is likely to be overlooked, particularly when the minimal diarrhea, which is the actual key to the diagnosis, is ignored. All of these manifestations of a generalized disease are common to both ulcerative colitis and to ileitis, omit the diarrhea and the general symptoms resemble brucellosis, rheumatic fever, subacute bacterial endocarditis, lupus erythematosus disseminatus or Libman Sachs disease. Rosenblate, Goldsmith and Strauss cite a case of regional ileitis unrecognized for four years, during which time, because of a soft systolic mitral murmur over the precordium, the case had been considered as one of chronic heart disease or subacute bacterial endocarditis, in spite of reported negative blood cultures. Sprague et al report a case of long standing fever in a girl of 15 years of age who presented the clinical picture of an irregular fever with temperatures ranging between 101 and 104 F with polyarticular rheumatic like arthritic manifestations and perirectal abscesses, the whole clinical picture being due to a regional ileocolitis.

The significance of the cardiac murmur may be overemphasized. At least 5 cases in this series had had rheumatic fever and bore residual systolic mitral or aortic murmurs. Add to the old rheumatic cardiac murmur a low grade fever and the possibility of ileitis or segmental colitis is rarely considered. This is all the more important because in one case in this series an acute endocarditis

actually occurred as a complication while the patient was under observation. A young woman with recognized regional ileitis was under observation preparatory to operation. An increased degree of fever was associated with the development of a systolic murmur at the cardiac apex; the murmur was heard with daily increasing intensity, a low rough murmur was audible and was transmitted to the mitral area. A blood culture was negative at the height of the fever. Penicillin was instituted, the fever subsided in even days and the cardiac murmur ceased to be audible. The patient was later successfully operated upon for regional ileitis. No further cardiac manifestations occurred.

The larger and sometimes the smaller joints of the hands or feet may be implicated in diarrheal diseases, the so-called 'arthritis dysenterique', but the significance of the arthritis as an intestinal manifestation is usually overlooked in favor of a diagnosis of rheumatic fever with joint involvement.

The mild or moderate eosinophilia that frequently accompanies ileitis and colitis may suggest and be confused with periarteritis nodosa.

The eye complications which occur occasionally in ileitis but more commonly in ulcerative colitis again are usually overlooked and are classified in the tuberculous or rheumatic category (Sprague et al). Ophthalmologists of wide experience are conscious of the fact that inflammatory lesions of the eyes are not uncommon complications of intestinal diseases. Iritis, iridocyclitis, keratitis, corneal ulceration, phlyctenular conjunctivitis all constitute focal manifestations of intestinal diseases.

#### DIFFERENTIAL DIAGNOSIS

When a suggestive 'string sign' is present, or when a disarrangement of the pattern of the intestinal mucosa is noticeable in the roentgenograms various other possible intra-abdominal diseases come into consideration in the differential diagnosis. When the upper ileum and the



jejunum show puddling and segmental dilations and disturbed mucosal patterns, sprue and the vitamin deficiency diseases deserve close notice. Here the differentiation between sprue and ileo jejunitis is often difficult (fig 38)

Intra abdominal Hodgkin's disease by its fever and low leukocyte count and its possible intestinal involvement may



Fig 38—Diffuse Ileitis. Whole Ileum Involved. Scattered skip lesions. fistulous tract in lower ileum between loops of small bowel constipation. Patient preserved his well being

simulate ileitis. Multicentric intestinal lymphosarcoma with its characteristic areas of intestinal dilatation, with hemorrhagic fever and perforation and rapid wasting may be confused with ileitis. Both Hodgkin's disease and intestinal sarcomatosis are intensely malignant and progressive diseases with early fatal issue in contrast to the more mild and low grade continuous course of chronic regional ileitis.

The "string sign" of the terminal ileum may be simulated by (figs. 39 and 40) hyperplastic ileocecal tuberculosis, carcinoid of the terminal ileum, endometriosis of the terminal ileum, enterogenous cysts about the terminal ileum, actinomycosis, intestinal



Fig. 39—Anemic Di enteri. Characteristic Deformity of the Cecum Incomplete Filling of Terminal Ileum. This radiograph is almost identical with that seen in nonspecific granulomatous ileo-coitis.

Primary intestinal tuberculosis as a manifestation of bovine infection is exceedingly rare. In a complete survey of all the postmortem and surgical pathologic material at the Mount Sinai Hospital over the last fifteen years, only 4 cases of primary intestinal tuberculosis survived a critical analysis (Crohn and Yarnis). In 2 of the these cases,

the ileum alone was the site of typical milium tuberculosis, tubercle bacilli were found in the sections of the histologic studies. All other cases were those of ileo cecal or ileal tuberculosis secondary to open lesions in the pulmonary areas with caseating cavity formation and positive sputum or positive gastric residue findings for tubercle bacilli.



Fig 40—Appearance of Appendix Filled with Barium Resembling String Sign of Ileitis

Cases of regional ileitis regularly have negative chest findings by physical examination and by radiographic study. Cases of secondary intestinal tuberculosis have strongly positive chest plates with evidence of cavity formation and with positive sputum findings. That intestinal tuberculosis can simulate exactly the string sign of ileitis has been amply demonstrated (Crohn and Yarnis). This point has again been recently emphasized by Connell Blackburn.

found the Mantoux Test negative in 19 of the 22 cases and x-rays of the chest uniformly negative in regional ileitis.

Argentophilic carcinoid tumors infiltrating along the mucosa of the terminal ileum represent a favorite site for that lesion and can simulate the clinical picture and the radiographic string sign of ileitis. In one case in my experience, the terminal ileum was resected under the supposition that it represented a case of ileitis; the gross specimen was passed from hand to hand and was accepted as one of ileitis by many long experienced in handling such surgically resected specimens. Only on histologic study was the true character of the lesion identified as one of carcinoid tumors of the terminal ileum with metastases to the mesenteric lymph nodes.

Up to 1939 237 cases of argentaffine tumors of the small intestine had been reported in the literature, obstruction occurred in 24.4 per cent of these. Pennington and Priestly report a case of carcinoid tumors of the terminal portion of the ileum invading the mesentery. The abdominal mass involved several loops of the terminal segments of the ileum. The lesion was malignant, several metastatic tumors being present in the liver. McLeod adds 2 cases to 283 reports in the literature (1944); intestinal obstruction in the terminal ileum and/or in metastases were commonly observed. Dockerty and Ashburn in a review of the literature describe carcinoid tumors (so called) of the ileum. They detail 30 cases of carcinoid of the terminal ileum, 18 of which had metastasized locally or distally; small orange submucosal nodules were arranged linearly with a minimal degree of ulceration. Regional lymph nodes were involved in 11 cases; in 5 metastases to the liver were present, suggesting a low grade type of carcinoma with low virulence.

Endometrial transplants on the serosa of the terminal ileum with or without invasion of the intestinal walls occasionally may be confused with ileitis; the string sign is almost identical. The intestinal bleeding with the menses may, if present, suggest the true nature of the lesion. A

case of endometriosis as a cause of intestinal obstruction has been described (McGuff et al), in which the segment 5 cm proximal to the ileo cecal valve was involved by endometrial transplants. The clinical picture resembled in general a case of regional ileitis with intestinal obstruction.

Enterogenous cysts about the ileo cecal angle are exceedingly rare, but may simulate roentgenographically the constriction of terminal ileitis (Rosenberg).

Actinomycosis, intestinal, with abdominal wall fistula may simulate ileitis. The finding of the characteristic ray fungus in the discharges from the fistula will save much confusion and often embarrassment. The possible confusion between ileitis and sarcoidosis has already been discussed (page 16). Sarcoidosis of Boeck involving the small intestine as a granulomatous lesion was described recently by Watson and his associates. The identity of ileitis and sarcoidosis was even suggested by Hadfield. Snapper has, however, made the point that he has never encountered a case of sarcoidosis with a lesion of the intestine, nor have his many cases of ileitis ever shown sarcoidosis. Blackburn states that in ileitis there is no record of swelling of the fingers and toes, lymphadenopathy or the "uveo parotid syndrome" such as is observed in typical sarcoidosis. Sarcoidosis is a generalized disease affecting the reticulo endothelial structural system. Intestinal involvement by sarcoidosis without generalized implication of other organs would be difficult to accept.

## 10      Considerations Regarding Therapy in Regional Ileitis

---

IT IS AN OPEN QUESTION whether any conservative or expectant form of medical treatment is warrantable in regional ileitis. True, it has been shown that in this series twelve out of 222 cases progressed to what seemed to be spontaneous resolution with the disappearance of roentgen evidence of the disease. True too, that it has been demonstrated that in a percentage of cases the lesion remains in the terminal ileum showing over the course of years little tendency to extend in an oral direction. These cases develop cicatricial stenosis, the true "string sign" in roentgen study and some of them eventually progress to intestinal obstruction. They may remain stationary without developing skip lesions year after year. Occasionally abdominal wall fistulas have been demonstrated to have closed spontaneously. Thus there would seem to be some rationale for the development of a conservative form of medical therapy.

On the other hand in the usual case of regional ileitis when first seen for diagnosis advanced skip lesions higher up in the ileum can often be demonstrated proximal to the parent distal area of involvement. And many if not most of the cases show over the course of years not only proximal intestinal extension but if conservatively observed, demonstrate a continuity of clinical symptoms that bespeak a dynamic disease having an active life cycle. If not stayed by surgery the repetitive series of fever, diarrhea, anemia

case of endometriosis as a cause of intestinal obstruction has been described (McGuff et al), in which the segment 5 cm proximal to the ileo cecal valve was involved by endometrial transplants. The clinical picture resembled in general a case of regional ileitis with intestinal obstruction.

Enterogenous cysts about the ileo cecal angle are exceedingly rare, but may simulate roentgenographically the constriction of terminal ileitis (Rosenberg).

*Actinomyces*, intestinal, with abdominal wall fistulas may simulate ileitis. The finding of the characteristic ray fungus in the discharges from the fistula will save much confusion and often embarrassment. The possible confusion between ileitis and sarcoidosis has already been discussed (page 16). Sarcoidosis of Boeck involving the small intestine as a granulomatous lesion was described recently by Watson and his associates. The identity of ileitis and sarcoidosis was even suggested by Hadfield. Snapper has however, made the point that he has never encountered a case of sarcoidosis with a lesion of the intestine, nor have his many cases of ileitis ever shown sarcoidosis. Blackburn states that in ileitis there is no record of swelling of the fingers and toes, lymphadenopathy or the "uveo-parotid syndrome" such as is observed in typical sarcoidosis. Sarcoidosis is a generalized disease affecting the reticulo endothelial structural system. Intestinal involvement by sarcoidosis without generalized implication of other organs would be difficult to accept.

in this series have either been operated upon or are awaiting a decision for surgical intervention, because of the progression of symptoms and the appearance of complications.

Since the definitive method of handling recurrences after operation has not yet been determined, these cases, too while being observed, must have some plan of conservative therapy.

The indications for conservative medical therapy should apply to

1 Those cases who refuse surgical intervention and insist on palliation

2 Cases of long standing low grade intensity whose pathologic lesions show no tendency to progression

3 Postoperative recurrences whose future course of treatment has not yet been determined

For all other cases, constituting the largest group surgical intervention seems indicated as soon as the diagnosis has been established.

### Diet

A bland low fat protein rich diet should be instituted. Abundant proteins and the liberal use of carbohydrates are necessary to maintain nutrition against the wasting of the disease and against the hypoproteinemia that may ensue as a result of protracted fever and diarrhea. Coarse fruits and vegetables should be eliminated soft vegetables well cooked should be pureed or passed through a collander. Fruit juice may be allowed in limited quantity soft fruits such as bananas apple sauce soft pears, ripe peaches are allowable. The greatest attention is paid to meats and fishes cheeses protein milk and eggs as foods with high nitrogen content. The diet should be supplemented by the administration of protein or casein hydrolysate with high amino acid content 6 8 or 12 ounces a day taken in milk water fruit juices cocoa substitutes or in any manner that makes them possibly palatable to taste.



and fistulas indicate a pathologic process that, by itself, suggests no cessation of the underlying infection. It is in this latter type of case that delayed surgical intervention is so often disappointing and probably accounts for the high percentage of recurrences after operation.

If one could determine, by study of a case over a course of years, that a particular patient had a stationary type of lesion which did not show progression, one might be warranted in attempting some form of conservative therapy. By that same token, if the patient were one whose lesion showed progression, then medical therapy should be rapidly replaced by surgical intervention. But would the delay, protracted over a course of several months if not years, be warrantable? And are not the surgical failures to be charged up against procrastination when only too late one becomes conscious of complications, sequelae and progression of the lesion? Clark and Dixon have stated that any one who has seen the irremediable damage sustained by an affected segment through prolonged delay must realize that there is no known medical treatment that will dissolve a fibrotic stenosis, or will close a fistula that develops in the later stages of ileitis. Lahey and Sanderson make the categorical statement that there is no medical treatment worth consideration in regional ileitis.

The percentage of spontaneous cures in this present series was small, actually 5.4 per cent. The percentage of surgical cures is high, even when reckoned conservatively, and is grossly well over 80 per cent. Then why contemplate palliative treatment at all, why not operate as soon as the diagnosis has been established? Those twelve cases of spontaneous healing represent those who have refused operation and preferred to undergo expectant therapy. They are the small minority who have won out by their own persistence. But it is questionable whether the success in this minimal group can be utilized to formulate a plan of expectancy treatment that will largely apply to the massed group of cases. The remainder of the group of 210 cases

6 30 p.m. *Dinner*

Choice of soup

Choice of main dish Lean beef (boil broil or roast) chicken turkey or fish Portion of strained cooked vegetable Portion of potato, rice or spaghetti Bread or roll with butter or jelly Dessert Jello cornstarch pudding or apple sauce Milk cocoa or weak tea.

9 30 p.m.

Milk soft drinks may be taken in small quantities if not too cold and most of fizz is expelled (or see 10 30 and 3 30)

### *Diet*

#### *Foods Allowed in Diarrhea*

All food should have their roughage reduced to a smooth form by putting through a sieve so that no rough indigestible portions are left to irritate the bowel

*Milk* At least a pint a day Milk need not be boiled Cultured milks may be used acidophilus buttermilk, etc Mellin's food or malted milk may be added to milk

*Note* Most persons tolerate milk very well However, milk may cause diarrhea in those few persons who are 'allergic' to milk Those who find themselves "allergic" to milk (diarrhea) should omit it

*Eggs* Two or three daily Use in any form except raw or soft boiled

*Cheese* Any mild cheese may be used cottage pot or sour cream cheese gruyere swiss, etc Small portions of sour cream

*Cereals* All highly milled grains that are well cooked farina cream of wheat, wheatina, cornmeal white rice, ralston, etc

*Note* When using ready to serve cereals, check advertising and reject all cereals containing bran Safe ready to serve cereals are cerevin cornflakes grape-nut flakes pabulum puffed rice, puffed wheat Spaghetti noodles and macaroni in moderation

*Meats* All lean meats boiled baked or broiled ground lean steak roast beef steak liver of all kinds veal, lamb, smoked ham chicken squabs turkey

*Fish* All white fish boiled baked or broiled halibut cod, flounder, sea bass etc shell fish oysters and clams

In one case of high ileo jejunitis where operation was contraindicated, a gain of 50 pounds of weight and a return to fair health and strength followed a conscientious forced attempt at superalimentation with protein foods and amino acid commercial preparations

## CONSERVATIVE PALLIATIVE TREATMENT OF REGIONAL ILEITIS

### *General Directions*

Live in an atmosphere which is peaceful Try to control anxiety, worry or mental upset Plan every day so as to have a rest period and as much quiet as possible

Fresh air and sunshine are fine aids to a quick recovery

No competitive games or strenuous exercises Plan recreation which does not produce nervous tension or great fatigue, no tennis match, no card games for money, no horse back riding

Have a full and highly nutritious diet by eating three meals with light mid morning, mid afternoon and evening lunch That is, eat once every two and a half hours throughout the day and little or nothing after 10 P.M. The diet which is suggested for diarrheal diseases is as follows

### *Sample Menu*

#### 8 00 a.m. *Breakfast*

Orange or tomato juice  $\frac{1}{4}$  cup (dilute with water if liked better)  
Cereal with milk and sugar Bread (white) 2 slices (toasted) with butter Egg (1 or 2) boiled hard poached skinned or scrambled  
Milk cocoa or weak tea (1 cup)

#### 10 30 a.m.

Hard boiled egg and cracker and milk cocoa or weak tea and cracker

#### 1 00 p.m. *Lunch or Supper*

Choice of soup strained dried bean or split pea soup or cream soup (made with strained vegetables) or broth and any other strained soup

Choice of main dish Eggs or macaroni and cheese or portion of liver meat or fish

Serving of potatoes rice or noodles Bread (white) 2 slices with butter Dessert Strained cooked fruit plain cake or custard

#### 3 30 p.m.

Banana or  $\frac{1}{4}$  cup orange or tomato juice (or see 10 30 a.m.)

3 Avoid any gas forming food such as cabbage, corn, kohlrabi, broccoli, pickles and relishes of all kinds, also skins of baked apples and potatoes. Eat no beans, peas or lentils except in pureed form.

4 Avoid nuts, raisins, coconuts, figs, dates, etc, even if used in small amounts in puddings or cookies.

5 Avoid heavy fat foods: gravies, sausages, pastries, doughnuts, fried meats and fish.

6 All condiments, especially pepper, mustard, horse radish, vinegar, etc are best avoided. Therefore, do not use any highly seasoned and spiced foods, such as smoked meats and fish, frankfurters, most canned meats and fish, salt fish, olives, etc.

7 Avoid fish high in fat—salmon, shad, etc. and shell fish: shrimp, lobster and crab.

8 Avoid all grossly sweet foods: preserves, most candies, all rich cakes with icings and fillings.

9 Alcohol is an irritant and should be avoided.

10 Coffee should be omitted from diet, and tea served weak.

11 Care must be taken not to use certain foods which are laxative as large amounts of lemon, grapefruit or orange juice, raw vegetable juices, long iced drinks and many cups of too hot drinks as well as melons, grapes, plums, molasses, raw or soft boiled eggs.

### *Ambulation*

Most cases of regional ileitis being afebrile, may be allowed some physical activity. They are best treated as ambulant cases because the long course of the disease would make persistency in insisting upon bed rest a psychic and mental handicap. Most of the patients suffering from ileitis are not strong enough to work or attend higher educational schools or colleges but they are happier up and about and would probably gain little by prolonged bed rest. The acute cases and those with a febrile course should have bed rest during the active phase; thereafter early ambulation is advisable to preserve muscle tone.

*Bread* White bread (not too fresh), melba toast, zwieback, rolls, plain sweet breads like brioche, also all crackers made of white flour

*Butter* In moderation Thin sweet cream if well tolerated

*Fruits* With the exception of orange juice and ripe banana, all fruits must be cooked and sieved. Note cooked dried fruit and canned fruit should likewise be sieved. Use apples, pears, peaches, prunes, apricots, etc

*Vegetables* Use tomato juice. Sweet or white potatoes cooked any way but fried. Carrots, green peas, string beans, onions, spinach, beets, turnips, tomatoes, should be cooked and pressed through a sieve. In milder cases, cooked chopped spinach and beet tops, asparagus tips and fresh tender lettuce may be used. Note Canned infant foods are very satisfactory for use

*Desserts* Custards, Bavarian cream, blanc mange, bread, rice and tapioca puddings, fruit whip, gelatin jellies, plain, sponge and angel cake, plain coolies. Ices and ice cream are not desirable but may be used in moderation in mild cases

*Miscellaneous* Strained jams and preserves may be used in limited amounts. Grape, pear, apricot and pineapple juices may be used in small portions. Juices of all mild cooked fruits may be used with added dextrin maltose (not lactose which is laxative). Hard candies are to be encouraged

A certain amount of seasoning is allowable, that is, salt and pepper in moderation

### *Foods to Avoid During Diarrhea*

1 Avoid all raw fruits and vegetables except strained orange juice in small amounts and ripe bananas. All other fruits should be stewed

2 Avoid bran in (a) coarse breads (pumpernickel, whole wheat breads, muffins, health breads, etc.) (b) coarse cereal or (whole wheat, rice crisp, Swedish wafers, etc.) (c) coarse cereals (oatmeal, brown rice, barley, hominy, pott

some of the suppurative complications call for antibiotics that would tend to sterilize the bacterial content of the intestinal lumen. Some efficacy of the sulfonamides in controlling the symptoms and progression of the disease seems suggested in studies of the use of sulfasuxidine and sulfathaladine (Crohn) in ileitis though convincing proof is lacking and though these drugs in the largest percentage of cases fail to accomplish any striking therapeutic result. Sulfasuxidine (succinyl sulfathiazole) in doses of 0.2 grams per kilogram of weight is frequently employed the usual method is an initial dose of approximately 20 to 36 half gram tablets (10-18 grams) followed by dosage of five to nine tablets four times daily by oral administration. Sulfathaladine (phthalyl sulfathiazole) 0.1 grams per kilogram of weight usually 2, 3 or 4 half gram tablets four times daily are equally efficient.

Both of these drugs are bland and relatively free of untoward results. Agnuclocytosis has not been observed in this series nor has hepatitis been seen except in one instance. Many patients have taken one or both drugs for protracted periods in one instance 3000 tablets were continuously ingested over a course of six months without untoward results. In many instances the disappearance of external abdominal wall fistulous tracts has been observed apparently as the result of sulfa therapy by oral administration. This is in itself a large accomplishment even though as a rule the insoluble sulfonamides do not control the active symptoms or the progression of the disease in regional ileitis [61].

Penicillin has no apparent specific effect on ileitis, though it is useful in controlling the suppurative complications when walled off perforation does occur. As a preoperative and postoperative antibiotic form of therapy penicillin is in great favor with most surgeons. Streptomycin too when given intramuscularly 2 grams daily in divided six hour doses, and one gram orally coincidentally, may sterilize the intestinal tract and produce a sterile culture of the

(Femoral phlebitis and pulmonary embolism have not been noted in any cases of this series)

### *Various Medications*

In dehydration, particularly with diffuse draining fistulas, water and electrolyte replacement is best attained by the use of parenteral intravenous therapy. Saline solution, Ringer's solution, 5 per cent glucose in water or in saline and the protein hydrolysates or amino acid preparations are indicated in amounts of 1500 cc per day, or 100 to 150 grams of protein daily. Blood plasma and occasional transfusions of 500 cc are indicated for protein replacement and in instances of advanced anemia.

In chronic ileitis vitamin replacement is strongly indicated. Concentrates of vitamin B complex reinforced by thiamin 10 to 20 or more milligrams per day may be administered orally though the absorption in the face of diarrhea may be problematic. In all cases, intramuscular injection of crude liver extract and of vitamin B complex, 2 cc of each, should be given into the buttock every other day, larger doses, as in sprue, may even be indicated. It is rare that the other vitamins are indicated, there is no material deficit of vitamin A or carotene, nor of ascorbic acid nor of vitamin D in instances of ileitis since the diarrhea is rarely so profound as to create such vitamin deficiencies except after extensive and repeated surgical resections or short circuiting procedures. Synthetic vitamin K is indicated, however, in all instances of hemorrhage from the bowel wall even though the prothrombin time is well within the normal range. Five to ten milligrams of synthetic vitamin K daily intramuscularly, should as in ulcerative colitis, control the hemorrhagic manifestations.

The insoluble sulfonamides are indicated in regional ileitis, even though the therapeutic result has not been convincingly demonstrated. Whether the disease is bacterial in origin or nature has not been demonstrated. But the bacterial contamination of the intestinal ulcerations and

## 11 Surgical Treatment of Chronic Regional Ileitis

---

THE SURGICAL RELIEF of regional ileitis is indeed a complicated and rapidly changing picture. It should be interesting, chronologically to follow the trends of opinion over the last fifteen years and to note the various schools of thought and of procedure. For the subject is apparently still in a state of flux, as is natural where methods change, procedures differ, judgments do not agree and the follow up period is all too short.

In general, one may note historically certain salient or outstanding surgical procedures in the passing years of experience:

- 1 Resection in one stage was the initial procedure of choice.

- 2 The gradual evolution of a two stage procedure (short circuiting, followed subsequently by resection of the original lesion).

- 3 Adoption of a single stage operation, namely ileo transverse colostomy, with transection of the ileum, as a sufficient procedure to induce cure.

- 4 Introduction of the Mikulicz technique, for primary resection by the Lahey Clinic.

- 5 Persistence of the two stage operations favored by the Mayo group.

- 6 Preference of Bockus for primary resection.

- 7 Skepticism of some of the Boston surgeons for all types of procedures.



feces, and yet it is questionable whether streptomycin has any clear indications for its use in the chronic protracted form of ileitis. In the acute cases, and in the febrile forms, streptomycin may be pragmatically utilized though its effects have not yet been proven of clinical benefit. I have actually seen one case in which streptomycin produced a sterile culture of the feces without in any way controlling the activity of the symptoms. And yet, in a severe and febrile case its use experimentally could well be justified.

Radiation therapy has been rather extensively employed, it was originally invoked by Eggers for the control of cases of intestinal granulomas and has been recommended and used by many writers in the literature. My own experience with radiotherapy in ileitis has been very disappointing, though it has been limited to few cases. We have had opportunity to observe the use of extensive radiotherapy in 18 cases, the radiation therapy having been previously employed by clinicians in attempts to control the disease and its complications. Those cases that I have seen have all been failures, if there have been successes in the use of radiotherapy in ileitis they were not apparent in this series under study, nor when employed by me as an initial form of therapy.

In instances of intestinal obstruction the Miller Abbott tube is beneficially employed and by decompressing the intestinal lumen rapidly produces amelioration of the obstructive phenomena.

## 11 Surgical Treatment of Chronic Regional Ileitis

---

THE SURGICAL RELIEF of regional ileitis is indeed a complicated and rapidly changing picture. It should be interesting, chronologically to follow the trends of opinion over the last fifteen years and to note the various schools of thought and of procedure. For the subject is apparently still in a state of flux, it is natural where methods change, procedures differ, judgments do not agree and the follow up period is all too short.

In general one may note historically certain salient or outstanding surgical procedures in the passing years of experience:

- 1 Resection in one stage was the initial procedure of choice

- 2 The gradual evolution of a two stage procedure (short circuiting followed subsequently by resection of the original lesion)

- 3 Adoption of a single stage operation namely, ileo transverse colostomy with transection of the ileum, as a sufficient procedure to induce cure

- 4 Introduction of the Mikulicz technic for primary resection by the Lahey Clinic

- 5 Persistence of the two stage operations favored by the Mayo group

- 6 Preference of Boeckus for primary resection

- 7 Skepticism of some of the Boston surgeons for all types of procedures

8 Growing realization of the increasing incidence of recurrences following all types of surgical interference

1 The initial surgical procedure at Mount Sinai Hospital practiced by A. A. Berg was primary resection, as was carried out on the first 14 cases of terminal ileitis, this series was carried up to 32 instances of resection by Berg in 1936. Correspondingly, most of the early cases in the literature concerned themselves only with primary resections of the terminal ileum. In 1936, Koster, reviewing 100 operated cases from the literature, found resection almost universally carried out as the procedure of choice, adding 17 cases of his own, most of which had been resected. In 1939, Ravdin and Johnston, reviewing the surgical literature to date, found 290 instances of primary resection against only 88 instances of short circuiting procedures. In 16 cases, enterostomy alone had been performed.

2 Already in 1933 one notes the suggestion independently by Clute and by Homans and Hass of ileo transverse colostomy, that is, a short circuiting of the lesion as a sufficient procedure, rather than resection. Unfortunately, neither contribution mentions whether transection of the ileum proximal to the lesion had been performed. Even before this date, Estes and Holm had warned against such a procedure: they demonstrated in 2 human cases and by animal experimentation that huge dilatation of the side tracked and excluded loop of ileum could and did occur. This point of view, however, was subsequently contradicted by the studies of Ginzburg, Colp and Sussman. By surgical experience, substantiated by careful roentgenologic studies, these latter authors proved that a short circuiting procedure with transection of the ileum was a safe and satisfactory procedure that involved no complications from the excluded loop of ileum.

By 1934, some of the members of the Mayo group (Clark and Dixon) were already experimenting with short circuiting procedures, having utilized such a plan in 5 cases out of 13 operables. Again, transection of the ileum was not

mentioned. In 1939, Clark and Dixon suggested for the early stages of the disease a simple short circuiting procedure with transection of the ileum which carries a low mortality and provides an opportunity for the diseased segment of the bowel to improve by being placed at rest. Subsequent resection of the diseased area was utilized if activity of the original site of disease continued. If the disease process was far advanced primary resection, a radical extirpation of the diseased segment, was advocated.

Mixter (1935) utilized a one stage or a multiple stage procedure in 11 cases, but unfortunately this small series carried a mortality of 4 cases or 36 per cent. In 1936, Koster for the first time warns that if a short circuiting procedure is to be performed transection or "occlusion" of the ileum must be carried out. He still preferred primary resection since he suspected the undesirability of leaving in situ the original focus of infection.

The importance of transecting the ileum above the diseased segment was realized in the following case in this present series.

*Case II 8 (Serial No 146)* Male age 21 years. For eighteen months this young man had suffered with diarrhea, fever and abdominal cramps. A mass in the right lower abdominal quadrant had formed. He had been operated upon at that time and a short circuiting procedure had been carried out without transection of the ileum. Diarrhea persisted; there was a weight loss of 50 pounds; edema of the lower extremities was present due to hypoproteinemia. Clubbing of the fingers and severe secondary anemia (hemoglobin 67 percent) were evident. At operation a transection of the ileum above the persistent and residual ileitis was the only procedure performed. This was followed by a regain of the 50 pounds of weight lost and a subsidence of the peripheral edema of the lower extremities. Unfortunately two years later a recurrent ileitis proximal to the old anastomosis was capable of demonstration. A two stage resection of the recurrent ileitis resulted in cure.

3. The transition of the Mount Sinai group from primary one stage resection to two stage resection and finally to a short circuiting procedure only is interesting to observe. In 1936, though primary resection seemed to be in favor it

was noted that both Lewisohn and Colp had 6 cases of short circuiting procedures with good results. Lewisohn later reported (1938) 9 cases of short circuiting or two stage procedures with only one death. He again strongly emphasized that a lateral anastomosis between upper healthy ileum and the colon, without transection of the ileum above the diseased segment is a useless operation since the affected part of the bowel is not put at rest.

In the one fatal case, a primary ileo transverse colostomy with transection of the ileum had been primarily performed. After a sufficient interval of what was then regarded as the proper sequence, the second operation of resection of the primary residual focus of disease was carried out. The resected specimen was now fully healed and cicatrized. Unfortunately, the child died postoperatively of pulmonary embolism due to an old rheumatic endocarditis. The tragic death and the evidence that the secondary intestinal resection had been superfluous, led to a revision of the views regarding the needs of the secondary resection. Bowen and Day report the postmortem findings in a case of regional ileitis nine years after symptomatic recovery following ileo colostomy. The ileum, previously affected, was at the autopsy shown to be fibrosed and sclerotic, the lumen almost obliterated by stenosis and by multiple adhesions. Complete healing of the ileum had been affected by the short circuiting procedure.

In 1939, Ginzburg, Colp and Sussman published 14 cases of short circuiting operations for ileitis, transection of the ileum being regularly carried out. Again they were convinced of the lack of necessity for any subsequent resection, the initial lesion drying up and undergoing atrophy after the short circuiting, fistulae disappearing, abdominal masses resolving. The x-ray studies of Sussman convincingly demonstrated no trapping in the excluded loop and good intestinal function through the new stoma.

In 1941, Colp and Ginzburg published a report of ileocolostomy with exclusion as practiced in 22 cases of terminal

ileitis with no deaths, 3 of the series showed recurrent symptoms. Four of the cases were re-explored because of persistent diarrhea, in all 4, healing of the original site of infection in the terminal ileum was clearly demonstrable and no further resection was attempted. Six cases had abdominal wall fecal fistulas before operation, after the short circuiting procedure, five of the fistulas healed and ceased to discharge content. In 1942 Ginzburg and Garlock published 54 cases of ileo colostomy with exclusion with no operative mortality, with two recurrences of the disease in the proximal ileum and with 5 cases with persistent clinical symptoms. Contrasted with this were 23 cases of primary resections with four deaths and three recurrences.

In 1945, the largest group of cases from the Mount Sinai Hospital was published (Garlock and Crohn), covering a twelve year experience with the surgical treatment of chronic regional ileitis. The series comprised 164 cases of enteritis, terminal ileitis and combined ileo colitis, most of which had been operated upon by various surgeons in that institution (137 cases) the remaining 27 cases had been operated elsewhere but reported for observation because of persisting symptoms. The inclusion of these latter cases, which represent failures of previous procedures depresses the statistical value of the total end results of the series but had been included nevertheless in the interests of scientific accuracy. The results may be seen in table 10.

TABLE 10—Surgical Treatment of Chronic Regional Ileitis

	Number	Deaths	Recurrences
Ileo colostomy with exclusion	6	0	9 (13.8%)
One stage resection	55	9 (16.3%)	9 (16.5%)
Two stage resection	5	3 (1.7%)	8 (36.3%)
Combined ileo colitis	19	(10.5%)	3 (1.6%)

By a process of comparison the advantages of the ileo colostomy with exclusion are obvious. The mortality rate was nil, the recurrence rate (as of that date) was moderate but yet considerable. The group of cases comprising primary resection was less satisfactory. A considerable mor-

was noted that both Lewisohn and Colp had 6 cases of short circuiting procedures with good results. Lewisohn later reported (1938) 9 cases of short circuiting or two stage procedures with only one death. He again strongly emphasized that a lateral anastomosis between upper healthy ileum and the colon, without transection of the ileum above the diseased segment is a useless operation since the affected part of the bowel is not put at rest.

In the one fatal case, a primary ileo transverse colostomy with transection of the ileum had been primarily performed. After a sufficient interval of what was then regarded as the proper sequence the second operation of resection of the primary residual focus of disease was carried out. The resected specimen was now fully healed and cicatrized. Unfortunately the child died postoperatively of pulmonary embolism due to an old rheumatic endocarditis. The tragic death and the evidence that the secondary intestinal resection had been superfluous, led to a revision of the views regarding the needs of the secondary resection. Bowen and Day report the postmortem findings in a case of regional ileitis nine years after symptomatic recovery following ileo colostomy. The ileum, previously affected, was at the autopsy shown to be fibrosed and sclerotic, the lumen almost obliterated by stenosis and by multiple adhesions. Complete healing of the ileum had been effected by the short circuiting procedure.

In 1939 Ginzburg, Colp and Sussman published 14 cases of short circuiting operations for ileitis, transection of the ileum being regularly carried out. Again they were convinced of the lack of necessity for any subsequent resection, the initial lesion drying up and undergoing atrophy after the short circuiting fistulae disappearing, abdominal masses resolving. The x-ray studies of Sussman convincingly demonstrated no trapping in the excluded loop and good intestinal function through the new stoma.

In 1941, Colp and Ginzburg published a report of ileo colostomy with exclusion as practiced in 22 cases of terminal

failed to result in amelioration of symptoms or relief in 36.3 per cent of this smaller series. These are the distressing and depressing cases that create skepticism as to the success of any or all surgical procedures as adapted to the cure of regional ileitis.

4 The Lahey Clinic favors a Mikulicz type of resection for ileitis, the diseased ileum and the ascending colon are brought out on the surface of the abdomen, exteriorized and resected extraperitoneally. A secondary closure of the enterostomies is later performed. Marshall in 1940 and again later in 1943 reported a ten year survey of 55 patients surgically treated. The operative mortality was 5.5 per cent, in a follow up that covered 42 patients of the series there were five recurrences or 9.6 per cent. In the presence of abscess or infection a short circuiting procedure is preferred as a first stage with a later secondary resection. Note again that even here the secondary resection is regarded as essential.

Of this series, 35 cases underwent the Mikulicz procedure with but one death, a highly commendable record.

5 The group at the Mayo Clinic seem to prefer a two stage operation. In 1937 Pemberton and Brown reported 39 cases of ileitis treated by two stage short circuiting and resection with three deaths. They allowed three weeks to six months between the first and the second procedure. Transection of the ileum as part of the first operation was always mandatory. In 1938, Dixon reported 40 operated cases the procedures were evenly divided between primary resection two stage operations and short circuitings alone. Again the gross mortality was low, 10 per cent.

In 1939 Clark and Dixon questioned the utility of the short circuiting operation per se they considered 50 per cent at best the estimate of cure after such a procedure, and emphasized the high percentage of postoperative recurrences following short circuiting procedures as the sole type of operation.



tality was apparent, nor did there appear to be any compensation for the greater risk since the recurrence rate of resection was even greater than that of the exclusion operation. Here the longer follow up period of the earlier resected cases may well be a factor since recurrences were noted once eight years and once twelve years after the original operation, the intervening years having been a period of apparent perfect health. Bockus had already noted a recurrence of postoperative ileitis nine years after apparent good health.

It cannot be too strenuously emphasized that transection of the ileum proximal to the diseased area is a *sine qua non* to the success of any operation (Lewisohn). In this present group of 222 cases of regional ileitis, 8 cases had been operated elsewhere, without transection of the ileum. All of them were in distress because of persistent symptoms. In one case the mere transection of the ileum at a subsequent operation resulted in control of the diarrhea and a gain of 50 pounds of weight. In many of the others, radical excision of the lesion was necessitated by the omission of the former surgeon properly to exclude the primary focus of disease. The remarkable case published by Forbes and Duncan in which a true ileo vesical fistula was cured only after transection of the ileum had been secondarily accomplished is most instructive.

To return to the analysis of end results of 164 surgically treated cases as published by Garlock and Crohn. The two stage operation consisting of a primary ileo colostomy with transection of the ileum, followed some time later by a resection of the primary focus of disease, constituted the most distressing group. Naturally so, because these cases all represent failures of short circuiting procedures. The second stage of the operation was forced by the gravity of the persisting symptoms at the original site of disease. Actually, they represent failures of the first stage short circuiting procedure. The forced secondary operation was fraught with a moderate mortality (12.0 per cent) but

of resection was employed in two of his cases, probably the first time this modified form of resection had been mentioned in the literature. Cutler (1939) even went a step further and suggested that surgery was not a cure for ileitis, recurrences following radical surgery were frequent, patients may recover spontaneously. He suggested the retention of the case in the hands of the family physician as possibly a good solution.

In 1942, Warren and Miller reported 26 resections for ileitis with only one death, however, in the follow up, only 6 patients were free of disease, most of them incapacitated.

These reports published from Boston represent individual opinions. None of the series of cases was large the opinions expressed seem to be depressing though contrary to the more optimistic conclusions of those with far greater numbers of cases and with equally critical scientific analyses of their results. The recent critical editorial (Starr), "Is There an Adequate Therapy for Regional Enteritis?" concedes that surgery offers the best opportunity for palliation and the control of regional enteritis, but emphasizes that the end results leave much to be desired.

The most disappointing effects are seen when attempts have been made to perform resections of the ileum in continuity with end to end anastomosis of the small bowel. Procedures such as this one are insufficient unphysiologic and lead to skepticism (3 such cases in this present group). This same criticism does not apply to high jejunal involvement where local resection is eminently successful.

8 Recurrences. There is the growing realization of the increasing incidence of recurrences following all types of surgical interference increasing incidence because the longer the follow up the greater the possibility for late recurrences, since by now it is recognized that recurrences can and do occur eight, nine or even twelve years after the operative procedure.

Recurrences are mentioned in the literature almost as soon as operative procedures had been initiated (Hornum and

In 1941, Mayo and Judd reported a full 100 cases, 64 of which underwent the two stage operation, with a 3 per cent mortality, the average length of time between the two stages was 14.4 months. With multiple procedures in the complicated cases, Brown and Donald had a higher mortality of 9 per cent for a larger series (178 cases). They suggest surgical removal of the diseased segment of bowel "a two stage procedure is better than a one stage operation."

Thus, the Mayo group still prefer the two stage procedure, eschewing the single stage resection and not accepting a simple short circuiting operation as sufficient.

6 Bockus clings to his preference for single stage resections. He draws support from the report of Rhoads who collected from the literature 125 cases in 51 of which a side tracking procedure alone had been carried out with a high percentage of recurrence (47 per cent). Bockus feels that considerable progress in the course of the disease may take place between the two stages of a double procedure, a point which is flatly contradicted by the experience of other writers. He believed in 1940 that a one stage operation could be safely performed. Again in 1944 he feels that the chances of cure are 2 to 3 times greater with primary resection than with short circuiting procedures. His medical cases were submitted to resection with a mortality of 7.7 per cent and only one late recurrence out of 18 cases. He retains throughout his confidence in primary resection, his series of cases is, however, small.

7 Pessimism as regards results in the operative treatment of ileitis seems to spring from some, but not all, of the Boston surgeons. Already in 1935 Mixer in a questionnaire to various surgeons reported a 20 per cent recurrence rate after operative procedures. In a small series of 11 difficult cases with complicated and multiple procedures, his mortality rate was 36 per cent. Mixer restricted his surgical procedures to, and advocated the use of resection in one or two stages. The Mikulicz technic

of resection was employed in two of his cases, probably the first time this modified form of resection had been mentioned in the literature. Cutler (1939) even went a step further and suggested that surgery was not a cure for ileitis, recurrences following radical surgery were frequent, patients may recover spontaneously. He suggested the retention of the case in the hands of the family physician as possibly a good solution.

In 1942, Warren and Miller reported 26 resections for ileitis with only one death, however, in the follow up, only 6 patients were free of disease, most of them incapacitated.

These reports published from Boston represent individual opinions. None of the series of cases was large the opinions expressed seem to be depressing though contrary to the more optimistic conclusions of those with far greater numbers of cases and with equally critical scientific analyses of their results. The recent critical editorial (Starck)

'Is There an Adequate Therapy for Regional Enteritis?' concedes that surgery offers the best opportunity for palliation and the control of regional enteritis but emphasizes that the end results leave much to be desired.

The most disappointing effects are seen when attempts have been made to perform resections of the ileum in continuity with end to end anastomosis of the small bowel. Procedures such as this one are insufficient unphysiologic and lead to skepticism (3 such cases in this present group). This same criticism does not apply to high jejunal involvement where local resection is eminently successful.

8 Recurrences. There is the growing realization of the increasing incidence of recurrences following all types of surgical interference increasing incidence because the longer the follow up the greater the possibility for late recurrences since by now it is recognized that recurrences can and do occur eight nine or even twelve years after the operative procedure.

Recurrences are mentioned in the literature almost as soon as operative procedures had been initiated (Hornum and

Hass, Shcater and Jackson, Oppenheimer, Felger and Schenk, Koster et al [15 per cent of recurrences]) These recurrences occur almost uniformly in the ileum just proximal to the new anastomosis, 6 to 8 inches in extent, the pathology of the recurrence grossly and histologically resembling the initial lesion. Skip areas above the local recurrence are unusual, though exceptionally these recurrences may extend continuously or in an interrupted fashion throughout the remaining ileum and the jejunum. Internal fistulous tracts may originate at the anastomosis, more rarely from the involved segment of recurrent ileitis.

It is difficult to explain the rationale and method of propagation of such recurrences, particularly when the operation had been performed by a competent surgeon, one of experience in recognizing the highest possible skip lesion. In the best of hands, where the transection isolating the ileal lesion has been placed high and far distant above any suspiciously involved areas, approximately 15 to 20 per cent of recurrences are conceded.

Two possible methods of propagation of the recurrences suggest themselves.

(1) The surgeon is unable to recognize the uppermost skip lesion and makes his anastomosis too low in the ileum. That this mistake is due to ignorance or inadvertence or inexperience is gainsaid by the fact that recurrences occur at the hands of the most careful, meticulous and experienced surgeons. Obviously, if it be a mistake in judgment, it occurs because the mucosal or submucosal infiltration is so minimal as to preclude recognition by palpation or inspection of the wall of the small gut which is utilized for the anastomosis. It seems likely that such skip areas may indeed escape the vigilance of even the most adept surgeons. If this be the explanation then the loop of ileum chosen for the anastomosis must be high and far away from any suspected pathologic involvement. Such a high choice invites the danger of nutritional disturbances by sacrificing too much mucosa necessary for food absorption and by inviting the possibility of diarrhea.

(b) The recurrences are due to the persistence of a nidus of residual infection, outside the intestinal wall, most likely in the retained and inflamed mesenteric lymph nodes. If such were the case, then recurrences should occur with greater frequency after short circuiting operations than after radical resections, a fact which is contradicted by the table of figures. In the radical excisions of the lesion, it is usual to remove the succulent enlarged lymph nodes with the mesentery and with the original lesion, a deep angular resection of the mesentery being performed by most experienced surgeons. However, even with such a radical excision the recurrence rate (19.5 per cent) is higher than with short circuiting procedures (13.8 per cent) (Garlock and Crohn). If the lymph nodes actually constitute the source of residual infection and reinfection, it is necessary to postulate the spread against the normal lymphatic flow which is from the intestine to the lymph nodes rather than the reverse. This is contrary to physiologic knowledge though retrograde reinfection and retrograde metastases do occur as in carcinoma. Further it has been shown on secondary explorations after short circuiting procedures that when re-examined the originally inflamed lymph nodes are now diminished in size and show retrogression of the infection.

In this present series of 222 cases of ileitis the types of operation which had been carried out are given in table 11.

TABLE 11—Types of Operations for Ileitis

	(Per cent Series)
Palmay resection	36 cases
Ileo transverse colostomy with transection of ileum	5 cases
Ileo transverse colostomy without transection of ileum	1 case
Mikulicz procedure	1 case
Ileo sigmoidostomy with transection of ileum	4 cases
Resection of ileum in continuity	1 case
Ileo transverse colostomy with transection followed by second stage resection of lesion	10 cases
Total	108 cases

The rate of recurrence for this group will not be quoted because so many of these cases were operated elsewhere.

and came under observation because of secondary aggravation of symptoms. To a large extent they include a "trouble" group, specially referred because of postoperative distress symptoms. To attempt to draw conclusions as to the incidence of or percentage of recurrence from such a group of unfortunates would only enhance pessimism regarding surgery in ileitis and would not be a true over all critical analysis of the problem. A better idea of mortality and recurrence rate is obtained by referring back to the figures (table 12) quoted by Garlock and Crohn in their review of 137 cases personally chosen and supervised and operated at the Mount Sinai Hospital, New York City.

TABLE 12—Operation and Recurrence in Ileitis (Garlock and Crohn)

	Cases	Deaths	Recurrences
Ileo colostomy with transection	5	0	6 (105%)
One stage resection	4	6 (133%)	6 (154%)
Two stage resection	16	2 (12.5%)	4 (25%)

### DIAGNOSIS AND TREATMENT OF RECURRENCES

After operation, the patient usually regains strength and weight and appetite. Hemoglobin values improve, diarrhea is relieved, fever ceases. True, the diarrhea after short circuiting procedures may persist to a distressing degree for a few weeks, even months, but almost invariably when weight and strength return the number of defecations is diminished to a normal of two to three daily movements without pain or urgency.

A recurrence of the disease is marked by the return of fever, by loss of weight, increasing anemia and increasing diarrhea. The radiographic demonstration of irregularity of the mucosal pattern of the "new" terminal ileum, narrowing of the lumen, rigidity and distortion in the 6 to 8 inches of small bowel just proximal to the new anastomosis confirm the existence of a recurrence (fig 41). These recurrences of disease are usually localized in the area just proximal to the anastomosis. Rarely do they extend upward by skip lesions to involve the upper small intestine, and very rarely do they involve the colon, sigmoid or rectum. The recurrence is best demonstrated by barium

enema, the possible involvement of a higher segment of small bowel must be established or excluded by the barium meal study of higher ileum and jejunum. Many of these recurrences are exceedingly mild, fever is absent, loss of weight minimal, diarrhea and anemia being the only presenting symptoms. Even the diarrhea is undependable as a sign of recurrence since postoperative diarrhea even in successfully operated cases is not unusual. One depends




FIG. 41.—Recurrent Ileitis Proximal to a Distal Ile Transverse Colostomy. Stricture new terminal ileum proximal dilation of small bowel. (on creative medical therapy patient under observation eight years) (all general condition)

therefore to a large extent upon the radio-graphic interpretation of the loop proximal to the anastomosis. Such interpretation is not infallible, just as the original radio-graphic diagnosis of a lesion in the terminal ileum is not always accurate. Errors of omission and of commission are likely to occur (figs 42 and 43).

Recurrences are likely to be noted soon after operation in fact most of them occur within the first year of convalescence as noted in this present series (table 13).






Fig 42—Extensive Recurrent Ileitis Proximal to Ileo Transverse Colostomy Note the involvement of several loops of the “terminal ileum”

TABLE 13—*Recurrences in Present Series after Operation*

Immediate recurrences	6 cases
Within 6 months	4 cases
Within 1 year	4 cases
Within 2 years	1 case
Within 3 years	3 cases
Within 4 years	2 cases
Within 5 years	3 cases
Within 6 years	— cases
Within 7 years	1 case
Within 10 years	1 case
Within 16 years	1 ca

Total 98 cases

It is not to be implied from this table that recurrences are limited in time, postoperatively. In fact there is probably no time limit of follow up which precludes the possi-



Fig. 43—Recurrent Ileitis Proximal to Ileocolostomy. Recurrence six weeks after operation.

bility of a recurrence. Bockus refers in detail to a case in which, presumably nine years previously, a resection of the terminal 10 inches of the ileum had been performed for a lesion undoubtedly regional ileitis in nature. The patient was well for nine years, suffered recurrent symptoms and was re-explored surgically. A recurrent ileitis involving the area of ileum proximal to the ileo transverse colon anastomosis was demonstrated. My own experience has furnished two instances of very late recurrences.

*Case 1* 1 S (Serial No 170) Male age 39 years. Nine years previously he underwent a resection of the terminal ileum and ascending colon.

for a regional ileitis involving the last eight inches of the small bowel. The patient remained well and free of symptoms for approximately nine years or until six months ago when he developed a spontaneous fecal fistula breaking through the anterior abdominal wall at the site of the scar of the previous surgical incision. Exploratory laparotomy now demonstrated a recurrent ileitis involving the proximal ten inches of the 'new' terminal ileum. A two stage ileo sigmoidostomy followed by a resection of the involved recurrent segment of ileum and its connected transverse colon resulted in restitution of health.

*Case 2, V R A* (Serial No 83) Male age 34 years. In a second case one which constituted one of the earliest cases of ileitis seen and diagnosed the terminal ileum had been resected for regional ileitis in 1930. The uneventful recovery was followed by twelve years of perfect health and efficiency. Abrupt recurrent symptoms resembling intestinal obstruction required an emergency operation. At the exploration an obstruction at the site of the anastomosis and due to a severe acute recurrent ileitis was demonstrated. An attempt to resect the lesion in the face of the obstruction resulted in a fatal termination.

Recurrences with gross intestinal hemorrhage are rare, with abdominal wall fistulation somewhat more frequent. Recurrences in the colon, in the rectum and in the upper small intestine have been noted, fortunately infrequently, but represent a sad picture since little or a further surgical nature can be attempted. Sometimes permanent ileostomy above the recurrent lesion is the only means of saving life.

The very rare involvement of the colon in recurrent ileitis is a very interesting negative phenomenon. For though there is no equivalent of an ileo cecal valve of Bauhin at the new anastomosis and though the recurrent ileitis empties unguardedly into the exposed colon, the natural immunity of the colonic mucosa to the ileal infection seems to preserve it from involvement by extension. This was impressively demonstrated in one particular case in ileitis had been resected 18 years previously, the nature of the lesion being little suspected. Recurrence of symptoms occurred, at the subsequent operation eighteen years later the recurrent lesion and attached ascending colon were resected en masse the colon was free of disease though probably for most of the long period the recurrent ileitis emptied its content into the large bowel at the anastomosis.

## THE TREATMENT OF RECURRENT ILEITIS

Fortunately most of the recurrences are mild. Many of them can be controlled by conservative medical treatment. Such therapy consists of drugs that control the frequency of the diarrhea, particularly opiates, combating the anemia by intramuscular injection of crude liver extract and vitamin B complex, the administration of amino acid preparations or protein hydrolysates by mouth in large amounts, 6-8 ounces per day, rest, warm climate and sunshine. The insoluble sulfa derivatives, sulfasuxidine and sulfathalidine are usually administered though their value is often questionable. Occasionally a brilliant therapeutic result is achieved. An increasing number of cases of recurrent ileitis seem to do well and retain only mild clinical symptoms under conservative medical therapy. So much so that surgical intervention should be reserved only for those cases which refuse to improve under general therapeutic measures or which develop complications such as fever and abdominal wall or perirectal fistulas. A remarkable instance of spontaneous cure of recurrent ileitis and jejunitis is seen in the case reported by Pessagno. A boy, aged 13, underwent an obstructive resection of terminal ileum and cecum for regional ileitis. A recurrence of the symptoms plus external fistulas forced a secondary resection. A persistent or recurrent fistula now complicated by obstruction necessitated a third operation. At the second operation an all extensive involvement of the terminal ileum and the jejunum were noted. In spite of the extensive evidences of recurrences throughout the small bowel and the almost hopeless prognosis, the patient when last seen was in excellent condition and making a rapid recovery, appetite good, one or two stools a day, wounds entirely healed. Unfortunately the period of follow up was limited to less than a year.

A similar spontaneous regression of a recurrent ileitis may be observed in an interesting case in this present series.

*Case D G* (Serial No 53) Male age 37 years Nine months ago operated for a supposed acute appendicitis. At exploratory operation an acute ileitis was observed. Six months ago reoperated because of recurrent symptoms ileo transverse colostomy performed for terminal ileitis. At the time of operation two ileo ileal internal fistulas were demonstrated and another fistula from ileum to sigmoid was demonstrated. Six months later radiographic studies showed an extensive recurrence in the loop of ileum proximal to the new anastomosis. One year later the patient reported being in good health and free of symptoms. X ray examination at this time showed spontaneous regression of the recurrent ileitis and no evidence of the ileo sigmoid fistula. Two years later the follow up of this interesting case reported perfect health except for slight diarrhea.

*Surgical intervention in recurrent ileitis becomes necessary when fistulas appear, when the recurrence shows signs of extension, when the general nutrition is compromised and the anemia becomes progressive. Continued fever is further an indication for surgery.*

The surgical procedures for the cure of recurrent ileitis consist of a further short circuiting procedure, usually an ileo transverse colostomy distal to the previous anastomosis again with transection of the intervening ileum so as to exclude the newly involved segment. Some surgeons also transect the transverse colon distal to the first anastomosis, bringing the proximal cut end out on the abdominal wall as a temporary or mucous colostomy. A second stage procedure is now required to excise the recurrent ileitis and the attached transverse colon. Other surgeons, following the principle of short circuiting, merely make the new anastomosis, transect the ileum so as to exclude the involved segment and make no attempt further to resect the recurrent lesion. It is too soon to say which is the preferable surgical approach. It is certainly true that one operation is always better than a two stage procedure, providing it suffices to heal the recurrence. The mortality of such operation for recurrent ileitis is exceedingly small, and the improvement in well being weight and strength is very satisfactory. Unfortunately, too many instances can be cited of extensive and rapidly extending recurrences that defy surgical inter-

vention. Two such cases in this series will demonstrate the almost malignant nature of the rapid and universal involvement of the small bowel by recurrent ileitis.

*Case I M* (Serial No 229) Male, 17 years of age. Six months of abdominal pain and diarrhea. Regional ileitis involving the terminal 18-24 inches of the terminal ileum was demonstrated radiographically, and at the exploratory operation ileo transverse colostomy with excision of the ileum. Recurrent symptoms occurred exactly one year later with rapidly downhill course. At exploratory laparotomy thirteen months after the original short circuiting procedure, a diffuse ileo jejunitis extending up to the duodenum was demonstrated. No further surgical procedure was attempted.

*Case A M* (Serial No 110) Female single 29 years of age. Eight years of alternating diarrhea and constipation loss of 50 pounds of weight. Eleven months ago resection of terminal 2 feet of ileum (and ascending colon) for terminal or regional ileitis. Six months later because of rapid recurrence of symptoms the patient again underwent surgical exploration. The entire ileum proximal to the anastomosis and the jejunum were the seat of extensive recurrent inflammatory disease. Persistent abdominal wall fecal fistulas resulted after the last and futile exploration. The prognosis is absolutely hopeless.

Secondary recurrences do occur, fortunately infrequently. These are usually met when the indications for surgery become mandatory by a third anastomosis, usually an ileo sigmoidostomy, again with transection of the ileum above the secondary recurrence. These cases, too, though infrequent, occasionally do well, the diarrhea following such repeated surgery is not excessive.

The most recent suggestion regarding treatment for extensive diffuse ileitis or ileo jejunitis comes from a Western group. Vagotomy has been experimentally tried for the relief of symptoms and possibly the cure of severe ulcerative colitis. The idea has been extended to cover also the possible very serious forms of ileo jejunitis. Considering that a vagotomy is followed by atony when applied to the stomach it may well be that the putting at rest of the small intestine by the same surgical procedure can be similarly accomplished. Certainly the idea is worth consideration.\*

*Case D G* (Serial No 53) Male age 37 years Nine months ago operated for a supposed acute appendicitis At exploratory operation an acute ileitis was observed Six months ago reoperated because of recurrent symptoms ileo transverse colostomy performed for terminal ileitis At the time of operation two ileo ileal internal fistulas were demonstrated and another fistula from ileum to sigmoid was demonstrated Six months later radiographic studies showed an extensive recurrence in the loop of ileum proximal to the new anastomosis One year later, the patient reported being in good health and free of symptoms X ray examination at this time showed spontaneous regression of the recurrent ileitis and no evidence of the ileo sigmoid fistula Two years later the follow up of this interesting case reported perfect health except for slight diarrhea

Surgical intervention in recurrent ileitis becomes necessary when fistulas appear, when the recurrence shows signs of extension, when the general nutrition is compromised and the anemia becomes progressive Continued fever is further an indication for surgery

The surgical procedures for the cure of recurrent ileitis consist of a further short circuiting procedure, usually an ileo transverse colostomy distal to the previous anastomosis again with transection of the intervening ileum so as to exclude the newly involved segment Some surgeons also transect the transverse colon distal to the first anastomosis, bringing the proximal cut end out on the abdominal wall as a temporary or mucous colostomy A second stage procedure is now required to excise the recurrent ileitis and the attached transverse colon Other surgeons, following the principle of short circuiting, merely make the new anastomosis, transect the ileum so as to exclude the involved segment and make no attempt further to resect the recurrent lesion It is too soon to say which is the preferable surgical approach It is certainly true that one operation is always better than a two stage procedure, providing it suffices to heal the recurrence The mortality of such operation for recurrent ileitis is exceedingly small, and the improvement in well being weight and strength is very satisfactory Unfortunately, too many instances can be cited of extensive and rapidly extending recurrences that defy surgical inter

vention. Two such cases in this series will demonstrate the almost malignant nature of the rapid and universal involvement of the small bowel by recurrent ileitis.

*Case I M* (Serial No 229) Male 17 years of age. Six months of abdominal pain and diarrhea. Regional ileitis involving the terminal 18-24 inches of the terminal ileum was demonstrated radiographically and at the exploratory operation ileo transverse colostomy with exclusion of the ileum. Recurrent symptoms occurred exactly one year later with rapidly downhill course. At exploratory laparotomy, thirteen months after the original short circuiting procedure a diffuse ileo jejunitis extending up to the duodenum was demonstrated. No further surgical procedure was attempted.

*Case A M* (Serial No 110) Female single, 29 years of age. Eight years of alternating diarrhea and constipation loss of 50 pounds of weight. Eleven months ago resection of terminal 2 feet of ileum (and ascending colon) for terminal or regional ileitis. Six months later because of rapid recurrence of symptoms the patient again underwent surgical exploration. The entire ileum proximal to the anastomosis and the jejunum were the seat of extensive recurrent inflammatory disease. Persistent abdominal wall fecal fistulas resulted after the last and futile exploration. The prognosis is absolutely hopeless.

Secondary recurrences do occur, fortunately infrequently. These are usually met, when the indications for surgery become mandatory by a third anastomosis, usually an ileo sigmoidostomy, again with transection of the ileum above the secondary recurrence. These cases, too, though infrequent, occasionally do well, the diarrhea following such repeated surgery is not excessive.

The most recent suggestion regarding treatment for extensive diffuse ileitis or ileo jejunitis comes from a Western group. Vagotomy has been experimentally tried for the relief of symptoms and possibly the cure of severe ulcerative colitis. The idea has been extended to cover also the possible very serious forms of ileo jejunitis. Considering that a vagotomy is followed by atony when applied to the stomach it may well be that the putting at rest of the small intestine by the same surgical procedure can be similarly accomplished. Certainly the idea is worth consideration.\*



IN THE ORIGINAL description published in 1932, cases of acute regional ileitis were recognized and described. The clinical picture was similar to that of acute appendicitis, namely, generalized abdominal colic, pain, tenderness over the right lower quadrant, fever up to 101 or 102 F. The white blood count was elevated. A mass without active abscess formation was fairly constant. At operation, the findings were those of a mass of adherent and inflamed loops of ileum, the terminal coil being red, edematous, soggy and brilliantly congested, the mesentery of the gut was thickened and edematous and contained enlarged inflamed mesenteric lymph nodes. Clear or cloudy fluid in small amount was present in the peritoneal cavity. In rare cases, an abscess cavity containing thick aqueous but not foul smelling pus was encountered. A general statement was made that some cases resolved without operative interference, others passed transitionally into the chronic phase of the disease. It was presumed that the acute was but one phase of the chronic disease. The number of cases was too small and the follow up too recent to allow of any scientific statements other than to describe the clinical and pathologic picture. Jackman shortly thereafter reported 2 almost identical cases in which the acute picture of ileitis represented only a flare up of the process upon the basis of an old chronic inflammatory lesion in the terminal ileum. Both cases were successfully resected. Erb and Farmer reported 4 cases of acute ileo colitis in children. Meyer and Rossi described 8 cases of regional ileitis, 4 of which were of the

acute type, 3 of these 4 acute cases underwent spontaneous resolution. Rooley described a case of thickening of the terminal ileum apparently an acute regional ileitis in a child aged 5 years.

The ample literature that followed from the publications of many surgeons, Clute, Ladd, Harris, Bell and Brunn and our own present group in this series amply bore out the facts as set down in the earlier publication. Sufficient time and experience and follow up has, however elapsed to warrant a more complete clinical picture to elaborate upon the life cycle and prognosis of such cases. The surgical attack upon acute ileitis has passed through several phases and may now be recapitulated in its present and probably its final formulation.

The group in this present series comprised 16 cases, none of them more recent than one year. All of them observed a sufficient time to allow for the creation of a life history of this the acute phase of ileitis.

The sex distribution is about equal, the ages vary from  $2\frac{1}{2}$  to 37 years, an average of  $23\frac{1}{2}$  years at onset in contrast to  $27\frac{1}{2}$  years for the chronic phase of the disease. The difference in age incidence probably represents the gap between acute onset and the final development of the chronic picture. Many cases of acute ileitis occur in young children. Thus the well described case of Erb and Farmer in a  $2\frac{1}{2}$  year old child with complete autopsy findings, that of Harris in a 5 year old female child, Schiff 14 years of age. Warren and Sommers observed that the acute form of ileitis is more severe in young children since the diameter of the bowel is smaller and edema easily produces obstruction.

Three of 8 cases reported by Meade occurred in children 6, 8 and 9 years of age respectively. Megret's case involved a child 3 years of age with successful resection of the involved ileum at the time of exploration.

The pathologic involvement as seen at operation showed

the terminal ileum as the constant seat of involvement (table 14)

TABLE 14—*Extent of Pathologic Involvement*  
(Present Series)

4 inches of terminal ileum	2 cases
5 inches of terminal ileum	2 cases
6 inches of terminal ileum	1 case
8 inches of terminal ileum	2 cases
15 inches of terminal ileum	1 case
18 inches of terminal ileum	2 cases

Skip lesions as seen in chronic ileitis are not mentioned by the operating surgeons, an important observation, since it is a clear indication of the fact that the primary seat of the disease is regularly at the terminal ileum, the skip lesions representing a secondary and later extension in time and in site. The lesion at operation is characteristic, the terminal loop of ileum being bright red, edematous, soggy, congested, the mesentery thickened, the lymph nodes enlarged in some cases, not remarked in others. In one case, an abscess cavity was observed, surrounding a perforation of the terminal loop of ileum, the abscess containing fresh pus. The serosa of the ileum is covered by a sero-fibrinous exudate free fluid being seen in the peritoneal cavity. The pathologic or histologic findings in acute ileitis are almost identical with those of chronic ileitis. A greater degree of edema is present, leukocytic invasion is more prominent. The villi show loss of epithelium and become mere shadows. Very little slough of tissue is to be noted. The serosa is edematous, the blood vessels tremendously engorged. The lymph nodes are extremely edematous with marked hyperplasia of the germinal centers (HARRIS).

The clinical manifestations of acute ileitis show but little variation. The onset of symptoms varies from one to three days in most cases, extending up to five weeks. The acute, often fulminating, onset is characterized by pain in the general abdomen or in the right lower quadrant, the pain, colicky in nature present in all cases, accompanied

by fever, ranging from 100 to 103 F, the curve being constantly elevated and not remitting. Nausea and vomiting are not outstanding symptoms, each of them mentioned only in 2 cases—an important point in differentiation from acute appendicitis. A mass in the right lower quadrant of the abdomen was clearly palpable in 3 of the cases.

Diarrhea occurred in 11 of the 16 cases; constipation is clearly mentioned once only. The diarrhea may be and is often transient; may last one day, and be followed by normal stools and constipation, or may last throughout the acute phase of the disease.

Hematemesis and tarry stools occurred in one case; transient intestinal obstruction once. Other unusual manifestations are purpura hemorrhagica (once), erythema nodosum (once), arthralgias (once); these latter unusual signs and symptoms though rare in the acute phase link this abruptly with that of the more fully developed clinical picture of chronic ileitis.

The radiographic picture—an examination which is usually undertaken some days or weeks after the exploratory laparotomy—may demonstrate positively the involvement of the terminal ileum. In 8 of the 16 cases which were subjected to roentgen study the localization and the extent of the lesion could be clearly noted. The string sign so characteristic of the fully developed chronic phase of the disease is missing because the cicatrization and the contraction of the lumen of the ileum has not yet had time to develop. The mucosal pattern of the terminal loop of ileum is disturbed, the lumen narrowed, the proximal gut dilated, the contour irregular and shaggy. The extent of involvement as noted by the radiologic survey corresponds almost identically with that seen at operation (figs. 44 and 45).

The diagnosis concerns itself practically with the differentiation of acute ileitis from acute appendicitis. The one and only possible clue lies in the presence of or absence of diarrhea. Since as a rule ileitis is accompanied

by diarrhea (11 of 16 cases) and appendicitis by constipation. Unfortunately, both symptom complexes afford exceptions to the rule. Ileitis may be accompanied by constipation, though rarely (once in this series only), acute appendicitis may occur with and in the presence of diarrhea. To all intents and purposes, however, appendicitis is asso-



Fig. 44—Acute Terminal Ileitis Involving only Two Inches of the most Distal Ileum Proximal to the Ileocecal Valve. Confirmed by operation.

ciated with constipation and vomiting, diarrhea has been noted though exceptionally. Endemics of acute gastroenteritis with severe diarrhea may be complicated by a rare incidence of acute suppurative appendicitis (Crohn [60]).

Acute mesenteric lymphadenitis simulates both acute ileitis or acute appendicitis and cannot by any clinical symptoms be identified. Whether acute mesenteric lymph-

denitis and acute ileitis are phases of the same disease process if not identical, has been suggested and is still an open question (Jackson). Involvement of the terminal ileum in acute mesenteric lymphadenitis has been mentioned (Strombeck), but the transition of acute lymphadenitis to chronic regional ileitis has not been observed.



FIG. 45.—Acute Ileitis Spontaneous Healing. One of two brothers. The other brother with subacute onset developed the clinical picture of chronic granulomatous ileitis.

The prognosis in acute ileitis is good; only one case with fatal termination was observed following an ill advised attempt at resection during the acute severe toxic febrile state of the disease. Four cases were not subjected to any operation. 2 of these cases are well to date (4 and 5 years of follow up); one has passed into the chronic granulomatous stage; one has been lost to follow up.

The remaining 11 cases were explored; appendectomy performed in eight instances with three subsequent and

by diarrhea (11 of 16 cases) and appendicitis by constipation. Unfortunately, both symptom complexes afford exceptions to the rule. Ileitis may be accompanied by constipation, though rarely (once in this series only), acute appendicitis may occur with and in the presence of diarrhea. To all intents and purposes, however, appendicitis is asso-



Fig. 44—Acute Terminal Ileitis Involving only Two Inches of the most Distal Ileum Proximal to the Ileocecal Valve. Confirmed by operation.

ciated with constipation and vomiting, diarrhea has been noted though exceptionally. Epidemics of acute gastroenteritis with severe diarrhea may be complicated by a rare incidence of acute suppurative appendicitis (Cohn [60]).

Acute mesenteric lymphadenitis simulates both acute ileitis or acute appendicitis and cannot by any clinical symptoms be identified. Whether acute mesenteric lymph-

TABLE 16—Catalogue of Cases of Acute Ileitis from the Literature

	No. of Cases	Sub and nec	Died	Expected	Entire post mortem
1934 Jackson	2			1	
1935 Fieb and Farmer	4	2	"		
1936 Probst and Cruikshank	3	2			2
Meyer and Posa	5	3		2	
Koster et al	6	6			
1937 Brown I. W.	6	3			
Muller	2	1			
Pisgard and Henke	1	1			
1938 Kros I.	1	1			
1939 Lehman		2			
1940 Harris	1			1	
Flanagan and Johnson	1		1		
1941 Smith	2	4		1	
1942 Ebrill	1				
	—	—	—	—	—
	9	2	2	2	

Three questions arise: (a) Is acute ileitis a distinct entity or is it a part of the picture of chronic regional ileitis? (b) What is the consensus as to the eventual outcome of cases of acute ileitis? (c) What is the proper surgical or medical treatment of acute ileitis?

#### (a) *Distinct Entity or Part of Chronic Regional Ileitis?*

Acute ileitis is undoubtedly an individual phase of regional ileitis. It occurs most frequently as an initial phenomenon capable of complete subsidence in from 25 per cent to approximately 50 per cent of the cases, in the remainder advancing to the classic picture of chronic granulomatous ileitis. On the other hand in furtherance of the concept of identity of acute and chronic ileitis acute ileitis may often be regarded as an abrupt exacerbation of the chronic phase of the disease. In many cases a previous history of diarrhea and abdominal pain can be elicited if carefully sought after an occasional bout of temperature, a previous rectal operation. Out of this low grade generally unrecognized and unacknowledged course will occur an acute severe episode this exacerbation will again subside and the future course will be identical again with that



consequent abdominal wall fistulas (two of which closed spontaneously) Following the exploratory operation 2 of these cases became well and free of symptoms and of radiographic evidence of disease, the other 9 passed into the granulomatous phase of the disease Of these 9 cases which passed into the chronic inflammatory stage 4 are still not operated upon, 4 have undergone a short circuiting surgical procedure (ileo transverse colostomy) and are well, one was subsequently resected and is well The aftermath of these cases may be summarized as in table 15

TABLE 15—*Follow up of Acute Ileitis*

Unoperated—4 cases	Cases
Spontaneous resolution	—
Chronic granulomatous phase	1
No follow up	1
Operated—1 <sup>o</sup> cases	
Resection—death	1
Spontaneous resolution—well	0
Chronic granulomatous phase	4
Subsequent short circuiting operation	4
Subsequent resection	1
	—
Total	16

It will be noted that the operative procedures, apart from the initial surgical exploration, were all carried out months or several years subsequent to the acute phase of the disease Most of the cases during the acute onset were subjected to appendectomy only In only the one instance where death occurred was resection attempted during the initial grave onset of the acute ileitis, this death occurred elsewhere in spite of the caution and warning of the gravity of such a bold attack

The literature on acute ileitis is replete with good illustrative cases, most of which have been carefully observed and reported with prolonged follow up At least 50 to 60 cases have been adequately catalogued, the clinical picture following in most instances the pattern as originally described and as amplified and confirmed by subsequent authors (table 16)

TABLE 1b—Catalogue of Cases of Acute Ileitis from the Literature

	No. of Cases	Subsidence	Died	Excised	Enterostomy
1934 Jackman	2			1	
1935 East and Farmer	4	2	2		
1936 Probst and Gruenfeld	3	2			2
Meyer and Koss	3	3		2	
Kost et al.	6	6			
1937 Brown P. W.	6	3			
Maiden	3	1			
Bisvard and Henke	1	1			
1938 Gross I.	1	1			
1939 Lehman	1	1			
1940 Harris	1			1	
Eliaeson and Johnson	1	1	2		
1941 Smyth	3	4		1	
1942 Fibrill	1				
	39	33	5	3	—

Three questions arise: (a) Is acute ileitis a distinct entity or is it a part of the picture of chronic regional ileitis? (b) What is the consensus as to the eventual outcome of cases of acute ileitis? (c) What is the proper surgical or medical treatment of acute ileitis?

#### (a) *Distinct Entity or Part of Chronic Regional Ileitis?*

Acute ileitis is undoubtedly an individual phase of regional ileitis. It occurs most frequently as an initial phenomenon capable of complete subsidence in from 25 per cent to approximately 50 per cent of the cases, in the remainder advancing to the classic picture of chronic granulomatous ileitis. On the other hand in furtherance of the concept of identity of acute and chronic ileitis, acute ileitis may often be regarded as an abrupt exacerbation of the chronic phase of the disease. In many cases a previous history of diarrhea and abdominal pain can be elicited if carefully sought after, an occasional bout of temperature, a previous rectal operation. Out of this low grade generally unrecognized and unacknowledged course will occur an acute severe episode, this exacerbation will again subside and the future course will be identical again with that

of chronic regional ileitis (Brown, Lehman, Elfrson and Johnson, Smithy) Lvall even expresses the view that chronic regional ileitis represents only the end stages of acute ileitis with subsequent fibrosis and cicatrization "The process is primarily an acute one which tends to subside so that the appearance thought by some to indicate a chronic inflammation are really attempts at repair"

(b) *The Eventual Outcome of Cases of Acute Ileitis*

In my own series, only 25 per cent (4 cases) showed complete resolution as judged by subsidence of all symptoms, by the change of a positive x ray picture to that of a normal roentgenogram. The follow up, covering a sufficient number of years (10 to 14 years in some instances), allows of the presumption that these patients will continue well, even though we now know that recurrences of ileitis (after even supposedly curative operations) may occur after an equally long period of observation. In all of them, x ray studies were made soon after convalescence and the change from a positive to a negative x ray examination was noted. The time required for healing, or for a negative roentgenogram varied from 1 month to as long as 14 months in the most protracted case. The gradual disappearance of the lesion could be ascertained by comparing successive x ray exposures. Meyer and Rosi noted spontaneous resolution in 3 out of 4 cases of acute ileitis. They controlled their observations by radiographic studies, in one case involving a child of 7 years of age, seven weeks after the exploratory operation the involved ileum appeared roentgenographically negative and free of disease. Another case in an adult female was well one month after the operation radiographic examination revealed a normal terminal ileum. An adult colored male was observed for 1½ years after exploratory operation for acute ileitis and was and remained radiographically negative for any residual signs of the previous inflammatory focus in the terminal ileum.

In the series of Brown and Donald 7 out of 10 of the acute cases were well, 6 of them 3 or more years after the

simple exploratory operation. The series of Witten and Sommers included 15 cases of acceptable acute ileitis. 12 of these, or 80 per cent regressed, 2 progressed and one died. McKinnon reported the spontaneous cure in a case of acute regional ileitis, 2½ years after the exploratory operation the patient was well and the radiographic examination, which previously had shown involvement of the terminal ileum, was now entirely negative.

In the literature, the percentage of spontaneous resolution is considerably higher than in my present series. Thus table 16 shows resolution in 33 out of the 59 cases collected from the current publications. One may honestly question the validity of the follow up in some of these published cases. Quite obviously, many of the reported cures had only recently been observed and often blanket statements are made as to complete resolution without adequate control by radiographic study and long period observation. Nevertheless, the subsidence and complete cure of acute ileitis is convincingly shown by most of the cases in the reported literature, thus, Meyer and Rosi in a follow up of 1 month, 7 weeks and 1½ years, Brown 2 years 3 years and 4 years. Bisgard and Hencke 2 years. Rixford in discussing a paper by Mixer cites a case which had remained well for 8 years after an exploratory operation. In the series reported by Eliason and Johnson 4 cases have remained well for 2 to 6 years, 4 additional cases for not less than 1 year. One case by Smithy was well 6 years after exploration the second case showed some radiographic evidence of the disease 6 weeks after exploratory operation but was well with a negative x ray examination 13 months later. In Lehman's series of 7 cases, 4 have been free of symptoms of the disease from 14 months to 9½ years after the exploratory operation. Kross followed his one case for seven years of complete restitution. Fugh reports 2 cases of acute ileitis with spontaneous recovery. One case was operated upon and acute terminal ileitis observed, 6 months later the case was reoperated and scarcely any evidence of the original lesion

was found. In the second case, after an operation for acute ileitis, the patient was followed by repeated x-ray examination until a completely normal bowel was observed roentgenographically. Probst and Gruenfeld described acute ileitis of terminal ileum with severe hemorrhage, enterostomy. Three months later, at closing of the fistula, the process appeared healed. A small section of the previously involved ileum was removed for biopsy, the microscopic sections revealed only slight round cell infiltration of the subserosa. In a second case, only an enterostomy was performed for severe terminal ileitis (15 cm). Two years later the patient was entirely well.

It is presumed in the literature that all those cases which did not advance to spontaneous healing gradually developed into instances of chronic regional ileitis. Meyer and Rosi make the point that when the mesentery is involved, spontaneous resolution is less likely to occur. The experience of Lehman, however, contradicts this opinion since in 1 of his cases with involvement of the mesentery, spontaneous resolution did occur, in another, with no mesenteric involvement, the after course was one of chronic regional ileitis.

### (c) *The Medical or Surgical Treatment of Acute Ileitis*

The medical treatment of acute ileitis consists of the administration of a low residue diet, sulfasuxidine or sulfathaladine, over a long period of time, supplemented by the use of streptomycin. The efficacy of these antibiotics in effecting the cure may be questioned, since therapeutic conviction is not possible in the presence of a disease that shows ability spontaneously to resolve. The sulfa drugs may be continued indefinitely without danger. streptomycin 2 grams daily by intramuscular injection, supplemented by 1 gram per os, is certainly worth a trial. In many cases of acute fulminating ulcerative colitis, the value of streptomycin therapy is often strikingly convincing.

The surgical therapy at the time of the exploratory operation resolves itself into three possible courses of action

(1) Simple closure of the abdomen (2) Appendectomy alone closure of abdomen without drainage (3) A radical procedure (resection or short circuiting)

1 Where the preoperative diagnosis can be made with any sense of assurance such as the presence of diarrhea or a possible previous history of diarrhea, fever, or a rectal fistula, under such circumstances no exploration is justified. In the absence of vomiting or rebound tenderness or of masses, with low fever and a low leukocyte count, exploratory operation should be avoided. If operation is undertaken under a false diagnosis simple closure of the abdomen without drainage is advocated.

2 If, however the abdomen is opened and the exploration reveals an acute ileitis, shall the appendix be removed? There seems little justification for the removal of an appendix. Though the practice of appendectomy is common under such circumstances, the pathologic report if scrutinized is invariably either negative or that of 'chronic inflammation'. Ileitis does not affect the appendix, an organ which is attached to and constitutes a ventral prolongation of the cecum. The cecum is free of disease in all but exceptional combined types of the disease. The appendix is affected only when involved in a large inflammatory mass of loops of ileum or in an abscess derived from a localized perforation of ileum. This does not constitute acute appendicitis and does not call for appendectomy. It is questionable whether the removal of the appendix is a disadvantage that leads to abdominal wall fistula. In our series of 16 cases of acute ileitis 9 cases underwent exploratory operation in 3 instances the appendix was not removed with no fistula resulting. In 6 instances the appendix was removed in 3 of which abdominal wall fistulas were noted as a sequel. Fortunately, 2 of the 3 fistulas healed spontaneously.

Certainly, the removal of the appendix has no curative value as to the course of the disease. Mixer advised against appendectomy. Flaxson and Johnson favor appen-

dectomy only if the cecum and appendix seem to be involved in the pathologic process. In their series of 15 cases, the appendix was removed in all, yet but one fistula resulted. Drainage of the abdominal wound seems unnecessary and tends to fistula formation. Smithy feels that appendectomy is purely incidental, does not increase the risk of fistula formation, he feels that to neglect appendectomy may invite later involvement of the appendix in the pathologic process with eventual development of acute suppurative appendicitis.

My own experience does not favor appendectomy at the time of exploration, because such a procedure is unnecessarily superfluous and tends to fistula formation. Acute gangrenous appendicitis as a complication of chronic regional ileitis occurred subsequently only twice in this series of 222 cases.

3 Is a radical procedure, such as resection or a short circuiting operation ever indicated at the time of exploration? The earlier surgeons with experience in this field, often advocated radical resection at the time of exploration. Mixer was one of these earlier advocates of such a procedure, though his mortality was extremely high. Jackman reported 2 cases of acute ileitis with resection at the time of exploration both patients recovering. Harris reported what he thought was the earliest acute stage of regional enteritis in a female child, aged 5 years, with a short history of 3½ days. Resection was accomplished without complications or shock and resulted in cure. Probst and Gruenfeld performed simple enterostomy in 2 cases of acute ileitis, with recovery in both instances as observed in the later examinations.

Since the establishment of the fact that spontaneous healing was not only possible but likely in a large percentage of cases of acute ileitis, the consensus favors expectant waiting and noninterference (Koster et al, Brown, Mailer Eliason and Johnson, Smithy, Waugh, Meyer and Rosi).

That resection or a short circuiting procedure may even

tully become necessary is evident in the follow up of cases of acute ileitis. In our own series of 16 cases, 4 were eventually subjected to ileo transverse colostomy, one to resection of the chronic lesion, 4 cases are well along in the chronic phase but have as yet not consented to operation because of long periods of remission with freedom from symptoms. The necessity for later operation has been noted by many of the current writers (Meyer and Rosi, Eliason and Johnson, Ebbell), and is a logical sequel to the chronic granulomatous phase of the disease.



ILEO JEJUNITIS may be defined as that form or type of chronic enteritis in which the granulomatous process typical of that disease involves the jejunum either as an extension from the ileum or as independent lesion limited to the upper segments of the small bowel

Very soon after the original publication in 1932 of regional or terminal ileitis, it was recognized by Harris, Bell and Brunn that the upper ileum and the jejunum could be and were also involved in an almost identical pathologic process. The literature, year by year, has added new case histories most of them on the regional form of jejunitis. As early as 1934, Brown, Borgen and Weber reported 3 cases of enteritis with the lesion restricted to the jejunum. In 1936 Meyer and Rosi mentioned a case with the inflammatory process localized in the jejunum. In 1937 Pemberton and Brown already noted that 3 of 37 cases of regional enteritis were entirely restricted to localized segments of the jejunum. Tikelv and Lisa reported in detail 2 cases of high jejunitis, resected, ending fatally and carefully studied at autopsy. These were both localized lesions high in the jejunum and causing clinical intestinal obstruction. Rardin and Johnston in 1939 cited 16 cases of sole involvement of the jejunum in cases culled from the literature.

In 1941 Crohn and Lunich published an analysis of 17 cases of diffuse ileo jejunitis with a clinical analysis of the outstanding symptoms and the therapeutic approach to the problem. The differentiation of ileo jejunitis from

sprue was particularly emphasized. Sussman and Wachtel added the distinguishing roentgenographic features in an analysis of 23 cases of granulomatous jejuno ileitis.

Jejunitis or ileo jejunitis is not a separate or different disease, in all of its major characteristics it is but a sub




Fig 46—Distribution of Granulomatous Lesion in Ileo-Jejunitis (38 Cases). (A) Ileum and lower jejunum (1 case). (B) diffuse ileum and jejunum (2 cases). (C) diffuse ileo jejunitis with involvement of duodenum (2 cases). (D) upper ileum and lower jejunum (6 cases). (E) localized jejunitis (3 cases). (F) terminal ileum and upper jejunum (3 cases). Group B represents the greatest number of cases.

type of the more common form of regional ileitis. In its pathology as a granulomatous lesion, it is in no major way different from ileitis except in its anatomic distribution, one notes the same clinical characteristics, fever, diar-



Fig. 47—Terminal Ileitis and Typhinitis by Continuous Involvement in a 21 Year old Man. Conservative medical therapy over a course of four years.

rrhea, a mass by palpation, fistulas, rectal complications, anemia. But the degrees of involvement are different: the pathologic process is spread over a wider area, the local intensity is less marked in any one segment, fistulas and rectal complications are fewer. But the general constitutional disturbance is greater: nutritional absorption is markedly impaired, fever, the enlarged spleen, the club

ling of the fingers indicate the degree of absorption of the toxic products of the disease skeletal growth is impaired Hypoproteinemia is a characteristic of the advanced pathologic state



Fig. 48—Involvement of Lower Jejunum and Upper Ileum At onset gross hemorrhage melena diarrhea and continuous fever

The course is low grade often less severe a considerable degree of spontaneous healing is possible the disease is less amenable to surgical therapy than is terminal or regional ileitis

Ileo jejunitis should be considered and studied in its various forms in its anatomic and pathologic variations as illustrated in the 28 cases of ileo jejunitis which constitutes the present series (fig. 46)

1 As an extension upward of an etiologically older process in the ileum

2 As a diffuse primary ileo jejunitis involving all of the ileum and the jejunum in its original distribution




Fig. 49—Pathologic Involvement of Upper Ileum and Jejunum

3 As constituted solely by a localized involvement of the jejunum in whole or in part

4 As isolated jejunal involvement plus another isolated segment of involved ileum the two processes being entirely discontinuous

1 Involvement of the Jejunum by Upward Extension from a Chronologically Older Process of Regional Ileitis

The usual form of this type is one of older terminal or regional ileitis in which by skip lesions or by continuous

extension the jejunum partakes of the same process which had originated in the lower ileum. Only 2 of the 36 cases in this series illustrate this type of extension, the older process chronologically resides in the terminal ileum (fig. 47). In these cases the severity of the process is so great that either initially or in the course of time the process of disease extends upward to involve the whole ileum and some of the lower jejunum usually as a continuous process (fig. 48). Jejunitis as a complication of ileitis is also seen after unsuccessful surgical procedures in which repeated recurrences take place. Following a resection or a short circuiting of the initial terminal ileitis a recurrence takes place in the new terminal segment of ileum. Another resection is carried out followed by a new recurrence. A third resection may occasionally be attempted and further recurrences may now carry the process up into the jejunum for the sacrifice of involved recurrent ileum may necessitate the resection of a wider area until the whole ileum has been dispensed with and the jejunum is invaded by the malignant extension of the disease process.

## 2 *As Diffuse Ileo jejunitis*

This is the most common form seen and is constituted by a group of 24 of the total 38 cases under study. Here the whole ileum and the whole or most of the jejunum are involved in a continuous process. Occasionally, in the uppermost jejunum, the process is interrupted by the appearance of skip lesions separated by skipped areas of normal mucosa the lower jejunum and the whole ileum being the seat of one continuous process of disease (fig. 49). In 2 of these cases the duodenum fourth and third portions were also involved by extension as seen radiologically and at operation.

Chronologically it is impossible to determine which segment is the primary seat of the disease. In practically all of this group the disease had universally involved the whole of the small bowel when first recognized so that it was impossible to state whether the initial process was in

the upper jejunum or in the lower ileum (fig 50) Presumably, in this severe type, the whole small bowel is



Fig 50—Pathologic Lesion Involving Lower Jejunum Fistula between loops of small intestine

simultaneously involved. Twice, by radiologic examination, the stomach was suspected of participating in the process, but such involvement was never proven by either gastroscopy or by exploratory operation. Barbour and Stokes in 1936 described a case which at autopsy was shown to have 13 areas of alternate thickening and thinning of the




Fig 51—Localized Jejunitis Occupying Uppermost Area of Jejunum. Alternate constrictions and dilation, enlarged mesenteric lymph node. Characteristic cobblestone appearance identical with that seen in the terminal form of ileitis.

entire small intestine from pylorus to the ileocecal valve. Holloway's Case 5 showed definite involvement of the third part of the duodenum as well as the terminal ileum and intermediate segments. Rees has described the case of a 30 year old Negroess who at operation presented the picture of multiple intestinal obstruction at three points in the upper jejunum due to segmental jejunitis. resection en masse was surgically successful.

### 3. As Isolated Jejunitis

This form is constituted by 9 of the total 28 cases in the



the upper jejunum or in the lower ileum (fig 50). Presumably, in this severe type, the whole small bowel is



Fig 50—Pathologic lesion involving lower Jejunum. Fistula between loops of small intestine

the duodenum was involved in this instance. In a second case the terminal duodenum and 18 inches of proximal jejunum showed "thickened and beefy walls", here gastro jejunostomy was performed. In the third case, 2½ feet of the proximal jejunum showed, on exploration, chronic inflammatory changes, again, gastro jejunostomy was performed. Cases of primary regional enteritis involving the jejunum following trauma were reported by Blowne and McHardy and by Spellberg and Gray.

#### 4 *The Combination of Isolated Jejunitis Plus an Isolated Segment of Ileitis (3 cases)*

In this type one notes two widely separated and discontinuous lesions, one in the upper jejunum as a sole lesion occupying 8 to 10 inches of jejunum, then far down usually in the terminal ileum is seen another separate and independent lesion of ileitis, the intervening mucosal area being free of extensions or of skip lesions. In a case of W. E. Ladd cited by Homans and Hass a child had two diseased loops of small intestine, one in the jejunum and the other in the distal ileum (1933). Corr and Boeck as early as 1934 published a case of ileitis of 6 years' duration, the ileitis beginning 3 feet above the ileo cecal valve. A second lesion high in the jejunum was also present, the whole picture being complicated by a recto vaginal fistula. It is again impossible chronologically to state the relationship in time or in sequence between these two areas since they were both present when the diagnosis was first established. Presumably both areas of disease were present from the initial onset of the pathologic process for from our study of regional ileitis it would be deduced that were the higher lesion an extension of the older process in the terminal ileum intervening skip lesions would have been observed. This viewpoint is the logical deduction which follows from the study of the life cycle and the methods of extension of chronic regional ileitis.

These variable segments of involvement in ileo jejunitis either as continuous, as isolated areas or as combination

group In 3 of them, isolated areas of jejunum solely were the seat of the disease, one was 4 inches, one 2 feet and one 4 feet from the fossa of Treitz, a small isolated segment of jejunum 6 to 8 inches in length being the seat of the pathologic process (figs 51 and 52) In the other 6 cases,




Fig 52—Localized Jejunitis Hammock like dilation between two areas of constriction Successful resection See figure 50 representing resected specimen

the lower jejunum was alone involved but the process was observed to extend also into the uppermost segment of the ileum

Instances of isolated jejunitis have been published by Johnson, by Brewster, by Gendel and Beaver, by Kohn, and by Lyons and Garlock James recently published 3 cases of regional jejunitis of unusual interest In one case, the whole jejunum from duodenum to ileum was involved An anastomosis was made between the stomach and the upper ileum in order to short circuit the lesion,

No new or significant factors as to the causation of ileo jejunitis are known as distinct from regional ileitis. The same fundamental ignorance as to causation and etiology which covers ileitis applies also negatively to ileo jejunitis. In none of the 36 cases in the group which forms the basis of this study was physical trauma attributed as a causative factor. Trauma, however, has exceptionally been recited in the literature (Browne and McHardy, Spellberg and Gray).

### PATHOLOGY

The pathology of ileo jejunitis differs but little from that of regional ileitis. Grossly the small intestine is bluish red in appearance the serosa streaked by inflamed lymphatics. The wall is thickened, coarse and firm, where long standing healing has taken place the lumen may be contracted and the wall of the gut puckered. On opening the resected specimen the mucosa is cobble stone in appearance as seen in ileitis cross hatching of longitudinal and transverse ulcerations creating that appearance. The mucosa is thrown up and prominent in places appearing as and even assuming a polypoid excrecent form. True inflammatory polyps occur frequently and can be visualized in the roentgenograms. The ulcers in the mucosa penetrate to the inflamed thickened submucosa these ulcers rarely penetrate more deeply and do not with rare exception form fistulous tracts. The whole wall submucosa muscularis and serosa is involved in the granulomatous and cicatricial process as evidenced by thickening edema congestion. Since much healing is capable in the upper small bowel, areas of contraction, moderate or extreme may appear with intervening dilated and distended loops of intermediate bowel (fig 53). Where two or more areas of constriction occur the intermediate bowel may be much distended. *This over distended loop may give rise to necrosis of the bowel wall with abscess formation.* Or a perijunal abscess may form in the thickened and inflamed mesentery the abscess being walled off and localized. When

types constitute the various forms in which the disease occurs. The symptoms, the physical signs, the complications are more or less the same in all the forms, the radiologic study constitutes a means for the identification and recognition of the anatomic distribution, as checked against the findings at the exploratory laparotomy where such a surgical confirmatory procedure has been invoked.

### ETIOLOGY

In this series of 36 cases, 11 were females, 27 were males, indicating a preponderance of male involvement of almost 3 to 1. It is possible that the series is too small to allow of a scientifically accurate conclusion, but throughout the series the major involvement of males is a striking characteristic. (In regional ileitis the proportion of males to females is 5 to 4.) Sussman and Wachtel found a similar incidence of 3 males to 1 female.

The age incidence in this series is striking as indicating a preponderance of involvement in the earlier decades of life (table 17).

TABLE 1 —*Ileo Jejunitis* Age Incidence

Age at onset Years	No of Cases
1-10	1
10-20	17
20-30	9
30-40	7
40-50	1
50-60	1
60-70	0
Total	36 cases

The main brunt of the disease seems to fall in the second decade of life, the third and the fourth decades together contributing an almost equal number. After the age of 40, only a rare exceptional case is seen. The extreme youth of most of these cases of ileo jejunitis is a striking fact. In the series of 23 cases of granulomatous jejuno ileitis published by Sussman and Wachtel, the average age incidence was 27.0 years, the range varying from 9 years to 65 years of age.

of the adjoined mesentery, the subacute stage by a granular and hypertrophic mucosa containing punched out ulcerations with smooth or irregular edges the ulcers being either longitudinal or transverse to the axis of the bowel early polypoid proliferation and evident skip lesions are discernible. The chronic stage is characterized by thickening and fibrosis alternating areas of atrophic and hypertrophic mucosa, multiple areas of stenosis the serosa being covered by a shaggy exudate. This description includes the findings at four autopsies and the intestinal biopsies obtained at three of eight exploratory laparotomies and represents an accurate and painstaking study of the pathologic changes in the three various phases of the disease.

Several forms of isolated jejunal ulcers appear in the literature, though there is some question as to the identity of such ulcers with the granulomatous forms which are recognized as ileo jejunitis. Richardson in 1922 was responsible for the first American description he cited 2 cases of jejunal ulcer with stricture and perforation. Buckstein gave a good description of primary ulcer of the jejunum. Robinson and Wise in 1940 reviewed the whole literature and culled the large number of 36 cases of solitary ulcers of the small bowel of which 7 were jejunal. Dowdle adds a case of multiple primary nonspecific jejunal ulcers with duodenal dilatation the bowel was obstructed not much below the ligament of Treitz with dilatation and thickening of the intestine above the strictured area.

#### SYMPTOMATOLOGY

An acute form of ileo jejunitis or phlegmonous jejunitis occurs in the literature though personally I have never seen such a case. An excellent description is given by Clark and Wright of 2 personally observed cases of acute phlegmonous enteritis the first case was characterized by an acute onset a mass in the left upper quadrant of the abdomen high fever leukocytosis it terminated in death. At autopsy an area of acute jejunitis was seen 30 cm from the pylorus and extending distally for 20 cm. In the

such an abscess forms, it may become adherent to the abdominal wall at the site of the scar of a previous incision and discharge its content. Fecal fistulas to the abdominal wall are, however, rare. The lymph nodes of the mesentery are enlarged and succulent, similar to those seen in distal terminal ileitis. Microscopically, the process is identical with that of ileitis and of intestinal granulomata in gen-



FIG. 53—Diffuse Ileo Jejunitis with Major Involvement of Jejunum. Alternating areas of constriction and dilation scattered throughout the small intestine. Marked hypoproteinemia, severe diarrhea, progressive loss of weight. Prognosis poor.

eral. Miliary like tubercles consisting of lymphoid and epithelioid mononuclear and plasma cells are common and frequently enclose giant cells, presumably of foreign body nature. The lymphoid follicles are hyperplastic with active germinal centers. In their excellent article, Sussman and Wachtel describe the pathology as well as the radiologic appearance of the disease as occurring in three stages, namely acute, subacute and chronic. The acute stage is characterized by edema and thickening of the mucosa and

of the adjoined mesentery, the subacute stage by a granular and hypertrophic mucosa containing punched out ulcerations with smooth or irregular edges, the ulcers being either longitudinal or transverse to the axis of the bowel, early polypoid proliferation and evident "skip lesions" are discernible. The chronic stage is characterized by thickening and fibrosis, alternating areas of atrophic and hypertrophic mucosa, multiple areas of stenosis, the serosa being covered by a shaggy exudate. This description includes the findings at four autopsies and the intestinal biopsies obtained at three of eight exploratory laparotomies and represents an accurate and painstaking study of the pathologic changes in the three various phases of the disease.

Several forms of isolated jejunal ulcers appear in the literature, though there is some question as to the identity of such ulcers with the granulomatous forms which are recognized as ileo jejunitis. Richardson in 1922 was responsible for the first American description, he cited 2 cases of jejunal ulcer with stricture and perforation. Buckstein gave a good description of primary ulcer of the jejunum. Robinson and Wise in 1940 reviewed the whole literature and culled the large number of 6 cases of solitary ulcers of the small bowel, of which 7 were jejunal. Dowdle adds 1 case of multiple primary nonspecific jejunal ulcers with duodenal dilatation. The bowel was obstructed not much below the ligament of Treitz with dilatation and thickening of the intestine above the strictured area.

#### SYMPTOMATOLOGY

An acute form of ileo jejunitis or phlegmonous jejunitis occurs in the literature though personally I have never seen such a case. An excellent description is given by Clark and Wright of 2 personally observed cases of acute phlegmonous enteritis. The first case was characterized by an acute onset, a mass in the left upper quadrant of the abdomen, high fever, leukocytosis. It terminated in death. At autopsy an area of acute jejunitis was seen 50 cm from the pylorus and extending distally for 20 cm. In the



such an abscess forms, it may become adherent to the abdominal wall at the site of the scar of a previous incision and discharge its content. Local fistulas to the abdominal wall are, however, rare. The lymph nodes of the mesentery are enlarged and succulent, similar to those seen in distal terminal ileitis. Microscopically, the process is identical with that of ileitis and of intestinal granulomata in gen-



FIG. 53—Diffuse Ileo Jejunitis with Major Involvement of Jejunum. Alternating areas of constriction and dilation scattered throughout the small intestine. Marked hypoproteinemia, severe diarrhea, progressive loss of weight. Prognosis poor.

eral. Miliary like tubercles consisting of lymphoid and epithelioid mononuclear and plasma cells are common and frequently enclose giant cells presumably of foreign body nature. The lymphoid follicles are hyperplastic with active germinal centers. In their excellent article, Sussman and Wachtel describe the pathology as well as the radiologic appearance of the disease as occurring in three stages, namely acute, subacute and chronic. The acute stage is characterized by edema and thickening of the mucosa and

fever, (b) diarrhea, (c) severe abdominal pain, (d) gross hemorrhage, (e) symptoms of obstruction, (f) anorexia, (g) loss of weight, (h) anemia

*Fever* was present in 17 of the 36 cases in this series, the fever is intermittent in type, higher in the late afternoon or evening, may rise to 103 F, though usually the range is between 100 and 102 F temperature, chills are occasionally noted. The fever may be so mild as to be almost imperceptible and may not be accompanied by malaise or prostration. The fever may be paroxysmal remaining continuously for several weeks or months, or may be interrupted by periods of apyrexia.

*Diarrhea* is the most constant symptom, occurring in 31 of the 36 cases in this series, constipation is noted once only, normal bowel movements in the remaining four instances. The diarrhea is never extreme, averaging usually two to four movements per twenty four hours. The stools are mushy or semisolid, mixed with thin mucus, occasionally watery. Pus is never seen, gross blood in flecks only exceptionally. The guaiac reaction was positive in 10 cases, negative in the remainder. In fact, it is not unusual to have negative guaiac reactions in the feces of proven cases throughout the course of the disease.

*Abdominal pain* is another almost constant symptom, having occurred in 30 of the 36 cases in this group. The abdominal pains are crampy in nature, diffused over the abdomen, or more likely to be localized over the left upper quadrant. The pains may be mild and associated with the desire to defecate or severe and definitely predefecatory in timing. With the passage of fluid feces and gas accompanied by an audible gurgling the cramp is allayed. When obstructive phenomena are present, the abdominal cramps are more or most severe occasionally accompanied by vomiting.

*Gross hemorrhage* occurred once in this present series. In a previously studied series it occurred three times as an initial symptom out of 17 observed cases. The gross

second case, the acute lesion was 50 cm above the ileo cecal valve in the upper ileum, the area involved again extending for 20 cm into the jejunum. They reviewed the literature and cite 12 cases in the duodenum or proximal jejunum, 22 cases in the jejunum alone, and 7 in the ileum, many of which, however, resembled granulomata and were associated with foreign bodies, fish bones and diverticula.

Very recently, Husebye presented 14 cases of acute phlegmonous jejunitis as seen by x-ray, in six instances the diagnosis rested essentially on the roentgen appearance of the films, seven of the fourteen patients recovered. The roentgen findings were characteristic and consisted of (1) Reduced diaphragmatic movements on one or both sides (2) Scout films revealed distended loops of gut over the left abdomen (3) Erect films revealed fluid levels (4) By barium meal, the mucosal relief was roughly jagged with irregular circular folds partly sharp and partly blunt.

Apparently, this form of acute phlegmonous jejunitis is an extremely severe one with a high mortality and with grave symptomatology. Involvement of the duodenum as part of a high jejunitis has recently been described by James in 2 out of 3 cases. In one case, the process involved the distal duodenum and the whole of the jejunum. In a second case, the process involved the terminal duodenum and 18 inches of the upper jejunum, the duodenum just at the fossa of Treitz being kinked and adherent to the loop of jejunum.

The classic form of chronic ileo jejunitis or jejunitis is that of a low grade granulomatous inflammatory lesion, involving primarily the jejunum either continuous with a similar process in the ileum or associated with an independent lesion in the distal ileum.

The onset is usually gradual and prolonged, lasting from six months to fourteen years or more. The actual onset is often difficult to estimate since the occasional slight diarrhea, mild fever or loss of weight is little noticeable.

The outstanding clinical symptoms were (a) low grade

case, who had remained stunted in growth and devoid of secondary sex characteristics later recovered fully from the diffuse ileo jejunitis from which he had suffered, grew to be a strong, well developed man married and was the father of two children.

The menses are rarely skipped in ileitis as they are in severe ulcerative colitis. Several cases of ileo jejunitis have married, borne normal children without detriment to their general health or difficulty, and without the induction of a recurrence of the disease process.

Aphthous stomatitis (3 cases) furunculosis, erythema nodosum, epidermolysis bullosa were each seen once. Phlyctenular conjunctivitis was twice seen. Manifestations of hypovitaminosis are not often seen, occasionally, one notes bleeding gums glossitis as evidence of nicotinamide deficiency, riboflavin deficiency is marked by cheilosis though these are extremely rare. Vitamin A deficiencies night blindness, the bone changes of vitamin D deficiency have not been noted.

Killian and Ingelfinger<sup>11</sup> studied the nutritional problems presented by a patient with extensive ileo jejunitis. A 23 year old boy with a previous history of a loss of 40 pounds of weight was torpedoed during the recent war and was exposed in a life raft for eight days. He was found in a state of extreme malnutrition emaciated dehydrated with pitting edema of the lower extremities. The tongue was smooth, fever up to 102 F, hemoglobin 54 per cent, red blood cells 1 200 000 hemitocrit 22 per cent. Serum protein was markedly reduced to 4.85 grams per 100 cc. albumin 2.34 globulin 2.51 grams per cent. Clubbing of the fingers was present, the glucose tolerance test showed hypoglycemia with a low tolerance curve. By x ray examination the lower jejunum and the ileum showed marked irregularities and points of narrowing with localized dilatation and constriction. Serum calcium was 8.0 and 8.2 milligrams per cent. Inorganic phosphorus 2.5 and 2.8 milligrams. Vitamin A absorption of 20,000 I. U. units was only slightly below normal.

hemorrhage takes the form of melena, though once it was accompanied by a brisk hematemesis. The hemorrhage while profuse is never sanguiniform, and usually not associated with shock.

*Symptoms of obstruction* are frequent and occurred in a mild form in 10 of the 36 cases. The intestinal obstruction is of low grade, none of the cases of this or of our previous series of 17 cases, having come to operation because of advanced obstructive symptoms. The obstruction is usually inflammatory in nature and is due to edema of the bowel wall, possibly accompanied by a perijunal or a mesenteric abscess and subsides as the inflammatory process recedes. Vomiting was noted in 5 cases. Browne and McHardy mention obstruction as one of the symptoms in their case of a Negro boy who developed a localized jejunitis following a severe abdominal trauma.

*Anorexia and weight loss* are common manifestations. The loss of appetite is spotty and occurs during acute exacerbations of the chronic process, at other times appetite may be well maintained.

The weight loss may be very severe amounting in some of our cases to as much as 50 pounds in a stout girl. The average loss of weight in our previous series was 17½ pounds. During the healing phases, with remissions of the fever and diarrhea the weight may be rapidly regained, 15, 25, or even 50 pounds may be added during prolonged remissions of the symptoms.

The *nutrition* is usually poor, skin dry, mucous membranes pallid. An extreme degree of malnutrition with emaciation and loss of strength marks the end stages of lethal cases. Retarded growth and retarded sex maturation are commonly observed in those young adolescents who are so frequently affected by the disease at a prepubertal age. With the recovery from the severe phase, and with healing of the lesion advancing weight, growth in height, and secondary sex maturation may replace the hypogonadism seen in the adolescent years. One of our earlier

little from the normal, a very mild leukocytosis up to 8,000 to 10,000 or even 18,000 white blood cells may be noted though most usually the count is within the lower range of normal. Eosinophilia as seen in ulcerative colitis is unusual, though it was seen in 3 patients of our previous series, up to 14 per cent of eosinophiles being present in the blood smears. A mild secondary anemia was present in 12 out of the 36 cases in this series, the range being usually between 72 and 80 per cent hemoglobin (Sahli). A macrocytic hyperchromic type of anemia occurred once in our previous series, with a color index of 1.3. In this series of 36 cases hyperchromic anemia was not observed even though the anemia was at times severe and the intestinal absorption severely compromised as noted by hypoproteinemia. The hyperchromic anemia in ileo jejunitis is usually not beneficially influenced by intramuscular injections of crude liver extract the reticulocyte response attaining a level of only 7 per cent this in contrast to its more successful therapeutic result in cases of nontropical sprue.

The occurrence of macrocytic anemia in association with intestinal lesions particularly intestinal strictures and anastomoses was amply demonstrated in the literature by Faber Hale White by Meulenracht and by Huist. Butt and Watkins mention 7 cases of diffuse ileitis with macrocytic anemia in 6 of which a fecal fistula co existed. With proper reconstructive surgical intervention the macrocytic type of anemia could be returned to a normal blood picture. They rightly ascribe the hyperchromic anemia to the abnormal interference with absorption and improper utilization of the effective hemopoietic fraction of the diet. Plum and Warburg in a publication entitled Hematological Changes Especially Megalocytic Anemia in Regional Ileitis describe 4 cases under their personal observation in which the blood picture was typical of that of a hyperchromic anemia. The hemoglobin values ranged as low as 20 per cent red blood cell count as low as 1 410 000 cells.

levels, fasting serum carotene almost normal. The patient improved rapidly under high protein diet and amino acid concentrates. The intravenous administration of amino acids did not improve the nitrogen balance or create nitrogen retention in the metabolism studies.

Loss of fat soluble vitamins was especially emphasized in Albright and Stewart's report. The case, a woman aged 30 years had undergone four successive operations for ileitis and recurrent ileitis, including two extensive intestinal resections. Prolonged severe abdominal cramps and diarrhea resulted in tetany and a state of marked vitamin deficiency. The prothrombin time was found to be only 13 per cent of normal, indicating a marked associated vitamin K deficiency. Vitamin C was within normal range. Fat soluble vitamin A was somewhat deficient 82 units per 100 cc of serum (normal 10-20 units) carotenoids somewhat diminished, and associated with a xeroderma of the breast and abdomen. Fat soluble steroids, particularly the quantity of 17 keto steroids in the urine were practically absent. The tetany responded rapidly to large doses of vitamin D, 500,000 USP units daily. The results of treatment in the form of liberal dosage with all of the fat soluble vitamin preparations was clinically dramatic.

An unusually exaggerated degree of hypoproteinemia was noted in a diffuse case of ileo jejunitis by Casten. At a previous operation an anastomosis between the upper jejunum and the transverse colon had been performed because of a diffuse inflammatory process involving all of the ileum and the lower jejunum. Extreme malnutrition and evidences of hypoproteinemia were soon evident. The total serum proteins fell to 3.1 grams per 100 cc, the albumin fraction being 1.1 Gm and the globulin 2 Gm (albumin globulin ratio 0.55). Most observers conclude that the critical level of serum protein is 5.0 Gm per 100 cc, edema will invariably result when the blood level falls below that point.

*Blood changes* The white cell count varies only a

little from the normal a very mild leukocytosis up to 8,000 to 10,000 or even 18,000 white blood cells may be noted, though most usually the count is within the lower range of normal. Eosinophilia, as seen in ulcerative colitis is unusual though it was seen in 3 patients of our previous series, up to 14 per cent of eosinophiles being present in the blood smears. A mild secondary anemia was present in 12 out of the 36 cases in this series the range being usually between 72 and 90 per cent hemoglobin (Sahli). A macrocytic hyperchromic type of anemia occurred once in our previous series, with a color index of 1.3. In this series of 36 cases, hyperchromic anemia was not observed even though the anemia was at times severe and the intestinal absorption severely compromised as noted by hypoproteinemia. The hyperchromic anemia in ileo jejunitis is usually not beneficially influenced by intramuscular injections of crude liver extract the reticulocyte response attaining a level of only 7 per cent, this in contrast to its more successful therapeutic result in cases of nontropical sprue.

The occurrence of macrocytic anemia in association with intestinal lesions, particularly intestinal strictures and anastomoses was amply demonstrated in the literature by Faber Hale White by Meulengracht and by Hurst. Butt and Watkins mention 7 cases of diffuse ileitis with macrocytic anemia in 6 of which a fecal fistula co-existed. With proper reconstructive surgical intervention the macrocytic type of anemia could be returned to a normal blood picture. They rightly ascribe the hyperchromic anemia to the abnormal interference with absorption and improper utilization of the effective hemopoietic fraction of the diet. Plum and Warburg in a publication entitled Hematological Changes Especially Megalocytic Anemia in Regional Ileitis describe 4 cases under their personal observation in which the blood picture was typical of that of a hyperchromic anemia. The hemoglobin values ranged as low as 20 per cent red blood cell count as low as 1,410,000 cells



levels, fasting serum carotene almost normal. The patient improved rapidly under high protein diet and amino acid concentrates. The intravenous administration of amino acids did not improve the nitrogen balance or create nitrogen retention in the metabolism studies.

Loss of fat soluble vitamins was especially emphasized in Albright and Stewart's report. The case, a woman aged 35 years had undergone four successive operations for ileitis and recurrent ileitis, including two extensive intestinal resections. Prolonged severe abdominal cramps and diarrhea resulted in tetany and a state of marked vitamin deficiency. The prothrombin time was found to be only 13 per cent of normal, indicating a marked associated vitamin K deficiency. Vitamin C was within normal range. Fat soluble vitamin A was somewhat deficient 82 units per 100 cc of serum (normal 10-20 units); carotenoids somewhat diminished, and associated with a xeroderma of the breast and abdomen. Fat soluble steroids, particularly the quantity of 17 keto steroids in the urine were practically absent. The tetany responded rapidly to large doses of vitamin D, 500,000 USP units daily. The results of treatment in the form of liberal dosage with all of the fat soluble vitamin preparations was clinically dramatic.

An unusually exaggerated degree of hypoproteinemia was noted in a diffuse case of ileo jejunitis by Craven. At a previous operation an anastomosis between the upper jejunum and the transverse colon had been performed because of a diffuse inflammatory process involving all of the ileum and the lower jejunum. Extreme malnutrition and evidences of hypoproteinemia were soon evident. The total serum proteins fell to 3.1 grams per 100 cc, the albumin fraction being 1.1 Gm and the globulin 2 Gm (albumin globulin ratio 0.55). Most observers conclude that the critical level of serum protein is 5.0 Gm per 100 cc, edema will invariably result when the blood level falls below that point.

*Blood changes.* The white cell count varies only a

wall occurred only once in this series. Internal fistula, jejunal to ileum, was once observed radiographically.

In cases of localized intestinal obstruction the whole abdomen may be distended. Visible peristalsis was not seen since the degree of obstruction is not complete. The edge of the spleen was palpable three times in this series. Club-



Fig. 54.—Clubbing of the fingers in a Case of Diffuse Ileo Jejunitis. Massive involvement of the whole jejunum and upper ileum. Palpable spleen.

bing of the fingers was observed twelve times in this series of 36 cases. Clubbing of the fingers is a significant observation and indicates the degree of constitutional alteration invoked by so diffuse an intestinal process (fig. 54).

color index 112, 119, 115, 105 at the point of greatest severity. In one case the jejunum seemed definitely to be involved. Burke and Hummel similarly quote a case of ileitis with resection and a secondary short circuiting operation and a consequent hyperchromic anemia.

Spigel, a disease without noticeable organic changes in the intestinal mucosa, but with completely deficient intestinal absorption is often characterized by a hyperchromic or microcytic type of anemia. Ileo jejunitis, by contrast, is a disease with extensive organic destruction of wide areas of intestinal mucosa, yet absorption is better maintained in spite of severe and extensive ulcerations and a hyperchromic type of anemia is rare.

The erythrocytic sedimentation rate may be somewhat accelerated, as seen in 5 patients. The total blood protein of the plasma is appreciably diminished in the more severe cases (5.8 Gm.), reaching a minimum of 4.0, 4.4, and 4.5 grams per 100 cc. in 3 cases and as low as 3.8 grams in a very severe case of ileo jejunitis. The albumin globulin ratio is disturbed, approaching a ratio of one but rarely completely reversed.

The urea, phosphorous, blood chlorides and blood sugar are but slightly if at all lowered. The blood calcium may be materially reduced, 2 of the patients in our present series having shown frank symptoms of tetany.

Gastric analysis showed normal titrable acidity in all cases except one of achylia gastrica.

*The physical examination of a case of ileo jejunitis may not be noteworthy. Some of the cases show well maintained nutrition, others may give evidence of pallor and extreme malnutrition in the later stages of a deteriorating status. A mass is occasionally present (6 cases), usually in the left upper abdomen and corresponds to the inflamed mesentery of the jejunum with the associated granulomatous loops of bowel. Such a mass may often be adherent to the anterior abdominal wall forming a localized abscess requiring incision and drainage. External fistula to the abdominal*

tory operation. The disorientation of the mucosal pattern of the jejunum and upper ileum is usually capable of recognition, segmentation and delayed puddling may occur similar to that seen in sprue, but in jejunitis there are frequently distended loops of bowel proximal to a localized partly



Fig 5b—Diffuse Ileo Jejunitis in a Boy 17 Years of Age. Exploratory operations (two). Impossibility of surgical relief with rapid downward course.

strictured area, or even evident skip areas separating involved segments of bowel (fig 5a).

The roentgen findings in ileo jejunitis are well described by Sussman and Wachtel. They were unable in the acute phase of the disease to differentiate jejunitis from sprue.

Rectal complications do not occur with the same frequency as in regional ileitis. In this series perirectal fistulas were present four times, a perirectal abscess twice, rectal fistula once.

#### ROENTGENOGRAPHIC RECOGNITION OF ILEO JEJUNITIS

The barium meal is utilized with reduced amount of



FIG. 55—Ileo Jejunitis Involving Upper Ileum and Lower Jejunum. Resection in continuity at the Mayo Clinic with brilliant result. Complete restoration of health.

barium sulphate and with frequent fluoroscopic and film controls is the method of choice. The degree and extent of involvement of the jejunum and ileum are definable with great accuracy as checked against the findings at exploratory

the barium rapid passage and poor mucosal pattern. In the chronic phase of the disease, greater rigidity and irregularity were noted, the barium seemed to make a cast. The



Fig. 58—Same patient as in Figure 57. Radiograph taken two and a half years later during which time subsidence of symptoms and a gain of 50 pounds in weight were noted. Observe by comparison the increased lumen of the jejunum and ileum but persistent irregularity in contour probably evidence of scarring. Note also absence of stenosis or stricturing.

rigid loops had a pipe stem appearance with occasional arcs of dilatation, multiple stenotic areas, and occasionally evident polyposis was suggested (figs. 56-58).

the same patchy segmentation was present in the loops of the bowel, the valvulae conniventes seemed further separated, the outlines of the jejunum were hazy. The loops of intestine showed persistently increased activity with



Fig. 57—Diffuse Ileo Jejunitis in a Woman 50 Years of Age Observed over a Course of Several Years. Radiograph taken in 1935 at the height of the active process showing involvement of the whole ileum and lower jejunum. See figure 58.

lack of flexibility. The terminal ileum was also involved in half of their cases.

In the subacute phase, the jejunal loops showed more pronounced alteration in contour, uneven distribution of

the barium, rapid passage and poor mucosal pattern. In the chronic phase of the disease, greater rigidity and irregularity were noted, the barium seemed to make a cast. The



Fig 58—Same Patient as in Figure 57. Radiograph taken two and a half years later during which time subsidence of symptoms and a gain of 50 pounds in weight were noted. Observe by comparison the increased lumen of the jejunum and ileum but persistent irregularity in contour probably evidence of scarring. Note also absence of stenosis or stricturing.

rigid loops had a pipe stem appearance with occasional areas of dilatation, multiple stenotic areas, and occasionally evident polyposis was suggested (figs. 56-58).



## DIFFERENTIAL DIAGNOSIS

The main point of interest lies in differentiation, high ileo jejunitis from sprue and from the secondary vitamin deficiency manifestations. Sprue is characterized by typical frothy and fatty stools, by a severe secondary anemia or by one of hyperchromic type with a high color index. Fever is absent, the stools do not contain blood, gross



Fig 59—Sprue. Characteristic Segmentation of Barium in Loops of Small Intestine. Contrast ileo jejunitis in figure 60

hemorrhage either as melena or as hematemesis is missing. The sprue case reacts well and promptly to deep injections of crude liver extract and vitamin B complex with the evident reversal of the mucous membrane disturbance of pattern and an attempt at least partially to resume a normal mucosal mosaic. The lower or terminal ileum is usually free of disease in sprue, but may often be involved in jejunitis. The intermittent areas of constriction and large

dilatation which is present as obstructive phenomena in jejunitis are absent in sprue (figs 59 and 60)

In jejunitis, marked general clinical improvement may be noted as a result of general hygienic measures and the administration of liver extract and vitamins, but the diffuse string signs throughout the jejunum and ileum are irre-



Fig 60—Linear Distribution of Barium in Ileo-Jejunitis in Contrast to Sprue in Figure 59

versible and remain permanent even though general good health and nutrition are restored. In a case of diffuse ileo-jejunitis observed over a course of 10 years, a complete subsidence of symptoms occurred with a regain of 50 pounds of weight. The repeated x-ray examinations revealed year after year the same continuous string signs occupying jejunum and ileum as seen at the time of the first observation. Evidently permanent scarring and cicatrization of

the intestinal wall had occurred and persisted even though clinically the patient had been restored to good health.

The absorptive capacity of the intestinal tract differs in sprue from that of ileo jejunitis. In sprue, the oral glucose tolerance test gives flat curves consistent with poor or deficient intestinal absorption. The same applies to tests for the intestinal absorption of vitamin A, as shown in experiments by Adlersberg. In ileo jejunitis, the oral glucose tolerance test is more likely to approach the normal rise and fall as is also the absorption of vitamin A from the intestinal tract. According to Adlersberg and Sabotka, ileo jejunitis and sprue can be reliably differentiated by postdigestive tests for the absorption of fats and vitamin A. In the active stages of sprue there is complete inability of the intestine to absorb fat and fat soluble vitamin A. During remissions, after adequate treatment, one observes fairly satisfactory absorption of both. They studied 5 cases of diffuse ileo jejunitis. The fasting lipids of the blood serum averaged 377 m<sub>g</sub> per cent in ileo jejunitis four hours after a fatty meal the blood serum lipids averaged 623 m<sub>g</sub> per cent a rise of 65 per cent, in contrast to sprue in which a rise of only 3 per cent was observed. After feeding 90,000 to 180,000 I U. of vitamin A, the level in the blood serum rose in ileo jejunitis from 63 I U. per cent to 94 I U. per cent, in contrast to active sprue where an actual loss was demonstrated.

The vitamin deficiency states with disturbances of fat metabolism such as accompany severe ulcerative colitis, give roentgenographic changes similar to those of sprue and often to those of mild diffuse jejunitis. They have been described by Snell and Comp, by Mackie and Pound, and by Adlersberg and Weintraub. Mild puddling, delay in jejunal loops, irregular outline of the upper ileum and lower jejunum, such as occur in sprue and in jejunitis may be present, there are no manifestations of delayed motility and no obstructive or polypoid changes in the roentgenograms.

Hodgkin's disease of the small bowel and diffuse multicentric sarcomatosis of the mesenteric lymph nodes with intestinal involvement are rare diseases, which simulate jejunitis. Both of these diseases are extremely malignant, the course is rapid and downhill and perforation and hemorrhage not uncommon.

Continuous fever of intestinal origin arising in ileo jejunitis or ileitis must be recognized in the differential diagnosis where more common diseases of febrile nature receive first presumptive consideration. Diseases characterized by continuous fever with low leukocyte counts include such syndromes as brucellosis, rheumatic fever, periuteritis nodosa, lupus erythematosus disseminatus. The advent of diarrhea so often overlooked or ignored indicates the possibility if not the likelihood of ileo jejunitis as the true cause of the febrile state [69].

## 14 Ileo-Jejunitis—Treatment

### COURSE AND PROGNOSIS

THE OUTLOOK in a case of ileo jejunitis depends to a large extent upon the degree and type of involvement of the small bowel. Where diffuse ileo jejunitis exists the prognosis differs materially from those cases in which localized or segmental involvement is exclusively present. In the localized form surgical resection or a short circuiting operation becomes a profitable procedure. Progress of the disease from one segment of bowel to a higher or a lower level of intestine has not been observed in this series, on the other hand recession of the lesion with healing, though with persistent x-ray deformity of the bowel, is possible and has been observed.

In the former series of 17 cases previously published (Yunich and Ciohn) 4 of 9 cases of diffuse ileo jejunitis had done well without recourse to surgery, gained strength and weight. Five conservatively treated cases gained some weight diarrhea diminished, anemia was overcome (fig. 61) the end result was less satisfactory.

In this series of 38 cases the end results were as shown in table 18 (follow up two to twelve years)

TABLE 18—Ileo-Jejunitis Follow-up Present State

	Cases
Clinically well	11
Clinically fair	6
Poor with persistent symptom	6
Died	6
No follow up	6

It will be noted that 13 of the 38 cases were clinically so improved and free of symptoms as to allow of the clinical impression of being restored to health. In most of these cases, clinical improvement was not paralleled by changes in the roentgenograms. However, a recent experience was




Fig. 61—Distribution of Anatomical Lesions in an Earlier Series of Cases. Note the continuous forms of ileo jejunitis, the extension forms, the skip lesions, and the purely localized forms of jejunitis.

most illuminating. Five years ago a patient was noted to have had diffuse involvement of the jejunum and upper ileum with active symptoms. On returning this year for follow up observation marked clinical improvement was noted with cessation of all active symptoms. On re-examination at this time the jejunal pattern was seen to be restored to normal, the upper ileum was contracted and disarranged

thus indicating the greater susceptibility of the upper jejunum to self healing. Such instances of apparent cure of jejunitis, confirmed by exploratory operation or by roentgenography, have previously been independently noted by Pessagno, by Wilensky, by Sussman and Wachtel, and by Kross in individual cases.

On the other hand, the severity of the disease is indicated in the fact that in this series 6 cases were only fair, 7 were poor and, above all, 6 had fatal terminations. These deaths occurred as a result of hasty surgical intervention (2 cases) or as a result of perforation, (1 case), of sepsis (*Friedlander bacillus*) (once), of inanition (1 case), of intercurrent disease 1 case. The unsatisfactory clinical outcome in 19 cases including 6 deaths is clear evidence of the severity and persistence of the disease process in progressively undermining nutrition and protein and mineral reserves.

The fact, however, that 13 cases did improve so as to resemble clinical cure, is impressive, and does indicate the fact that self healing in high jejunitis is possible. The regain of weight in these cases is impressive and the subsidence from fever and diarrhea and anemia most striking. In the series of 23 cases published by Sussman and Wachtel, the end results were quite similar insofar as of 17 cases observed through a prolonged follow up 5 cases became inactive, 3 died postoperatively of intestinal obstruction, 3 continued in an active phase with exacerbations, 6 continued with chronic low grade activity. One case was followed to spontaneous regression and to the restoration of good health and function. From this publication one would gain the impression that the prognosis in ileo jejunitis is most unfavorable.

#### MEDICAL TREATMENT

The conservative treatment of ileo jejunitis applies principally to the type in which diffuse involvement of the entire small bowel exists or to those in which the disease affects such large stretches of upper ileum and the whole

jejunum as to preclude any idea of surgical intervention. The conservative approach to therapy is aimed at establishing some degree of partial sterilization of the intestinal tract by the use of antibiotics and to maintain nutrition and electrolytic balance in the plasma and to combat anemia and vitamin deficiencies.

As antibiotics penicillin is useless streptomycin possibly of some avail. In the febrile cases of ileo jejunitis, streptomycin in doses of 2 or 3 grams daily intramuscularly administered would seem well worth the trial. The insoluble sulfonamides, sulfasuxidine and sulfathaladine in large dosage is regularly administered in this type of presumable intestinal infection. Whether such sulfonamides have curative value, or merely act as inhibiting agents upon secondary infection remains to be determined. Several patients have reported control of diarrhea and pain while taking the sulfonamides, with recurrence of the symptoms upon discontinuance of the oral sulfonamide therapy, in the main the specific therapeutic value of the sulfonamides by and large remains open to question.

Nutrition is maintained by a liberal soft nonirritating or bland diet not too strictly construed. High protein, high carbohydrate elements in the diet are stressed meats, fish, cheese, eggs, milk, pureed vegetables, and fruit juices, bread and butter being forced. (See page 116) The protein content of the diet may be reinforced by the use of the protein hydrolysates up to 4 to 6 ounces a day being urged upon the patient. The newer commercial preparations are no longer bad tasting are actually palatable, and tend to maintain and restore nitrogen balance. Occasional transfusions of whole blood or of plasma or the intravenous use of amino acid hydrolysates and in overcoming hypoproteinemia, raise the total proteins of the blood and restore when plentifully administered over a long period of time the relationship between the albumin and globulin fractions of the serum.



The intramuscular injection of crude liver extract plus vitamin B complex, 2 cc of each, every day or every other day, tends to overcome anemia, encourages blood formation and replaces the deficiency in vitamin absorption which results from the damaged intestinal tract. Even though



Fig. 62—Diffuse Ileo-Jejunitis in a 28 Year Old Man. Slowly progressive course, episodic flurries of fever, abdominal pain and diarrhea, marked clubbing of fingers, palpable ileon.

the beneficial effects of such injections are not as evident in ileo-jejunitis as in sprue, they should continue to be regularly administered.

The administration of folic acid by mouth is often to be recommended particularly in the types with hyperchromic forms of anemia.

Fresh air, sunshine and much rest particularly at the seashore or in the mountain air often acts very beneficially to effect general improvement in weight and strength.

### SURGICAL INTERVENTION

The indications for surgical interference are limited to those cases in which the lesion is localized to one or more



Fig. 63.—Diffuse Ileo Jejunitis in a Woman 28 Years of Age. Confirmed by Exploratory Laparotomy. Inoperable situation with present multiple abdominal wall fistulas requiring downhill course.

isolated segments of the small bowel such as the Group C or Group D of the original description (See page 160). Of the 36 cases in this group 15 were explored, in 7 only had an appendectomy been performed, 3 cases were subjected either to an intestinal resection or a short circuiting procedure. Surgical exploration is always warrantable and indicated where the true nature and extent of the disease is in some doubt. A case may seem to be segmental or localized in extent and at exploration may be discovered to be of the diffuse type. Or a case considered diffuse and

probably incapable of surgical help may be determined at exploratory operation to be segmental and capable of resection (figs 62-69)

Where no benefit is being derived by conservative medical treatment, an exploratory operation is often justified. The removal of the appendix is a superfluous manipulation.



Fig 64—Diffuse Ileo Jejunitis in an 18 Year Old Girl. Unsatisfactory clinical course progressive loss of nutrition hypoproteinemia

fortunately, in ileo jejunitis appendectomy rarely results in abdominal wall fistula.

In Group A, where the disease is in extension upward of ileitis so as to invade and include the jejunum no radical surgical procedure offers much hope of success.

In Group B, those cases in which the disease invades the whole of the jejunum and much or all of the ileum the same discouraging attitude toward reckless excision of the involved intestine holds equally true.

Massive resections of the small bowel with the sacrifice of most of its length has been occasionally encountered in

the literature, not for ileitis or jejunitis, but more often for strangulation for volvulus for superior mesenteric vein thrombosis or for intussusception. The average length



Fig. 63.—Diffuse Ileo Jejunitis in a Man 48 Years of Age. Note multiple areas of constriction chronic course. Episode of fever abdominal pain and obstructive phenomena. In the interval between attack preserved well being good nutrition and active working efficiency.

of the small intestine varies under normal conditions from 18 to 26 feet averaging 21 feet 6 inches.

Gray's Anatomy 20 feet 6 inches

Rost 18 to 26 feet averaging 19 feet

Treves 22½ feet in the male

23¼ feet in the female adult

Sunoff felt that half of the small intestine could be sacrificed without unduly endangering life and quoted 5 cases in which 18, 17 $\frac{1}{2}$ , 17, 16, and 13 $\frac{1}{4}$  feet of small bowel had been removed—all survived the operation, but died later




Fig. 66.—Diffuse Ile Jejunitis. Lipid Formation throughout Upper Ileum. Patient maintained good nutrition and weight, no evidence of anemia, normal blood figure. He leads a normal life, actively efficient.

Fate reported a case in which 12 $\frac{1}{2}$  feet of small bowel were removed for a traumatic stimulation of the mesenteric, only 12 inches of jejunum remaining, the patient was well four weeks later. In a case reported in 1940 (Todd, Dittenbriand and Montague) all but 3 feet of jeju

num had been surgically removed, resultant carbohydrate metabolism was normal, proteins not so well absorbed (67, 70, and 76 per cent in three periods) tetany was prevented by liberal administration of calcium and Vitamin D. In their study of the metabolism in their case, 20 per cent of the protein intake and 45 per cent of the fat intake




FIG. 6—Diffuse Ileo Jejunitis Involving Upper Ileum and Lower Jejunum

were lost in the feces—a total loss of 20 per cent of the caloric value of the ingested food. The fecal fat showed much free fatty acids indicating good splitting but poor absorption.

Coleman and Bennett report massive small intestinal resection with survival the resection involving 6 feet 7 inches in one case 14 feet 8 inches of small bowel in another

case Prioleau also supports the view that massive resection of the small intestine is possible with survival under fortunate circumstances. He reports 2 cases, one with a 40 per cent resection and one with a 53 per cent resection of the small bowel with survival, though the latter case died four months later. Flman and Reed report the



Fig. 68—Diffuse Ileo Jejunitis

resection of all but 3 feet of proximal jejunum and also the right half of the colon the patient gained 50 pounds of weight and experienced normal bowel function. The results of human resections indicate that approximately two thirds of the small bowel can be removed without serious risk to life (Berman et al). They report the removal of all but 18 inches of the jejunum and the right half of the colon for ergot poisoning and superior mesenteric vein thrombosis with complete recovery. The daily postoperative defecations

were limited to three formed stools per twenty four hours. At the 1948 meeting of the American Gastroenterological Association Althausen reported the resection for mesenteric thrombosis of 435 cm. of small intestine, including the




Fig. 69—Diffuse Ileo-Jejunitis Involving All of Ileum and Most of Jejunum

entire ileum and all but 15 cm. of the jejunum. The patient survived the operation tolerably well with retained nutrition as proven by extensive studies of metabolism and of food absorption.

While such massive resections of small bowel are thus



shown to be possible, with survival, the attempt to apply that principle to diffuse ileo jejunitis would likely be associated with an extremely high mortality because of the previous state of malnutrition and anemia, and is scarcely to be encouraged.

In Group C, those cases with high and isolated segmental involvement of the jejunum, localized resection of the lesion is not only possible but is highly successful. Such was the outcome in our 3 cases in this present series. Johnson resected 3 feet of upper jejunum beginning at the fossa of Treitz for localized jejunitis. Brewster resected 2 feet of jejunum for a lesion beginning  $2\frac{1}{2}$  feet caudad of the fossa of Treitz. Both cases made complete recoveries. Spellberg and Gray performed a gastroenterostomy for a tumefactive segmental jejunitis, because of poor result denoted by delay in the proximal loop, a secondary resection was performed with a successful outcome. Gendel and Berver had an identical experience. Lyons and Garlock resected 10 inches of upper jejunum for localized jejunitis, whose upper limit was the fossa of Treitz, with good result. The case of Koenig was equally successful after a resection of 3 feet of jejunum.

In high jejunitis multiple segments of involvement are not uncommon. It often becomes necessary to treat or resect each individual focus of disease as an independent lesion, in order to achieve a successful outcome. Dixon cites a striking case of recurring obstruction from multiple non neoplastic tumefaction of the jejunum. An interesting insight into the life history of localized jejunitis may be gained from the study of a case recently published by Husebye. In 1928 a 49 year old man was operated upon for an acute phlegmonous jejunitis; an entero anastomosis was performed at the ileo jejunal junction. The operation was followed by a long period of apparent good health. In 1945, or seventeen years later, the reappearance of heartburn and abdominal cramps was followed by the clinical picture of an 'acute abdomen'. The radiographic

examination showed the upper jejunum to be irregular, narrowed, tortuous the upper jejunum and the duodenum were distinctly dilated and retained bromum. At operation the roentgenographic findings were confirmed, the jejunum was diffusely scarred the lumen contracted and the proximal bowel and duodenum and stomach abnormally dilated. A gastro jejunostomy and a jejuno jejunostomy were performed the patient succumbed to the high intestinal obstruction and the extensive surgical procedures. Apparently in this case in spite of the entero anastomosis the healing of the short circuited segmental jejunitis had resulted in scar tissue contraction and in high intestinal obstruction.

In Group D where a high jejunitis is associated with an independent and similar lesion in the terminal ileum the operation of choice is localized resection of the upper jejunal lesion and a short circuiting of the lower ileal lesion (ileo transverse colostomy with transection of the proximal ileum above the diseased area). This present series includes three additional instances of combined lesions successfully operated by local resection of the jejunal affect and short circuiting of the terminal ileal segment all of them successfully accomplished.

In summary it will be seen that surgical intervention in ileo jejunitis is advocated both as an exploratory procedure and as a means of attack in those cases of high jejunitis with or without distal ileal involvement. In the localized or segmental forms of jejunitis, surgical intervention results frequently in restoration of health and intestinal function.

## 15 Ileo-Colitis—Combined Forms

---

**BOTH ILEITIS** and so called nonspecific or indeterminate colitis constitute diseases if not identical in nature, at least of the same family. The etiology of neither disease is known both occur in young people, with the same even sex determination, both run the same general life span of extension and are signalized by such common complications as rectal suppurative phenomena, erythema nodosum, fever, secondary anemia and nutritive disturbances. Both occur in pure form in the overwhelming proportion of cases but certain mixed forms must be recognized. Thus severe universal ulcerative colitis will involve in about 24 per cent of the cases, the most terminal portion of the ileum. This, however, does not constitute regional or terminal ileitis but is simply the backwash of the colitis through the ileo cecal valve into the reservoir of terminal ileum. Such ileitis does not extend upward by skip lesions and does not form internal nor external fistulas. On the other hand, true regional enteritis rarely crosses the ileo cecal valve but does persist proximal to the valve of Bauhin. When the ileum is anastomosed to the colon as occurs in all short circuiting operations and in resections for ileitis, recurrences take place in the new terminal ileum but rarely in the apposed colon even though now the guardianship of the ileo cecal valve is no longer present. In some instances the recurrence of ileitis at the free and open anastomosis has been followed for many years (in one case up to seventeen years) and yet when the anastomosis was taken down the transverse colon was free of disease sug-

estimating the fact that the colon is relatively immune to the "virus" of the ileitis

Relatively, yes, but not always so, since combined forms of the disease do occur, in this series of 306 cases of ileitis




Fig. 70—Combined Ileitis and Colitis (22 Cases) (A) terminal ileum and proximal colon (10 cases) (B) terminal ileum and transverse colon (2 cases) (C) terminal ileum and whole colon (except rectum) (6 cases) (D) ileum jejunum and whole colon (except rectum) (2 cases) (E) terminal ileum and distal colon (2 cases)

are included 22 cases of combined ileitis and colitis. Of these 22 cases, 11 represent combinations of true terminal

## 15 Ileo-Colitis—Combined Forms

---

BOTH ILEITIS and so called nonspecific or indefinite colitis constitute diseases if not identical in nature, at least of the same family. The etiology of neither disease is known, both occur in young people, with the same even sex determination both run the same general life span of extension and are signalized by such common complications as rectal suppurative phenomena, erythema nodosum, fever, secondary anemia and nutritive disturbances. Both occur in pure form in the overwhelming proportion of cases but certain mixed forms must be recognized. Thus severe universal ulcerative colitis will involve in about 24 per cent of the cases, the most terminal portion of the ileum. This, however, does not constitute regional or terminal ileitis but is simply the backwash of the colitis through the ileo cecal valve into the reservoir of terminal ileum. Such ileitis does not extend upward by skip lesions and does not form internal nor external fistulas. On the other hand, true regional enteritis rarely crosses the ileo cecal valve but does persist proximal to the valve of Bauhin. When the ileum is anastomosed to the colon as occurs in all short circuiting operations and in resections for ileitis, recurrences take place in the new terminal ileum but rarely in the apposed colon even though now the guardianship of the ileo cecal valve is no longer present. In some instances, the recurrence of ileitis at the free and open anastomosis has been followed for many years (in one case up to seventeen years) and yet when the anastomosis was taken down the transverse colon was free of disease, sug-

These findings of combined lesions were confirmed in 1936 by Crohn and Rosenak who cited 60 cases of regional ileitis, in 9 of which simultaneous involvement of the colon was found to exist. Musick further described a case in



Fig. 72—Regional Ileitis and Colitis Involving Cecum, Ascending, and Proximal Transverse Colon in a 21 Year Old Man. At operation the extensive involvement of the colon was fully realized. Ileo sigmoidostomy with creation of temporary mucosal colostomy of proximal sigmoid.

which 18 inches of ileum, 3 inches of cecum and 3 inches of sigmoid were simultaneously involved. An ileo ascending colostomy, right side, a Mikulicz successful outcome. Perianal with involvement

ileitis and various forms and distributions of ulcerative colitis, the other 11 cases are mixed types of diffuse ileitis and ileo jejunitis and variously distributed patterns of ulcerative colitis (fig 70). The sequence in the life history of the combined diseases is not known since when recognized



Fig 71—Regional Ileitis at Onset. Subsequent extension to ascending and transverse colon. Surgical resection recurrent in upper ileum and jejunum. General health well preserved, good working efficiency, consistent gain in weight. Only occasional flare up of mild diarrhea.

both large and small bowel are already involved. Actually, Colp as early as 1934 first pointed out the existence of a combined form of ileitis and colitis. In 1935, Mixer reported a case of terminal ileitis with extension into the cecum. In the discussion of Mixer's cases, Binney also quoted a case in which an extension of the process into the cecum caused perforation and a localized abscess.

These findings of combined lesions were confirmed in 1936 by Crohn and Rosenak who cited 60 cases of regional ileitis, in 9 of which simultaneous involvement of the colon was found to exist. Musick further described a case in



Fig 72—Regional Ileitis and Colitis Involving Cecum As ending in Proximal Transverse Colon in a 21 Year Old Man. At operation the extensive involvement of the colon was fully realized. Ileo-sigmoidostomy with creation of temporary mucosal colostomy of proximal sigmoid.

which 18 inches of ileum, 3 inches of cecum and 3 inches of sigmoid were simultaneously involved. An ileo ascending colostomy was performed on the right side, a Mikulicz procedure on the left colon with successful outcome. Pesagno (1937) reported regional ileitis with involvement



ileitis and various forms and distributions of ulcerative colitis, the other 11 cases are mixed types of diffuse ileitis and ileo jejunitis and variously distributed patterns of ulcerative colitis (fig 70). The sequence in the life history of the combined diseases is not known since when recognized



Fig 71—Regional Ileitis at Onset. Subsequent extension to ascending and transverse colon. surgical resection. recurrence in upper ileum and jejunum. General health well preserved. good working efficiency. consistent gain in weight. Only occasional flare up of mild diarrhea.

both large and small bowel are already involved. Actually Colp as early as 1934 first pointed out the existence of a combined form of ileitis and colitis. In 1935, Mixer reported a case of terminal ileitis with extension into the cecum. In the discussion of Mixer's cases, Binney also quoted a case in which an extension of the process into the cecum caused perforation and a localized abscess.

The most common form is constituted of those cases in which the terminal ileum is associated with a disease process involving the proximal colon, cecum and ascending colon up to the hepatic flexure (10 cases) (fig 73). Of equal frequency are mixed forms of terminal ileitis and transverse or distal colon involvement (10 cases). Finally



Fig. 73.—Combined Ileitis Ceci and Ascending Colitis. Restoration to perfect health.

one notes 2 cases of great severity where the whole of the small intestine and the whole of the colon to the sigmoid are almost continuously involved. It is of interest to note that the rectum and pelvic colon are rarely invaded; the form of colitis under consideration is the segmental or regional form with proximal involvement. For it is known that in regional colitis the parent lesion regularly occurs in the proximal colon and extends distally, usually over the course of years (Crohn and Berg). The life cycle of the disease has been studied, the inflammatory process stops

of the cecum Pugh reported 9 cases of ileitis with evidence of extension of the pathologic process into the large bowel. In 6 the cecum, in 2 the sigmoid and in one the transverse colon were implicated. The pre-eminence of the combined form of the disease seems to rest in the terminal ileum. Pemberton and Brown pointed out that if an ileo colostomy be performed for a combined lesion, at subsequent exploration the colonic lesion will be found healed, only the ileitis constituting the residual focus of the disease. This is an astute observation which is amply confirmed in the life histories of several cases in this present series. Clark and Dixon found that in 23 out of 44 cases of regional enteritis observed, the large bowel, mainly cecum and ascending colon were involved with the terminal ileum. And yet, they distinctly say "The distal portion of the ileum appeared to be the original site of invasion of the disease in practically all cases" (figs 71 and 72).

### ETIOLOGY

The age and sex distribution of combined ileo colitis is similar to that of the two independent diseases. The age incidence ranged from 16 years to 56 years, the average or mean age being 28 years. The sexes were evenly divided, 11 males and 11 females.

The pathologic distribution over the anatomy of the two tracts is the most interesting feature but the various forms and combinations are so heterogeneous as to make systematic classification difficult. An attempt at classification may be made, however, as in table 19.

TABLE 19—*Distribution of Combined Ileocolitis*

	Cases
Terminal ileum and proximal colon	10
Terminal ileum and transverse colon	2
Terminal ileum and distal colon	2
Terminal ileum and whole or distal colon to sigmoid	6
Ileum and jejunum and whole colon	3
Total	23

that form of terminal ileitis which is characterized by mass formation and by multiple internal fistula formation. The spleen was palpable in 2 cases, clubbing of the fingers occurred once in this group.

Internal intestinal fistulas occurred in four instances, as demonstrated by x ray examination, one from ileum to




Fig. 74.—Fistula between Diseased Terminal Ileum and Cecum. Fistula discernible as a fine line in the radiograph.

the cecum, one to the transverse colon, two to the sigmoid (fig. 74). Again internal fistulas from terminal ileum to other loops of small intestine such as characterize true terminal ileitis have not been noted. In only one case was a fistula present to the anterior abdominal wall.

Rectal complications such as mark both ileitis and ulcerative colitis are again present in this combined form as would be expected. Perirectal suppurative fistulas and ischio-rectal abscesses occurred in 9 of the 22 cases, a recto-vaginal fistula in one instance. Intestinal obstruction at the terminal ileum occurred once only in this group.

or carries for long periods at the sigmoid flexure, extending into the rectum rarely or only late in the course of an intractable and particularly virulent type of invasion and progression

That even the rectum is not immune is evidenced by the fact that in 2 of the fatal cases that succumbed after multiple and unsuccessful resections, the terminal phase was marked by severe proctitis. Two other living cases have severe grades of proctitis, following unsuccessful surgical procedures

### CLINICAL FEATURES

Except for the greater severity of the disease, the combined forms of ileocolitis duplicate in general those of regional enteritis and of ulcerative colitis. The general features are those of fever, which was present at some time in nearly all of the cases, again low grade and intermittent, present during the active phases of the disease, absent during remissions. The temperature may reach as high as 104 F in exceptional cases, but usually varies between 101 and 102 F. Malnutrition is extreme, losses of up to 50 or 65 pounds occurring in the more severe types of cases. A mild leukocytosis (10,000 to 12,600) may be present with an abnormal increase of nonsegmental forms in the blood, indicating severe toxicity. Secondary anemia is marked with hemoglobin values usually ranging between 60 and 80 per cent Sahli. Exceptional hemoglobin readings of 30 and 55 per cent have been noted, the red blood cell count corresponding. Sedimentation rate in the blood is usually moderately increased.

Hypoproteinemia is unusual, for even in severe cases the total serum proteins average between 60 and 63 grams per 100 cc with only an occasional slight inversion of the albumin globulin ratio. Occasional hypocalcemia has been observed.

### THE PHYSICAL SIGNS

Rarely an abdominal mass is present, for the dispersion of the disease process in the ileum and colon differs from

tion was performed but within one and a half years a recurrence, severe in nature, initiated by a melaena was observed in the whole colon, including even the rectum. This is an excellent example of a localized process of combined ileitis and colitis in which the recurrence involved the whole colon without recurrence in the residual ileum. In a second case cited by Taylor the resection of a strictly localized ileitis involving only 30 cm. of the terminal ileum recurred in three months with involvement of the whole colon with fatal termination.

#### MEDICAL TREATMENT

The medical treatment of the combined forms is not dissimilar to the care of ileitis or colitis, namely bed rest, sunshine, bland nonroughage diet, vitamin replacement by oral and parenteral route with the liberal use by injection of crude liver extract and vitamin B complex. Parenteral use of plasma, protein hydrolysates, and multiple transfusions are highly to be recommended. Of this series, 3 cases were not operated upon, one is well, spontaneously cured, the other 2 are too extensively involved to allow for any surgical resection or short circuiting operation.

#### SURGICAL APPROACH

The surgical approach to the cure of the disease is extremely varied and complicated and differs in each case depending upon the individual factors and the degree and extent of involvement of the various segments of large and small bowel.

Four cases have had an ileo transverse colostomy with transection of the ileum performed, thus short circuiting a circumscribed lesion involving the ileum and proximal colon. One case is well, 2 have continued to do poorly because of persistence of the original disease, the fourth has had extensive recurrences in the remainder of the colon.

Resection of the original lesion in terminal ileum and proximal colon seems to have been the operation of choice.

## PROGNOSIS

The prognosis in combined ileo colitis is poor, as might be expected where two such severe diseases are concurrently present. The end results to date are as shown in table 20.

TABLE 20—*End Results in Ileo Colitis*

	Cases
Well	7
Improved	2
Poor	9
Ceased	3
No follow up	1
	—
Total	22

Again it is interesting to note that 7 cases are well. Of these, only one case represents a spontaneous recovery under conservative management. This was an instance of terminal ileitis with concurrent involvement of the transverse colon. Under nonspecific protein therapy, using typhoid vaccine as an agent, clinical improvement was rapid, and eleven years later this man is apparently perfectly well and free of symptoms. The recent x ray study shows a perfect mucosal pattern in the colon throughout, the terminal ileum remains slightly distorted in outline.

The remaining 6 cases which are "well" represent successful surgical resections or short circuiting procedures.

Two cases are improved. 9 cases are classified as unimproved. Of these 9 instances of poor results, 5 represent surgical failures usually after multiple attempts, 4 have done badly because of such an extensive involvement of ileum, jejunum, and colon as to preclude any radical surgical approach.

Three further cases have ceased, 2 of them after multiple resections until only an ileo sigmoidostomy remained followed by recurrences in rectum and upper small bowel. The remaining case died of uremia, associated with multiple renal calculi and kidney parenchymatous damage.

Taylor describes a case with involvement of the terminal ileum, ascending colon up to the hepatic flexure. A resec-

vation. Cases successfully operated elsewhere would naturally have no necessity to seek further help, they are probably numerous throughout this country and elsewhere, where the combined types of ileitis and colitis are recognized and surgeons of ability are willing to devise some surgical procedure which fits the individual pattern and gives promise of fruition. Such a successful outcome is seen in the case of Musick, in which ileum, cecum and sigmoid were all involved. An ileo ascending colostomy was employed to short circuit the ileo-cecal area of disease, a Mikulicz procedure was utilized to resect the area in the sigmoid. The man was back at work within three months restored to health.

In a very recent publication the involvement of the ileum and jejunum secondary to extensive ulcerative colitis is stressed (Olsan and Sussman). This observation confirms the occasional case, rare fortunately, in which one sees involvement of the jejunum in a very serious type of ulcerative colitis. The widespread inflammatory invasion should be regarded as essentially colitis and not true ileitis or jejunitis.



in most instances. In this series, 9 cases had been submitted to such a procedure with greatly varying results. Three of the 9 cases so operated are well today, many years after the original procedure. Two have died as a result of extensive recurrences with subsequent multiple operations and eventually involvement of the rectum. The extension of the process to the rectum or to the upper jejunum always marks the terminal stage of the disease.

The remaining 4 cases that underwent resection as an initial procedure have fared badly because of extensive recurrences in colon mainly, one also in jejunum, but have not been subjected to further surgical procedures.

Four cases were submitted to ileo sigmoidostomy. One is well, one died of uremia some months after operation, 2 have done poorly, and are awaiting resection at some future date.

Two cases remain as instances of extreme surgical necessities. One has had a high ileostomy as an attempt at a life saving procedure. One had had elsewhere a jejuno sigmoidostomy in an attempt to short circuit all of the ileum and all of the colon down to the sigmoid. The latter is a particularly poor result with marked hypoproteinemia and malnutrition. All in all, the results are not good. Five well out of 22 cases is not very promising. However, it should be remembered that most of these cases were referred because of the disappointment over the results of initial surgical attack previously performed elsewhere and lacked success, although the procedures had been carried out by most competent surgeons. Of the cases in which the diagnosis and surgical indications were solely in my own hands from the initiation (7 cases) 4 are well, one has died, one is a poor surgical result and one is awaiting further resection after an ileo sigmoidostomy with good promise. Perhaps these 7 cases give a more true picture of the eventual prognosis. The remainder of the series hardly constitute a basis of prognostication since they had already suffered the handicap of being failures before coming under obser-

## References

- 1 Abercrombie J Pathological and Practical Researches on Diseases of Stomach the Intestinal Tract and Other Viscera of the Abdomen Edinburgh Waugh & Innes 1825 Pp 235
- 2 Adlersberg D and Sobotka H Fat and vitamin A absorption in sprue and jejuno ileitis Gastroenterology 1 307 1943
- 3 ——— and Weingarten M Enteropathy in deficiency states New York State J Med 41 14-19 1941
- 4 Albright F and Stewart J D Hypovitaminosis of all fat-soluble vitamins due to steatorrhea New England J Med 3 239-241 1940
- 5 Althausen T L Uveyama K and Simpson R G Digestion and absorption after massive resection of the small intestine Presented at the annual meeting Am Gastroenterological Assoc April 30 1948 at Atlantic City N J
- 6 Anshutz C Ueber unspezifische entzündliche Geschwulste des Dickdarms Deutsch Zsch Chir 143 37 1934
- 7 Arnheim L Regional ileitis with perforating abscess and peritonitis J Mt Sinai Hosp 61 1935
- 8 Asencio Camacho F Regional enteritis in Puerto Rico Gastroenterology 6 483 1946
- 9 Ashley L B Meyers S G and Reynolds L Regional entero-colitis J Michigan State M Soc 41 109 1943
- 10 Babson W W Terminal ileitis with obstruction and abscess complicating pregnancy New England J Med 35 544-547 1946
- 11 Barbour R F and Stokes A B Chronic cicatrizing enteritis A phase of benign non specific granuloma of the small intestine Lancet 1 299-303 1936
- 12 Barger J A Wesson H R and Jackman R G Studies in the ileo-cecal valve (ileo-cecus) Surg Gynec & Obst 1 33 1940
- 13 Barker W H and Hummel L E Macrocytic anemia in association with intestinal strictures and anastomoses Review of literature and report of two new cases Bull Johns Hopkins Hosp 64 215-256 1939
- 14 Bell H G Chronic cicatrizing enteritis California and West Med 41 39 1934
- 15 Berman J K Brown H M Foster R T and Grisell T L Massive resection of the intestine J A M A 135 918-919 1947



- 37 Brown D C and McHardy C Primary lesion of jejunum *J A M A* 115 225 1940
- 38 Buckston J Primary ulcer of the jejunum *Radiology* 33 299-304 1939
- 39 Butt H R and Watkins C H Occurrence of macrocytic anemia in association with lesions of the bowel *Ann Int Med* 10 292 1936
- 40 Camel M R Ileocecal tuberculosis *Radiology* 41 344-351 1943
- 41 Cation D Nutritional disturbances in regional enteritis *Surgery* 6 708-716 1939
- 42 Cave H W Inflammatory lesions of the small intestine *Surgical aspects* *J A M A* 127 456 1943
- 43 Clark F and Wright A Acute phlegmonous enteritis *Arch Surg* 34 997 1937
- 44 Clark R I Jr and Dixon C F Regional enteritis *Surgery* 2 27-304 1939
- 45 Clute H M Regional ileitis A report of two cases *S Clin North America* 13 561-567 1933
- 46 Coffen T H Non specific granuloma causing intestinal obstruction *J A M A* 8 1303 1925
- 47 Coleman F I and Bennett D A Massive intestinal resection *Am J Surg* 9 429-438 1943
- 48 ——— and ——— Fecal fistulas following acute appendicitis *Ann Surg* 84 837 1926
- 49 Colp R A case of non specific granuloma of the terminal ileum and the cecum *S Clin North America* 14 443-449 1934
- 50 ——— Non specific granulomatosis of the intestine *Ann Surg* 107 74-81 1938
- 51 ——— Carlock J H and Ginzburg H Ileocectomy with exclusion for non specific ileitis *Am J Dig Dis* 9 64-68 1942
- 52 ——— and Ginzburg L Ileocectomy with exclusion in the treatment of regional ileitis *New York State J Med* 41 952-990 1941
- 53 Combe C and Saunders W A singular case of stricture and thickening of the ileum *Med Tr Roy Coll Phys Lond* 4 16-21 1813
- 54 Connell F G Regional ileitis *Am J Dig Dis & Nutrition* 3 438 1936
- 55 Corr I and Boeck W C Chronic ulcerative enteritis *Am J Dig Dis & Nutrition* 1 161 163 1934-5
- 56 Coutts W E and Opoz L Lymphogranulomatosis of mouth small and large bowel rectum and anus *Re Chil Hyg & Med* 2 83 1939
- 57 Crane A W A roentgenologic sign of mucous colitis *Am J Roentg and Rad Rev* 17 416 1917
- 58 Crohn B B The broadening conception of regional ileitis *A J Dig Dis & Nutrition* 1 91-93 1934-5
- 59 ——— Regional ileitis *Surg Gynec & Obst* 68 314-321 1939
- 60 ——— Editorial Acute appendicitis and diarrhea *Gastroenterology* 4 511 1945

- 16 Bernstein, M Konzelmann F W and Sudlick D M Boeck's sarcoid  
Report of a case with visceral involvement Arch Int Med 44  
721 1929
- 17 Binney, H Discussion of a paper by C G Mixer Regional ileitis  
Ann Surg 102 689 1935
- 18 Bisgard, J D and Henske J A Regional ileitis J A M A 103  
550 1937
- 19 Bissell A D Localized chronic ulcerative ileitis Ann Surg 99 957-  
966, 1934
- 20 Blackburn G Hadfield G and Hunt A H Regional ileitis St  
Barth Hosp Rep 2 181-224 1939
- 21 Blumenthal J S and Berman R Terminal ileitis with extension into  
cecum following non perforating abdominal trauma Minnesota  
Med 22 406 1939
- 22 Bockus H L Gastroenterology vol 2 Philadelphia W B Saunders  
Co, 1944
- 23 ——— Present status of chronic regional or catrizing enteritis  
J A M A 127 449-456 1945
- 24 ——— Johnson T A and Lee W E An appraisal of the results  
of surgically treated regional ileitis Liley Birthday Volume  
Springfield Ill C C Thomas 1940 1 p 53-83
- 25 ——— and Lee W E Regional (terminal) ileitis Ann Surg  
102 412-421 1935
- 26 Bonorino-Udaondo C Ilitis regionales Prensa Med Argent 27  
1141-1151 1940
- 27 ——— and Centono A M El pronostico en las enteritis segmen-  
tarias Prensa Med Argent 30 774-775 1943
- 28 ——— Maissa P A and DiLotto V El diagnostico radiologico de  
los ileitis egmentarias Prensa Med Argent 27 2111-2107 1940
- 29 Bowen W H and Day T D Post mortem in a case of regional  
ileitis nine years after symptomatic recovery following ileo  
colotomy Guy Hosp Reports 53 70-76 1939
- 30 Brewster H V Personal communication Gastroenterology 1 353  
1943
- 31 Braun H Ueber Entzündliche Erkrankungen des Netzes Arch Klin  
Chir 65 375 1901
- 32 Brown P W Barger J A and Weber N M Chronic inflammatory  
lesions of the small intestine (regional enteritis) Am J Dig  
Dis & Nutrition 1 426-431 1934-5
- 33 ——— and Donald C J Jr Irrguosis of regional enteritis Am  
J Dig Dis 9 87-91 1942
- 34 ——— and Pemberton J de J Solitary ulcer of the ileum and ulcer  
of Meckel's diverticulum Proc Staff Meet Mayo Clinic 11 259  
1936
- 35 ——— and Schieffley C H Chronic regional enteritis occurring in  
three siblings Am J Dig Dis 6 257 1939
- 36 Brown S Chronic non specific regional enteritis Am J Surg 51  
487-495, 1945

- 37 Browne D C and McHardy G Primary lesion of jejunum *J A M A* 110 2257 1940
- 38 Buckstein J Primary ulcer of the jejunum *Radiology* 33 299-304 1939
- 39 Butt H R and Watkins C H Occurrence of macrocytic anemia in association with lesion of the bowel *Ann Int Med* 10 222 1936
- 40 Camiel M R Ileo cecal tuberculosis *Radiology* 44 344-351 1945
- 41 Casten D Nutritional disturbances in regional enteritis *Surgery* 6 708-716 1939
- 42 Cave H W Inflammatory lesion of the small intestine *Surgical abstract J A M A* 127 456 1945
- 43 Clark E and Wright A Acute phlegmonous enteritis *Arch Surg* 31 997 1937
- 44 Clark R L Jr and Dixon C F Regional enteritis *Surgery* 5 277-304 1939
- 45 Clute H M Regional ileitis A report of two cases *S Clin North America* 13 561-567 1933
- 46 Coffen T H Non specific granuloma causing intestinal obstruction *J A M A* 81 103 1925
- 47 Coleman E F and Bennett D A Massive intestinal resection *Am J Surg* 59 429-438 1943
- 48 ——— and ——— Fecal fistulas following acute appendicitis *Ann Surg* 84 837 19 6
- 49 Colp R A case of non specific granuloma of the terminal ileum and the cecum *S Clin North America* 14 443-449 1934
- 50 ——— Non specific granulomata of the intestine *Ann Surg* 107 74 81 1938
- 51 ——— Carlock J H and Ginzburg H Ileo colostomy with exclusion for non specific ileitis *Am J Dig Di* 9 64-68 1942
- 52 ——— and Ginzburg I Ileo colostomy with exclusion in the treatment of regional ileitis *New York State J Med* 41 985-990 1941
- 53 Combe C and Saunders W A singular case of stricture and thickening of the ileum *Med Tr Roy Coll Phys Lond* 4 16-21 1813
- 54 Connell F G Regional ileitis *Am J Dig Di & Nutrition* 3 438 1936
- 55 Corr P and Boeck W C Chronic ulcerative enteritis *Am J Dig Dis & Nutrition* 1 161-163 1934-35
- 56 Crott W F and Opozol L Lymphogranulomatosis of mouth small and large bowel rectum and anus *Rev Chil Hyg y Med* 8 195 1959
- 57 Crane A W A roentgenologic sign of mucous colitis *Am J Roentg and Rad Ev* 1 416 1917
- 58 Crohn B B The broadening conception of regional ileitis *A J Dig Dis & Nutrition* 1 97-111 1934-35
- 59 ——— Regional ileitis *Surg Gynec & Obst* 68 314-311 1939
- 60 ——— Editorial Acute appendicitis and diarrhea. *Gastroenterology* 4 511 1945

- 16 Bernstein, M Konzelmann F W and Sidlick D M Boeck's sarcoid  
Report of a case with visceral involvement Arch Int Med 41  
721 1929
- 17 Binney, H Discussion of a paper by C G Mixter Regional ileitis  
Ann Surg 102 689 1935
- 18 Bisgard J D and Henske J A Regional ileitis J A M A 109  
550, 1937
- 19 Bissell, A D Localized chronic ulcerative ileitis Ann Surg 99 957-  
966 1934
- 20 Blackburn, G Hadfield C and Hunt A H Regional ileitis St  
Barth Ho p Rep 72 181-224 1939
- 21 Blumenthal J S and Berman R Terminal ileitis with extension into  
cecum following non perforating abdominal trauma. Minnesota  
Med 22 406 1939
- 22 Boekus H L Gastroenterology vol 2 Philadelphia W B Saunders  
Co 1944
- 23 ——— Present status of chronic regional or catrizing enteritis  
J A M A 127 449-456, 1945
- 24 ———, Johnson T A and Lee W E An appraisal of the result  
of surgically treated regional ileitis Lahey Birthday Volume  
Springfield Ill C C Thomas 1940 Pp 53-83
- 25 ——— and Lee W E Regional (terminal) ileitis Ann Surg  
102 412-421 1935
- 26 Bonorino Udaondo C Ileitis regionales I rensa Med Argent 27  
1141-1151 1940
- 27 ——— and Centono A M El pronostico en las enteritis segmen-  
tarias I rensa Med Argent 30 774-775 1943
- 28 ——— Mai sa I A and D Alotto V El diagnostico radiologico de  
los ileitis egmentarias I rensa Med Argent 27 2011-2107, 1940
- 29 Bowen W H and Day T D Post mortem in a case of regional  
ileitis nine years after symptomatic recovery following ileo-  
colotomy Guv's Hosp Reports 89 70-76 1939
- 30 Brew ter H A Personal communication Gastroenterology 1353  
1943
- 31 Braun H Ueber Entzündliche Cecchwalte de Netzes Arch Klin  
Chir 63 378, 1901
- 32 Brown, P W, Bergen J A and Weber N M Chronic inflammatory  
lesions of the small intestine (regional enteritis) Am J Dig  
Dis & Nutrition 1 426-431 1934-35
- 33 ——— and Donald C J Jr Diagnosis of regional enteritis Am  
J Dig Dis 7 87-91 1942
- 34 ———, and Emberton J de J Solitary ulcer of the ileum and ulcer  
of Meckel's diverticulum Proc Staff Meet Mayo Clinic 11 259  
1936
- 35 ——— and Schneffly C H Chronic regional enteritis occurring in  
three sibling Am J Dig Dis 6 157 1939
- 36 Brown S Chronic non peptic regional enteritis Am J Surg 51  
487-492, 1945

- 31 Browne D C and McHarris G Primary lesion of jejunum *J A M A* 115 2257 1940
- 32 Buckstein J Primary ulcer of the jejunum *Radiology* 33 299-304 1939
- 33 Butt H R and Watkins C H Occurrence of macrocytic anemia in association with lesions of the bowel *Ann Int Med* 10 227 1936
- 34 Cannel M R Ileo-cecal tuberculosis *Radiology* 41 344-351 1943
- 35 Casten D Nutritional disturbances in regional enteritis *Surgery* 1 708-716 1939
- 36 Case H W Inflammatory lesions of the small intestine Surgical aspects *J A M A* 127 436 1945
- 37 Clark F and Wright A Acute phlegmonous enteritis *Arch Surg* 34 997 1937
- 38 Clark R L Jr and Dixon C F Regional enteritis *Surgery* 3 217-304 1939
- 39 Clute H M Regional ileitis A report of two cases *S Clin North America* 13 561-567 1933
- 40 Coffen T H Non specific granuloma causing intestinal obstruction *J A M A* 85 1303 1925
- 41 Cokman F I and Bennett D A Massive intestinal resection *Am J Surg* 9 429-438 1943
- 42 ——— and ——— Fecal fistulas following acute appendicitis *Ann Surg* 84 937 1926
- 43 Colp R A case of non specific granuloma of the terminal ileum and the cecum *S Clin North America* 14 443-449 1934
- 44 ——— Non specific granuloma of the intestine *Ann Surg* 10 74 81 1938
- 45 ——— Carluck J H and Ginzburg H Ileocolostomy with exclusion for non specific ileitis *Am J Dig Dis* 3 64-68 1947
- 46 ——— and Ginzburg L Ileocolostomy with exclusion in the treatment of regional ileitis *New York State J Med* 41 982-990 1941
- 47 Combe C and Saunders W A singular case of stricture and thickening of the ileum *Med Tr Roy Coll Phys Lond* 4 16-21 1813
- 48 Connell F G Regional ileitis *Am J Dig Dis & Nutrition* 3 438 1936
- 49 Corr I and Boeck W L Chronic ulcerative enteritis *Am J Dig Dis & Nutrition* 1 161-163 1934-35
- 50 Coutts W F and Ojczko I Lymphogranulomatosis of mouth small and large bowel rectum and anus *Rev Chil Hyg y Med* 85 1939
- 51 Crane A W A roentgenologic sign of mucous colitis *Am J Roentg and Rad Rev* 1 416 1917
- 52 Crhn B B The broadening conception of regional ileitis *A J Dig Dis & Nutrition* 1 91-99 1934-35
- 53 ——— Regional ileitis *Surg Gynec & Obst* 68 314-321 1939
- 54 ——— Editorial Acute appendicitis and bruchia *Gastroenterology* 4 511 1945



- 61 ——— Clinical use of sulfasuxidine *Gastroenterology* 1 140-146 1943
- 61a ——— The use of sulfonamides in ileitis *Gastroenterology* 4 11-13, 1945
- 62 ——— The relationship of trauma to diseases of the gastrointestinal tract *Gastroenterology* 8 735-742, 1947
- 63 ——— Trauma and Disease In Brahdly and Kahn Philadelphia, Lea & Febiger 1941 P 185
- 64 ——— Benign diseases of the small intestine *Gastroenterology* 2 385-394 1944
- 65 ——— and Berg A A Right sided (regional) colitis *J A M A* 110 32-38 1938
- 66 ——— Ginzburg L and Oppenheimer, G D Regional ileitis A pathologic and clinical entity *J A M A* 99 1323, 1932
- 67 ———, and Rosenak, B D A combined form of ileitis and colitis *J A M A* 106 1-7 1936
- 68 ——— and Yarnis, H The anatomical position of the ileum in health and disease *Radiology* 33 325-330 1939
- 69 ——— and ——— Continuous fever of unknown origin *Ann Int Med* 26 858 1947
- 70 ——— and Yumich A M Ileo jejunitis *Ann Surg* 113 371-380 1941
- 71 Cutler E C A neglected entity in abdominal pain and a common disease Cicatricial enteritis *New York State J Med* 39 328 1939
- 72 Cushway B C Chronic cicatrizing enteritis Regional ileitis (Crohn) *Illinois M J* 66 525 1934
- 73 Dalziel T K Chronic interstitial enteritis *Brit M J* 2 1068-1070, 1913
- 74 Dixon C F Regional enteritis *Ann Surg* 108 857-866 1938
- 75 Dowdle E Multiple primary non specific jejunal ulcers with chronic duodenal dilatation *Ann Surg* 116 348 1942
- 76 Ebrill D A case of regional ileitis in childhood *Brit J Surg*, 32 512-514 1945
- 77 Eggers C Discussion of Gordon *Ann Surg* 97 132 1933
- 78 Elason E I and Johnson J Acute Regional Enteritis *Int Clin N S* 3 1 123-133 1940
- 79 Elman R and Read J A Nutritional recovery following removal of all but three feet of jejunum and half of the colon *J Missouri M A* 42 145-146 1945
- 80 Erb L H and Farmer A W Ileo colitis *Surg Gynec & Obst* 61 6-14 1935
- 81 Erdmann J F and Burt C V Non specific granuloma of the gastrointestinal tract *Surg Gynec & Obst* 57 71-80 1933
- 82 Erskine F B The pathologic relationship of mesenteric adenitis, ileitis and appendicitis *Am J Clin Path* 11 706-712 1941
- 83 Faber K Pernicious Anemia some follow up of terminal ileitis Hospital studende 601 1895 Fjerde Række Bind 111
- 83a ——— Perniciöse anæmie bei dunnarmstrickturen *Berl Klin Wchnschr* 34 643-646 1897

- 84 Fallis L S Massive intestinal hemorrhage in regional ileitis *A J Surg* 53 512 1941
- 85 Felger L and Schenk H L Recurrence in segmental enteritis following radical resection *A J Surg* 49 307-315 1940
- 86 Felten J Non specific ulcerative colitis terminal (distal) ileitis and bacillary dysentery Their common pathogenesis *New York State J Med* 35 576-578 1935
- 87 ——— The relationship of bacillary dysentery to distal ileitis chronic ulcerative colitis and non specific intestinal granuloma *Ann Int Med* 10 645-669 1936
- 88 Fenster L Ileitis ulcerosa *Brun's Beitr* 161 462 1936
- 89 Fischer A Appendicitis und Granulomata *Ztschr f Chir* 55 143 1931
- 90 Fischer A W and Lurmann Ueber eine tumorbildende ulceros stenosierende und perforierende entzündung des unteren ileum *Arch fur Klin Chir* 177 638-650 1933
- 91 Forbes R D and Duncan J Some observations on regional ileitis and allied conditions *West J Surg Obst & Gynec* 45 362-367 1937
- 92 Frazer I and Haggart H Crohn's disease *J Royal Army Med Corps* 81 9-13 1945
- 93 Friedl Meyer M Kasuistischer Beitrag zur tumorbildenden ulcerösen stenosierenden Entzündung des unteren ileum (Ileitis terminalis) *Schweiz Med Wchnschr* 66 508-512 1936
- 94 Galambo A and Mittleman W Typical and atypical terminal ileitis *Am J Dis Dis & Nutr* 2 442-447 1935-36
- 95 Garlock J H Present status of regional ileitis *Am J Surg* 72 875 1946
- 96 ——— and Crohn B B An appraisal of the results of surgery in the treatment of regional ileitis *J A M A* 126 205-208 1947
- 97 Gendel S and Berber M G An unusual case of Meckel's diverticulum *Ann Surg* 121 981-983 1945
- 98 Ginzburg L Persistent abdominal fecal fistulas due to regional ileitis *Surgery* 5 515-518 1940
- 99 ——— Colp R and Sussman M Ileocolostomy with exclusion *Ann Surg* 110 648 655 1939
- 100 ——— and Garlock J H Regional ileitis *Ann Surg* 116 906-91 1949
- 101 ——— and Oppenheimer C D Non specific granulomata of the intestines (inflammatory tumors and strictures of the bowel) *Ann Surg* 98 1046 1933
- 102 Gishertz H Ist die Palikale Operation der unspezifischen umschriebenen Entzündung des Ileumendes notwendig? *Zentr Chir* 65 7687 1936
- 103 Golden A Primary systemic amyloidosis of the alimentary tract *Arch Int Med* 75 413-416 1945
- 104 Goldfarb S J The roentgen diagnosis of lesions of the small intestine *New York State J Med* 31 500-505 1934

- 61 ——— Clinical use of sulfasuxidine *Gastroenterology* 1 140-146 1943
- 61a ——— The use of sulfonamides in ileitis *Gastroenterology* 4 11-13, 1945
- 62 ——— The relationship of trauma to diseases of the gastrointestinal tract *Gastroenterology* 8 730-742 1947
- 63 ——— Trauma and Disease In Braddy and Kahn Philadelphia Lea & Febiger 1941 P 180
- 64 ——— Benign diseases of the small intestine *Gastroenterology* 2 380-394 1944
- 65 ——— and Berg, A A Right sided (regional) colitis *J A M A* 110 32-38 1938
- 66 ——— Ginzburg L and Oppenheimer G D Regional ileitis A pathologic and clinical entity *J A M A* 99 1323 1932
- 67 ——— and Rosenik B D A combined form of ileitis and colitis *J A M A* 106 1-7 1936
- 68 ——— and Yarnis H The anatomical position of the ileum in health and disease *Radiology* 33 320-330 1939
- 69 ——— and ——— Continuous fever of unknown origin *Ann Int Med* 26 858, 1947
- 70 ——— and Yunch A M Ileo jejunitis *Ann Surg* 113 371-380, 1941
- 71 Cutler F C A neglected entity in abdominal pain and a common disease Cicatricial enteritis *New York State J Med* 39 378, 1939
- 72 Cu hway B C Chronic cicatrizing enteritis Regional ileitis (Crohn) *Illinois M J* 66 520 1934
- 73 Dalziel T K Chronic interstitial enteritis *Brit M J* 2 1068-1070 1913
- 74 Dixon C F Regional enteritis *Ann Surg* 108 807-866 1938
- 75 Dowdle F Multiple primary non specific jejunal ulcers with chronic duodenal dilatation *Ann Surg* 116 348 1942
- 76 Ebrill D A case of regional ileitis in childhood *Brit J Surg* 32 512-514 1945
- 77 Eggers C Discussion of Gordon *Ann Surg* 97 132 1933
- 78 Eliason F L and Johnson J Acute Regional Enteritis *Int Clin N S* 3 1 123-133 1940
- 79 Elman R and Reid J A Nutritional recovery following removal of all but three feet of jejunum and half of the colon *J Missouri M A* 12 145-146 1940
- 80 Erb L H and Farmer A W Ileo colitis *Surg Gynec & Obst* 61 6-14 1930
- 81 Erdmann J F and Burt C V Non specific granuloma of the gastrointestinal tract *Surg Gynec & Obst* 57 71-80 1933
- 82 Erskine F B The pathologic relationship of me enteric adenitis, ileitis and appendicitis *Am J Clin Path* 11 706-712 1941
- 83 Faber K Pernicious Anemia som folge af tarmlidelse *Hospital studende* 601 1890 *Fjerde Roekke Bind* 111
- 83a. ——— Pernicose anemie bei dunndarmstrikturen *Berl Klin Wchnschr* 34 643-646, 1897

- 127 Jellen J Regional ileitis A review of fifty cases Am J Ro ntg and Rad 37 190-201 1937
- 128 Johnson W R Chronic non specific jejunitis with unu al features Gastroenterology 1 346 1943
- 129 Kantor J L Regional (terminal) ileitis Its roentgen diagnosis J A M A 103 2016-2021 1934
- 130 Kibava A K Po sible case of Crohn's diseas in a Ruando native East African M J 23 317 1946
- 131 Kiefer E D The diagnosis of regional ileitis (chronic idiopathic ulcerative enteritis) Surg Clin North America 23 862-872 1943
- 131a ——— Regional ileitis Clinics 3 506-515 1944
- 132 ——— and Ross J R Criteria in the management of chronic ileiti J A M A 129 104-109 1945
- 133 Kilham S T and Ingelfinger F J Nutritional problems presented by patient with extensive jejuno-ileitis Arch Int Med 73 466 1944
- 134 Kirklin B R Discussion of a paper by Brown I W Am J Roent gen 54 496 1945
- 135 Kirsner J B Owens F M and Humphrey F M Regional enteritis in father and son Gastroenterology 10 883-891 1948
- 136 Koenig A S Jr Chronic jejunitis Arch Path 10 187-190 1945
- 137 Koletsky S and Strecher R M Primary systemic amyloidosis Arch Path 27 267-283 1939
- 138 Konjetzny G F Discussion Zentr Chir 59 2534 1937
- 139 Korte W Ueber entzündliche Geschwulste am Darm Arch Klin Chir 118 138, 1921
- 140 Koster, H., Kasman I I and Sheinfeld W Regional ileitis Arch Surg 37 789 1936
- 141 Kross I Terminal Ileitis Conservative surgical treatment Am J Dig Dis & Nutr 313-314 1938-39
- 142 Ladd W E Quoted by Humans and Hass New England J Med 209 13,3 1933
- 143 Lahey F H and Sanderson E Lesions of the right colon involving right colectomy J A M A 120 1356 1942
- 144 Landois F Zirkumscription Enteritis am Dickdarm Zentr Chir 9 816 1923
- 145 ——— Ueber Ileitis Ulzera Zentr f Chir 61 1690 1937
- 146 Lehman F I Regional enteritis The acute phase Rev Gastro enterology 6 222 1939
- 147 Lenartowicz J and Rothfeld J Diffuse Boeck's sarcoid Arch Derm u Syph 161 504 1930
- 148 Lewisohn R Segmental enteritis Surg Gynec & Obst 66 215-22- 1938
- 149 ——— and Colp R In Prognosis in Regional Ileitis Crohn B B Am J Dig Dis & Nutrition 3 738 1939
- 150 Lakely D S and Liza J R Chronic granuloma Am J Dis Dis 6 113 1939

10. Golob, M Infectious granuloma of the intestines with special reference to the difficulty of pre operative differential diagnosis *M J & Rec* 135 390, 1932
- 106 Groen, J, and Pompen A W M Ileitis regionales *Geneeskundige Bladen Haarlem de evren F Bohn N V* 1936
- 107 von Haberer H Unspezifische Entzündung des Ileozökums *Muench Med Wchnschr* 81 479-483 1934
- 108 Hadfield, G The primary histological lesion of regional ileitis *Lancet* 2 773 775, 1939
- 109 Halligan E J and Halligan H J Acute free perforation as first sign of regional enteritis *Am J Surg* 57 493-497 1937
- 110 Holloway J W Regional ileitis *Ann Surg* 118 329 1943
- 111 Harris F I The earliest acute stage of regional enteritis Report of case in a child *Arch Pediat* 57 367-376 1940
- 112 ——— Bell, G H and Brunn H Chronic cicatrizing enteritis Regional ileitis (Crohn) A new surgical entity *Surg, Gynec & Obst* 57 637 1933
- 113 Holm C E The fate of the side tracked loop of ileum following lateral anastomosis for complete benign obstruction *Surg Gynec & Obst* 56 746 1933
- 114 Holman C C Survival after removal of twenty feet of intestine *Lancet* 2 597, 1944
- 115 Homans I and Hass G M Regional ileitis A clinical not a pathological entity *New England J Med* 209 1315-1324 1933
- 116 Horn R C Jr and Rhoads J E Regional enteritis involving Meckel's diverticulum perforation of the diverticulum and fistula formation *Ann Surg* 119 274-278 1944
- 117 Hurst A F Regional ileitis (Crohn's disease) I and II *Guy's Hosp Rep* 89 54-70 1939
- 118 ——— A case of Addison's anemia with subacute combined degeneration of the spinal cord and normal gastric secretion following chronic obstruction of the ileum *Guy's Hosp Rep* 87 175 1937
- 119 Husebye O W On roentgen diagnosis of jejunitis acuta phlegmonosa *Acta Radiolog* 29 71 1948
- 120 ——— Case of chronic non specific jejunitis and stenosis *Acta Radiolog* 29 516-518 1948
- 121 Jackman R J Anal abscess and anal fistula in association with regional ileitis Report of case *Proc Staff Meet Mayo Clinic* 19 154-156 1943
- 122 ——— and Buie L A Tuberculosis and anal fistula *J A M A* 130 630-632 1946
- 123 ———, and Smith N D Some manifestations of regional ileitis observed sigmoidoscopically *Surg Gynec & Obst* 76 444 1943
- 124 Jackman W A Localized hypertrophic enteritis as a cause of intestinal obstruction with a report of two cases *Brit J Surg* 22 27-32 1934
- 125 Jackson A S Regional enteritis *Surg Gynec & Obst* 6, 1-10 1937
- 126 Janus W L Regional jejunitis *Radiology* 50 532-534 1948

- 127 Jellen J Regional ileitis. A review of fifty cases. *Am J Roentgenol* 37 190-201 1937
- 128 Johnson W R Chronic non specific jejunitis with unusual features. *Gastroenterology* 1 346 1943
- 129 Kantor J L Regional (terminal) ileitis. Its roentgen diagnosis. *J A M A* 103 2016-2021 1934
- 130 Kibaya A K Possible case of Crohn's disease in a Ruando native. *East African M J* 23 317 1946
- 131 Kiefer E D The diagnosis of regional ileitis (chronic idiopathic ulcerative enteritis). *Surg Clin North America* 23 862-872 1943
- 131a ——— Regional ileitis. *Clinics* 3 506-515 1944
- 132 ——— and Ross J R Criteria in the management of chronic ileitis. *J A M A* 129 104-108 1945
- 133 Killian S T and Ingelfinger F J Nutritional problems presented by patient with extensive jejuno ileitis. *Arch Int Med* 73 466 1944
- 134 Kirklin B R Discussion of a paper by Brown P W. *Am J Roentgenol* 54 496 1945
- 135 Kirsner J B Owens F M and Humphreys E M Regional enteritis in father and son. *Gastroenterology* 10 883-891 1948
- 136 Koeng A S Jr Chronic jejunitis. *Arch Path* 10 187-190 1945
- 137 Koletsky S and Streeher R M Primary systemic amyloidosis. *Arch Path* 10 267-283 1949
- 138 Konjetzny, G F Discussion. *Zentr Chir* 59 2534 1937
- 139 Korte W Ueber entzündliche Geschwülste am Darm. *Arch Klin Chir* 113 138, 1921
- 140 Koster H Kasman L P and Sheinfeld W Regional ileitis. *Arch Surg* 37 789 1936
- 141 Kross I Terminal ileitis. Conservative surgical treatment. *Am J Dig Dis & Nutr* 5 313-314 1938-39
- 142 Ladl W E Quoted by Homans and Hass. *New England J Med* 109 1373 1933
- 143 Lahay F H and Sanderson E Lesions of the right colon involving right colectomy. *J A M A* 120 1356 1942
- 144 Landou F Zirkumskripte Enteritis am Dickdarm. *Zentr Chir* 0 816 1923
- 145 ——— Ueber Ileitis Ulcerosa. *Zentr f Chir* 64 1690 1937
- 146 Lehman E P Regional enteritis. The acute phase. *Rev Gastroenterology* 6 222 1939
- 147 Lenartowicz J and Rothfeld J Diffuse Boeck's sarcoid. *Arch Derm u Syph* 161 504 1940
- 148 Lewisohn R Segmental enteritis. *Surg Gynec & Obst* 66 215-227 1938
- 149 ——— and Colp R In Prognosis in Regional Ileitis. *Crohn B B Am J Dig Dis & Nutrition* 739 1936
- 150 Lohely D S and Liza J R Chronic granuloma. *Am J Dig Dis* 6 113 1939

- 105 Golob M *Infectious granuloma of the intestines, with special reference to the difficulty of pre operative differential diagnosis* M J & Rec 135 390 1932
- 106 Groen, J, and Pompen, A W M *Ileitis regionales* Geneeskundige Bladen Harlem de evren F Bohn N V 1935
- 107 von Haberer H *Unspezifische Entzündung des Ileozökum* Muench Med Wchnschr 81 479-483 1934
- 108 Hadfield G *The primary histological lesion of regional ileitis* Lancet 2 773 775 1939
- 109 Halligan E J and Halligan H J *Acute free perforation as first sign of regional enteritis* Am J Surg 37 493-497 1937
- 110 Halloway J W *Regional ileitis* Ann Surg 118 329 1943
- 111 Harris F I *The earliest acute stage of regional enteritis* Report of case in a child Arch Pediat 57 367-376 1940
- 112 ——— Bell G H and Brunn H *Chronic cicatrizing enteritis Regional ileitis (Crohn) A new surgical entity* Surg Gynec & Obst 57 637 1933
- 113 Holm C E *The fate of the side tracked loop of ileum following lateral anastomosis for complete benign obstruction* Surg Gynec & Obst 56 746 1933
- 114 Holman C C *Survival after removal of twenty feet of intestine* Lancet 2 597, 1944
- 115 Homans I and Hays G M *Regional ileitis A clinical not a pathological entity* New England J Med 209 1315-1324 1933
- 116 Horn R C Jr and Rhoads J F *Regional enteritis involving Meckel's diverticulum perforation of the diverticulum and fistula formation* Ann Surg 119 274-278 1944
- 117 Hurst A F *Regional ileitis (Crohn's disease) I and II* Guy's Hosp Rep 89 54-70 1939
- 118 ——— *A case of Addison's anemia with subacute combined degeneration of the spinal cord and normal gastric secretion following chronic obstruction of the ileum* Guy's Hosp Rep 87 175 1937
- 119 Husebye O W *On roentgen diagnosis of jejunitis acuta phlegmonosa* Acta Radiolog 29 71 1948
- 120 ——— *Case of chronic nonspecific jejunitis and stenosis* Acta Radiolog 29 516-518 1948
- 121 Jackman R J *Anal abscess and anal fistula in association with regional ileitis* Report of case Proc Staff Meet Mayo Clinic 18 154-156 1943
- 122 ——— and Buie L A *Tuberculosis and anal fistula* J A M A 150 630-632 1946
- 123 ——— and Smith N D *Some manifestations of regional ileitis observed sigmoidoscopically* Surg Gynec & Obst 76 444 1943
- 124 Jackman W A *Localized hypertrophic enteritis as a cause of intestinal obstruction with a report of two cases* Brit J Surg 22 27-32 1934
- 125 Jackson A S *Regional enteritis* Surg Gynec & Obst 65 1-10 1937
- 126 Jamieson W L *Regional jejunitis* Radiology 50 32-34 1948

- 17 Jellen J Regional ileitis A review of fifty cases Am J Roentg and Rad 37 190-201 1937
- 128 Johnson W R Chronic non specific jejunitis with unusual features Gastroenterology 1 346 1913
- 129 Kantor J L Regional (terminal) ileitis Its roentgen diagnosis J A M A 103 2016-2021 1934
- 130 Kabaya, A K Possible case of Crohn's disease in a Ruando native East African M J 23 317 1946
- 131 Kiefer E D The diagnosis of regional ileitis (chronic rhopathic ulcerative enteritis) Surg Clin North America 33 867-872 1943
- 131a ——— Regional ileitis Clinics 3 506-515 1944
- 132 ——— and Ross J R Criteria in the management of chronic ileitis J A M A 129 104-108 1945
- 133 Kallian S T, and Ingelfinger F J Nutritional problems presented by patient with extensive jejuno ileitis Arch Int Med 73 466 1944
- 134 Kirklin B R Discussion of a paper by Brown P W Am J Roentgen 34 496 1945
- 135 Kirsner J B, Owens F M and Humphrys F M Regional enteritis in father and son Gastroenterology 10 883-891 1948
- 136 Koenig A S Jr Chronic jejunitis Arch Path 40 187-190 1945
- 137 Koletsky S and Strecher R M Primary systemic amyloidosis Arch Path 37 267-283 1939
- 138 Konjetzny, G F Discussion Zentr Chir 59 2334 1932
- 139 Korte, W Ueber entzündliche Geschwülste am Darm Arch Klin Chir 118 138 1921
- 140 Koster H Kasman L P and Sheinfeld W Regional ileitis Arch Surg 3 789 1936
- 141 Kroes I Terminal ileitis Conservative surgical treatment Am J Dig Dis & Nutr 5 313-314 1938-39
- 142 Ladd W E Quoted by Homan and Hass New England J Med 309 1323 1933
- 143 Labey F H and Sanderson E Lesions of the right colon involving right colectomy J A M A 110 1356 1941
- 144 Landois F Zirkumskripte Enteritis am Dickdarm Zentr Chir 0 816, 1923
- 145 ——— Ueber Ileitis Ulcerosa Zentr f Chir 64 1690, 1937
- 146 Lehman E P Regional enteritis The acute phase Riv Gastroenterology 6 22, 1939
- 147 Lenartowicz J and Rothfeld J Diffuse Boeck's sarcoid Arch Derm u Syph 161 504 1930
- 148 Lewisohn R Segmental enteritis Surg Gynec & Obst 60 215-222 1938
- 149 ——— and Colp R In Diagnosis in Regional Ileitis Crohn B B Am J Dig Dis & Nutrition 3 738 1936
- 150 Likely D S and Liza J R Chronic granuloma Am J Dig Dis 6 113 1939



- 151 Llombart A, and Mineru, J Sur la localisation de la maladie de Nicolas Favre sur l'intestine grele, stenosis ileales multiples Ann d Anat Path 16 597 1939
- 152 Longcope, W T and Pierson, J W Boeck's sarcoid (sarcoidosis) Bull Johns Hopkins Hosp 60 223-296, 1937
- 153 Lowen, A Uber Appendicitis Fibro plastica Deutsch Ztschr f Chir 129 221-241, 1914
- 154 Lust F J Roentgenological studies of the mucosa of the normal terminal ileum Am J Roentg and Rad 45 63-68, 1941
- 155 Lyall, A Regional ileitis Glasgow M J 143 1-20 1945
- 156 Lyons, A S and Garlock J H Localized proximal jejunitis Arch Surg 53 702-704 1946
- 157 McGuff P Dockerty M B Waugh J M, and Randall L M Endometriosis as a cause of intestinal obstruction Surg Gynec & Obst 86 273 1948
- 158 McLeon C E Carcinoid tumors of the ileum Am J Clin Path 11 310 1944
- 159 McKinnon D C Spontaneous cure of acute regional enteritis Minnesota Med 74 744-746 1939
- 160 Mackie, T T Vitamin deficiency and the small intestine J A M A 117 910 1941
- 161 ——— and Pound R E Changes in the gastro-intestinal tract in deficiency states J A M A 101 813-816 1935
- 162 Mailer R Acute regional ileitis A report of two cases, with bacteriological findings Brit J Surg 25 517 1937
- 163 Marshall S F Regional ileitis New England J Med 222 375-382, 1940
- 164 ——— Regional ileitis Surgical management and results of operation S Clin North America 23 873-880, 1943
- 165 Mayo, C W and Judd F S Surgical resection in non specific ileocolitis J A M A 117 836 1941
- 166 Meide R H Jr Acute segmental enteritis Pennsylvania M J 44 1519-1526 1940-41
- 167 Megret R Ileite terminal aigue primitive chez un enfant de trois ans Mem Acad de Chir 65 380-382 1939
- 168 Meulengracht F Pernicious anemia in intestinal strictures Acta Med Scand 23 231-240 1929
- 169 Meyer K A and Ross P A Regional enteritis (non specific) S Clin North America 15 697-711 1935
- 170 Mixter C G Regional ileitis Ann Surg 102 674-694 1935
- 171 Mock H F Infective granuloma Surg Gynec & Obst 5 672-689 1931
- 172 Mokswoth H W L Granuloma of intestine Stenosis of ileo cecal valve Brit J Surg 21 370-372 1933-34
- 173 Monks G H Studies of surgical anatomy of the small intestine and its localization Ann Surg 12 543-569 1905
- 174 Morhead R I Regional enteritis with observations concerning pathogenesis North Carolina M J 9 216 1947

- 145 Morland A A case of arcoidosis of the lung with regional ileitis  
Tubercle London 29 3~ 1947
- 176 Morlock, C G Burgen J A and Pemberton J de J Regional  
enteritis following severe external violence Proc Staff Meet  
Mayo Clinic 14 631-635 1939
- 177 Moschcowitz E Amyloidosis in ulcerative colitis Ann Int Med  
10 73-88 1936
- 178 ——— Terminal ileitis Clinical recovery without operation four  
years after onset J Mt Sinai Hosp 7 77-80 1940-41
- 179 ———, and Wilensky A O Non specific granulomata of the intes-  
tine Am J M Sc 166 48-66 1923
- 180 Movniban B G A The mimicry of malign disease in the large  
intestine Edinburgh M J 21 228 1907
- 181 Mummery L Diseases of the Rectum and Colon London William  
Ward and Co 1923 P 538
- 182 Musick V H Combined non specific ulcerative colitis J Okla  
homa State M A 29 280-283 1936
- 183 Olson E S, and Sussman M L Non specific enterocolitis A J  
Roentg and Rad 60 471-486 1948
- 184 Oppenheimer G D Recurrent or residual progressive ileitis J A  
M A 110 1103-1104 1939
- 185 ——— Non specific localized granulomatous ulcers of the jejunum  
J Mt Sinai Hosp 1~ 1039-1041 1946
- 186 Pate J C Massive resection of small intestine Excision of 12 ft  
6 in with recovery J Florida M A 29 28-31 1942
- 187 Patricelli L Regional ileitis with Meckel's diverticulitis North  
west Med 14 117 1945
- 188 Pemberton J de J and Brown F W Regional ileitis Ann Surg  
10 855-870 1937
- 189 Iennington R E and Frielty J T Multiple carcinoid tumors of  
the small intestine Proc Staff Meeting Mayo Clinic 18 49-51  
1943
- 190 Penner A and Crohn B B Perianal fistulas as a complication  
of regional enteritis Ann Surg 109 867 1938
- 191 Passagno D J Regional ileitis with involvement of the cecum  
South M J 10 105-1055 1937
- 192 Peters K O Drei Weitere Falle von Darmphlegmone im Ileum  
Zentr Chir 61 1208 1934
- 193 Flum P and Warburg E Hematological changes especially meg-  
alocytic anemia in regional ileitis Acta Med Scand 10~ 449-  
475 1949
- 194 Topley J K Reproduction of ulcerative colitis in dogs Arch Surg  
43 551 1941
- 195 Prikau W H Massive resection of the small intestine Report of  
two cases Tr South Surg A 94-99 1944
- 196 Frohstein J J and Cruenfeld C F Acute regional ileitis Ann  
Surg 103 273-78 1936
- 197 Pugh H I Regional enteritis Ann Surg 1 845-861 1945

- 141 Ilombart A and Maneru J Sur la localisation de la maladie de Nicolas Favre sur l'intestine grele stenosis ileales multiples Ann d Anat Path 16 597 1939
- 142 Longcope W T and Pierson, J W Boeck's sarcoid. (sarcoidosis) Bull Johns Hopkins Hosp 60 223-296 1937
- 143 Lowen A Ueber Appendicitis Fibroplastica. Deutsch Ztschr f Chir 129 221-241 1914
- 144 Lust, F J Roentgenological studies of the mucosa of the normal terminal ileum Am J Roentg and Rad 45 63-68 1941
- 145 Lvall A Regional ileitis Glasgow M J 113 1-20 1945
- 146 Lyons A S and Garlock J H Localized proximal jejunitis Arch Surg 53 702-704 1946
- 147 McGuff P Dockerty M B Waugh J M and Randall L M Endometriosis as a cause of intestinal obstruction Surg Gynec & Obst 86 273 1948
- 148 McLeon C E Carcinoid tumors of the ileum Am J Clin Path 11 310 1944
- 149 McKinnon D C Spontaneous cure of acute regional enteritis Minnesota Med 744-746 1939
- 150 Mackie T T Vitamin deficiency and the small intestine J A M A 117 910 1941
- 151 ——— and Pound R E Changes in the gastro-intestinal tract in deficiency states J A M A, 101 813-816 1935
- 152 Mailer, R Acute regional ileitis A report of two cases with bacteriological finding Brit J Surg 25 517 1937
- 153 Marshall S F Regional ileitis New England J Med 222 350-352 1940
- 154 ——— Regional ileitis Surgical management and results of operation S Clin North America, 33 873-880 1943
- 155 Mayo C W and Judd F S Surgical resection in non specific ileocolitis J A M A 117 836 1941
- 156 Meade R H Jr Acute segmental enteritis Pennsylvania M J 44 1519-1526 1940-41
- 157 Megret R Ileite terminal aigue primitive chez un enfant de trois ans Mem Acad de Chir 65 380-382 1939
- 158 Meulengracht F Pernicious anemia in intestinal strictures Acta Med Scand 31-240 1939
- 159 Meyer K A and Rosi P A Regional enteritis (non specific) S Clin North America, 15 697-711 1935
- 160 Mixter C G Regional ileitis Ann Surg 10, 674-694 1935
- 161 Mock H E Infective granuloma Surg Gynec & Obst 672-689 1931
- 162 Molesworth H W L Granuloma of intestine Stenosis of ileocecal valve Brit J Surg 21 370-372 1933-34
- 163 Monks, G H Studies of surgical anatomy of the small intestine and its localization Ann Surg 4, 543-569 1905
- 164 Morehead R F Regional enteritis with observation concerning pathogenesis North Carolina M J 8 216 1947

- 175 Morland A A case of sarcoidosis of the lung with regional ileitis  
Tubercle London 28 32 1947
- 176 Morlock C G Borgen J A and Pemberton J de J Regional  
enteritis following severe external violence Proc Staff Meet  
Mayo Clinic 14 631-635 1939
- 177 Moschowitz E Amyloidosis in ulcerative colitis Ann Int Med  
10 73-88 1936
- 178 ——— Terminal ileitis Clinical recovery without operation four  
years after onset J Mt Sinai Hosp 77-80 1940-41
- 179 ———, and Wilenky A O Non specific granulomata of the intes-  
tine Am J M Sc 166 48-66 1923
- 180 Moynihan B G A The mimicry of malignant disease in the large  
intestine Edinburgh M J 21 928 1907
- 181 Mummery L Diseases of the Rectum and Colon London William  
Ward and Co 1923 P 538
- 182 Musick V H Combined non specific ulcerative colitis J Okla  
homa State M A 29 290-293 1936
- 183 Olsan E S and Sussman M L Non specific enterocolitis A J  
Roentgen and Rad 60 471-486 1948
- 184 Oppenheimer, G D Recurrent or residual progressive ileitis J A  
M A 110 1103-1104 1938
- 185 ——— Non specific localized granulomatous ulcers of the jejunum  
J Mt Sinai Hosp 12 1039-1041 1946
- 186 Pate J C Massive resection of small intestine Excision of 1<sup>2</sup> ft  
6 in with recovery J Florida M A 3 28-31 1941
- 187 Patricelli L Regional ileitis with Meckel's diverticulitis North  
west Med 44 117 1945
- 188 Pemberton J de J and Brown P W Regional ileitis Ann Surg  
10 855-870 1937
- 189 Pennington R E and Priestly J T Multiple carcinoid tumors of  
the small intestine Proc Staff Meeting Mayo Clinic 18 49 51  
1943
- 190 Penner A and Crohn B B Perianal fistulas as a complication  
of regional enteritis Ann Surg 109 867 1938
- 191 Pessagno D J Regional ileitis with involvement of the cecum  
South M J 30 1052 1055 1937
- 192 Peters K O Drei Wastere Fall von Darmphlegmone im Ileum  
Zentr Chir 61 1208 1934
- 193 Flum P and Warburg F Hematological changes especially meg-  
alocytic anemia in regional ileitis Acta Med Scand 101 440-  
445 1939
- 194 Poppe J K Reproduction of ulcerative colitis in dogs Arch Surg  
43 551 1941
- 195 Prioleau W H Massive resection of the small intestine Report of  
two cases Tr South Surg A 55 84-88 1944
- 196 Irobstein J J and Cruentfeld G E Acute regional ileitis Ann  
Surg 103 243-278 1936
- 197 Pugh H I Regional enteritis Ann Surg 111 845-861 1945

- 198 Pumphrey, R E Studies on the etiology of regional ileitis Proc Staff Meet Mayo Clinic 13 539-541 1938
- 199 Raffensberger, F C Recurrence in terminal ileitis following pregnancy Gastroenterology 10 1010 1948
- 200 Ravdin I S and Johnston C G (Shapiro) Regional ileitis A summary of the literature Am J M Sc 198 269-292 1939
- 201 ——— and Rhoads J F Regional ileitis and fibroplastic appendicitis Ann Surg 106 394-406 1937
- 202 Rees V L Regional jejunitis—Report of an unusual case Am J Surg 67 119-122 1945
- 203 Reichert F L and Mathes, M E Experimental lymphedema of the intestinal tract and its relation to regional cicatrizing enteritis Ann Surg 101 601-616 1936
- 204 Rhoads, J E The management of regional ileitis and certain other ulcerative lesions of the intestines Pennsylvania M J 42 1050-1053 1939
- 205 Richardson E P Jejunal ulcer Surg Gynec & Obst 35 1-10, 1922
- 206 Robinson D R and Wise W D Simple non specific ulcers of the jejuno ileum Surg Gynec & Obst 70 1097 1940
- 207 Rockey, E W Thickening of terminal ileum with mesenteric adenitis in children Northwest Med 32 145-147, 1933
- 208 Rodanich E C Kirsner, J B and Palmer, W L The relationship between lymphogranuloma venereum and regional enteritis an etiologic study of 4 cases with negative results Gastroenterology 1 657-689 1943
- 209 Rosenblate A J Goldsmith A A, and Strauss, A A A summary of regional ileitis J A M A 106 1797-1800, 1936
- 210 Rosenberg S A Enterocœcums at the ileo cecal junction Ann Surg 119 873 1944
- 211 Ros K Regional ileitis with report of a case M J Australia 1 321, 1936
- 212 Ruffin J M Baylin G J and Cayer D The clinical significance of alteration of the small intestine pattern as demonstrated by x ray Gastroenterology 4 289 1945
- 213 Sarnoff J Extensive resection of the small intestine Report of recovery after resection of fifteen feet of small intestine and hysterectomy Ann Surg 78 745-750 1923
- 214 Shearer J P and Jackson J T Recurrent regional (terminal) ileitis Ann Surg 106 459-461 1937
- 215 Schepers G W H The pathology of regional ileitis Am J Dig Dis 12 97-116 1940
- 216 Schiff E Regional enteritis Ann Pediatr 164 23 1945 165 281 1945
- 217 Sherrill J G and Hall, D P Regional ileitis Am J Surg 48 669-674 1940
- 218 Sinaiko E S and Neeches H Experiments in ulcerative enteritis Surgery 30 390 1946
- 219 Smithy, H G Conservatism in the surgical management of acute regional enteritis Surgery 13 122-130 1943

- 20 Snapper I Pseudotuberculosis in Man Amsterdam F B nn 1935
- 21 ——— Pompen A W M and Groen J Ileite regionale Van de Med 39 2-23 1936
- 22 Snavely J R Diarrhea and abdominal tenderness Bull Tulane Med Fac 6 22 1946
- 23 Snell A M and Camp J B Chronic idiopathic catarrhea Roentgenologic observation Arch Int Med 53 615-629 1934
- 24 Snielsen H and Ryan J Regional ileitis Presume of present knowledge and the addition of twenty two cases from Broome County New York Am J Surg 34 44-432 1941
- 25 Spellberg M A and Gray L W Regional enteritis of the proximal jejunum following trauma Surgery 17 343-350 1945
- 26 ——— and Ochner A The role of trauma as a possible etiologic factor in regional ileitis The effect of non penetrating trauma on the small intestine of dogs Am J M Sc 215 579 1947
- 27 Sprague P H, Anderson W S and Aaron T H Long standing fever due to regional ileocolitis Am J Dig Dis 11 295 1944
- 28 Sproull J A review of some features of regional ileitis with report of a case involving approximately three feet of the mid enteron Am J Roentgen 36 910-920 1936
- 29 Stafford E S Lymphogranuloma venereum Bull Johns Hopkins Hosp 62 399 1938
- 30 Starr A Editorial Is there an adequate therapy for regional enteritis? Surg Gynec & Obst 37 351-352 1948
- 31 Stierlin E and Chaoul H Klinische Roentgendiagnostik des Verdauungskanal Berlin J Springer 1938
- 32 Strombeck J P Ileitis terminalis Acta Chir Scandinav 80 Suppl 50 1-59 1957
- 33 ——— Terminal ileitis and its roentgen picture Acta Radiologica 22 827 1941
- 34 Sussman M L and Wahlst E Granulomatous jejuno-ileitis Radiology 39 49-53 1942
- 35 ——— and ——— Factors concerned in the abnormal distribution of barium in the small bowel Radiology 40 175 1943
- 36 Tallroth A Regional enteritis with special reference to its etiology and pathogenesis Acta Chir Scand 48 407-432 1943
- 37 Taylor A W Chronic hypertrophic ileo-cecal tuberculosis and its relation to regional ileitis (Crohn's disease) Brit J Surg 33 178-181 1945
- 38 Taylor R T Localized non specific ulcerative enteritis with some unusual features Am J Roentgen 35 884 1937
- 39 ten Kate J Two cases of terminal ileitis Nederl Tdchr v geneesk 80 5660-5664 1936
- 40 Tietze A Die Entzündlichen Geschwülste des Dickdarms Ergeb Chir u Orthop 1 11 1940
- 41 Toll W R Dittenbrant M Montague J R and West F S Digestion and absorption in a man with all but three feet of the small intestine removed surgically Am J Dig Dis 39 1940

- 198 Pumphrey, R E Studies on the etiology of regional ileitis Proc Staff Meet Mayo Clinic 13 339-541 1938
- 199 Raffensberger E C Recurrence in terminal ileitis following pregnancy Ga troenterology 10 1010 1948
- 200 Ravdin, I S, and Johnston, C G (Shapiro) Regional ileitis A summary of the literature Am J M Sc 198 269-292 1939
- 201 ——— and Rhoads J E Regional ileitis and fibro plastic appendicitis Ann Surg 106 394-406, 1937
- 202 Rees V L Regional jejunitis—Report of an unusual case Am J Surg 67 119-122 1945
- 203 Reichert F L and Mathes M E Experimental lymphedema of the intestinal tract and its relation to regional cicatrizing enteritis Ann. Surg 101 601-616 1936
- 204 Rhoads, J E The management of regional ileitis and certain other ulcerative lesions of the intestines Pennsylvania M J 12 1050-1053, 1939
- 205 Richardson E P Jejunal ulcer Surg Gynec & Obst 35 1-10, 1922
- 206 Robinson D R and Wise, W D Simple non specific ulcers of the jejunum Surg Gynec & Obst 70 1097 1940
- 207 Rockey E W Thickening of terminal ileum with mesenteric adenitis in children Northwest Med 30 145-147, 1933
- 208 Rodanich E C Kirsner J B and Palmer, W L The relationship between lymphogranuloma venereum and regional enteritis an etiologic study of 4 cases with negative results Gastroenterology 1 687-689 1943
- 209 Rosenblate A J Goldsmith A A, and Strauss V A A summary of regional ileitis J A M A 106 1797-1800, 1936
- 210 Rosenberg S A Enterogenous cysts at the ileo cecal junction Ann Surg 119 873 1944
- 211 Ross K Regional ileitis with report of a case M J Australia 1 321 1936
- 212 Ruffin J M Baylin G J and Caver D The clinical significance of alteration of the small intestine pattern as demonstrated by x ray Gastroenterology 4 289 1945
- 213 Sarnoff J Extensive resection of the small intestine Report of recovery after resection of fifteen feet of small intestine and hysterectomy Ann Surg 78 745-750 1923
- 214 Shearer J P and Jackson J T Recurrent regional (terminal) ileitis Ann Surg 106 459-461 1937
- 215 Schepers G W H The pathology of regional ileitis Am J Dig Dis 1 97-116 1945
- 216 Schiff E Regional enteritis Ann Pediatrics 164 23, 1945 165 281 1945
- 217 Sherrill J G and Hall D P Regional ileitis Am J Surg 48 669-674 1940
- 218 Sinaiko E S and Necheles H Experiments in ulcerative enteritis Surg 20 395 1946
- 219 Smithy H G Conservatism in the surgical management of acute regional enteritis Surgery 13 122-130 1943

# Index

- Abcesses
  - formation in ileo jejunitis 167
  - intra abdominal 84
  - multiple liver 83
- Actinomyces 1109 112
- Age incidence
  - in ileocolitis 204
  - in ileo jejunitis 166
  - in regional ileitis 28
- Agglutinations serum in ileitis 14
- Allergy possible causative agency 16
- Ambulatory treatment of ileitis 119
- Ankylosis 84
- Anemia
  - hyperchromic 173
  - pernicious 83
  - secondary 103 171
- Angulation intestinal in ileitis 57
- Appendectomy 47 197
- Appendicitis gangrenous 154
- Appendix possible etiologic relation hip 24
- Appetite 50
- Aphthous stomatitis 17
- Argentaffine tumors of ileum 111
- Arthralgia 14
- Arthritis 84
  - ly enterique 107
- Bacteria etiologic agency in ileitis 10 11
- Barium enema 98 99
- Barium meal 98 99
- Peeling oolite in ileitis 48
- Bile
  - excretion in regional ileitis 84
  - changes in ileo jejunitis 174
  - proteins in ileo jejunitis 176
  - transfusions 189
- Brucella 10
- Carcinoid of the terminal ileum 107 109 111
- Cardiac murmurs, significance of in ileitis 106
- Caseation lymph node 44
- Cecum involvement in ileitis 101
- Cholecystitis acute 83
- Cholelithiasis 83
- Clinical feature
  - ileocolitis 206
  - ileo jejunitis 169
  - regional ileitis 46
  - regional ileitis acute 144
- Clubbing of fingers 53 57 109 171 207
- Colitis
  - segmental 103
  - ulcerative 50 51 200
- Colo-ileitis *see* Ileocolitis
- Colostomy ileo-transverse 123 124 133 209
- Complications
  - eye 107
  - of regional ileitis 82
- Conjunctivitis phlyctenular 107 173
- Constipation 49 145
- Corneal ulceration 107
- Cultures stool 13
- Cyst enterogenous 102 109 112
- Deficiency states 93
- Dagnosis differential
  - of ileo jejunitis 182
  - of regional ileitis 103
- Diarrhea 46 70 103 145 171
- Diet
  - high protein in ileo jejunitis 169
  - in treatment of ileitis 115
- Duodenum involvement of in ileo jejunitis 161 163 170



- 242 Warren R, and Miller R H Regional enteritis New England J Med 226 589 1942
- 243 Warren S and Sommers S C Cicatrizing enteritis (regional ileitis) as a pathologic entity Am J Path 21 475-510 1948
- 244 Watson C J, Rigler L G Wangenstein, O H and McCartney J S Isolated sarcoidosis of the small intestine simulating non specific ileo jejunitis Gastroenterology 1 30-52, 1945
- 245 Waucho R L Terminal (regional) ileitis with report of a case Rev Gastroenterology 6 281-293, 1939
- 246 Weber H M In discussing article by Dixon C F and Weber H M Recurring obstruction from multiple non neoplastic tumor factions of the jejunum Proc Staff Meet Mayo Clinic 11 717-720, 1936
- 247 ——— Regional enteritis Roentgenologic manifestations Proc Staff Meet Mayo Clinic 13 545 1938
- 248 Weintraub S and Williams R A rapid method of roentgenologic examination of the small intestine Dept Radiology N Y Hosp In press
- 249 West F S Montague J R and Judy F R Digestion and absorption in a man with three feet of small intestine Am J Dig Dis & Nutr 5 690 1938
- 250 White W H Pathology and prognosis of pernicious anemia Cuy's Hosp Rep 47 149 1890
- 251 Wilensky A O Non specific granuloma of the intestine M J & Rec 135 445-446 1932
- 252 ——— Spontaneous healing in non specific granuloma of the terminal ileum Rev Gastroenterology 11 108 1944
- 253 Williams C Inflammatory tumors of the small intestine Virginia M Monthly 60 728-731 1934
- 254 Williams R H and Nickerson D A Sarcoidosis Proc Soc Exper Biol & Med 33 403-405 1935-1936
- 255 Wilmans Ein Fall von Darmstenose infolge chronisch entzündlicher Verdickung der Ileocecal Klappe Quoted by Tietze Fggeb d Chir u Orthop 12 212 1920
- 256 Wilson A K Gruman R B and Ashburn H C Regional ileitis South M J 35 881 1942
- 257 Yunch A M and Crohn B B Atypical regional ileitis Roentgenologic limitations Am J Dig Dis 8 185 1941



- Dysentery bacillary, 13, 14  
 Dysuria, 46  
 Efficiency loss of in ileitis, 52  
 Enterarteritis  
   obliterans, 45  
   subacute bacterial, 106  
 Endometrial transplants 111  
 Endometriosis 102 109  
 Entamoeba histolytica, 12  
 Enteritis  
   chronic electrizing 1  
   chronic ulcerative, 1  
   regional 1  
   segmental 1  
 Enterococcus : possible etiologic agent 11  
 Enterostomy emergency 124 132  
 Erythema 16 51 107 173  
 Erythema nodosum 57 143 173  
 Etiology  
   ileo colitis 204  
   ileo jejunitis 166  
   regional ileitis 4 9  
 Eye complications of ileitis 107  
 Familial incidence in regional ileitis 21  
 Fertility in ileitis 86  
 Fever 10 103 109 171 189 206  
 Fistula  
   abdominal wall 153 168 176  
   formation of 54 59 96  
   inguinal 60  
   internal terminal 59 61 177 207  
   to lumbar region 60  
   perianal 104  
   rectal and perirectal 59 67  
   recto vaginal 67 68 163 207  
   tuberculous 103  
   urinary bladder 59 64  
 Fibulous tracts 167  
 Folic acid 190  
 Foreign body 12 13  
 Frei test in diagnosis 14 15  
 Gastric analysis in ileo jejunitis 176  
 Grucher's disease, differential diagnosis 83  
 Giant cells, significance of, 4 13 41, 44  
 Gram positive cocci, 11  
*Granulomata, non specific* 2 4  
 Growth retarded  
   in ileitis 52  
   in ileo jejunitis 172  
*Gynecologic manifestations* 86  
 Healing spontaneous, 71 114, 134, 188  
 Hematemesis 46, 83 143  
 Hemorrhage 82 83 171  
 Hodgkins disease intestinal, 100 108  
 Hypoproteinemia 159 174 206  
 Hypovitaminosis 173  
 Ileitis terminal 1  
   regional see Regional Ileitis  
 Ileo colitis 1 5  
   clinical features 206  
   combined forms 200  
   pathologic distribution 204  
   physical signs 206  
   prognosis in 208  
   ex distribution 204  
   surgical treatment of 209  
   treatment medical 209  
 Ileo jejunitis 1 156 170  
   anemia in 171  
   blood changes in 174  
   blood protein in 176  
   blood transfusions in 189  
   clinical features 169  
   course in 186  
   differential diagnosis 176 182  
   high protein diet 189  
   isolated segmental involvement 165  
   mortality rates 186  
   nutrition in 172  
   pain abdominal 171  
   pathologic distribution 160

- physical examination 176  
 prognosis 186  
 resection massive 19\_  
 roentgenography of 178  
 sex distribution 166  
 skip lesions in 161  
 spontaneous cures 189  
 stomach involvement in 163  
 surgical treatment of 191  
 therapy of 189  
 treatment medical 188  
 upward extension 180  
 vitamin deficiency in 173-174  
 Ileo-jejunitis acute form 170  
 Ileo transverse colostomy 173 124  
     120  
 Ileum surgical transection of 126  
     198  
 Incubations animal 11  
 Intestinal  
     absorption in ile jejunitis 181  
     obstruction in ileitis 46 113  
 Intussusception 20 45 86  
 Iritis 107  
 Jejunitis  
     acute phlegmonous 170  
     isolated 163  
 John's disease 12  
 Keratitis 107  
 Life cycle in regional ileitis 113  
 Liver extract crude 120 190  
 Leucopodium 13  
 Lymphatic block experimental 21  
 Lymph nodes mesenteric 10 21  
     133 144  
 Lymphadenitis mesenteric 27 146  
 Lymphoedema experimental 6 21  
 Lymphopathia venereum 14  
 Lympho sarcoma differential diag-  
     nosis from ileitis 100 108  
 Malnutrition 208  
 Mass abdominal 54 57 104  
 Meckel's diverticulum 83  
 Melæna 46 83 209  
 Menstruation 53 80 173  
 Me enteric fat 37  
 Mesenteric lymphadenitis 27  
 Mesentery involvement in ileitis  
     152  
 Mikulicz surgical technique 123 121  
     130  
 Mortality rates in ileitis 82 121  
     197 129 130  
     in ileo jejunitis 186  
 Mucosal relief pattern 91  
 Mucosal type of ileitis 40 91  
 Nerve in ileitis 50  
 Necrosis in lymph node 44  
 Nerve tissue involvement in ileitis  
     41  
 Nonspecific protein therapy 208  
 Nomenclature enteritis  
     chronic ulcerating 1  
     chronic ulcerative 1  
     colitis 1  
     regional 1  
     segmental 1  
     terminal 1  
 Nutrition  
     in ileitis 70  
     in ileo jejunitis 17\_  
 Obstruction  
     of inferior vena cava 84  
     intestinal 50 57 70 114 12\_  
         163 172 177 207  
 Onset ileitis 46  
 Oration  
     experimental 153 192  
     short circuiting 154  
     two stage 130  
 Pain  
     abdominal 49 49 171  
     joints 53 81  
     prædefecation 48  
 Pathologic differentiation  
     in ileitis 204  
     in ileo jejunitis 160  
 Pathology  
     gross in ileo jejunitis 167

- in regional ileitis, 34
- microscopic in regional ileitis 41
- Penicillin therapy 121, 189
- Perforation, intestinal 46, 82
- Peritoneal fluid 82
- Peritonitis, 82
- Personality type in regional ileitis, 30
- Physical examination
  - in regional ileitis 54
  - in ileo jejunitis 176
- Polypos 102 167
- Pregnancy, relation to ileitis 86
- Prodroma of regional ileitis 47
- Prognosis
  - in acute ileitis 147
  - in combined ileo colitis 208
  - in ileo jejunitis 186
  - in regional ileitis 70
- Protein hydrolysate 115 120 209
- Psychic manifestations of ileitis 43
- Psychoneurosis in ileitis 30 46 43
- Purpura hemorrhagica 145
- Pyelophlebitis in ileitis 85
- Race incidence in regional ileitis 30 32
- Radiation therapy of ileitis 122
- Rectal complications 47 178 207
- Recto vaginal fistula see Fistula recto vaginal
- Recurrences in ileitis 124 128 130 131 134 147
  - treatment of 139
- Regional ileitis 1, 9
  - acute 142
    - clinical manifestations 144
    - prognosis 147
    - pathological involvement 144
    - roentgenography of 145
    - therapy of 152
  - anemia in, 51 103
  - blood calcium 84
  - blood changes 51
  - clinical features 46
  - complications of 82, 107
  - differential diagnosis, 100 103 108
  - familial incidence in, 21
  - intestinal obstruction in, 46
  - life cycle 113
  - medical therapy of, 113
  - mortality rates 82 125 127, 199
  - nutrition in 70
  - onset 46
  - pain abdominal, 48 49
  - pathology microscopic 41
  - pathology gross 34
  - perforation intestinal 46 82
  - personality type 30
  - physical examination 54
  - pregnancy 86
  - prodroma 47
  - prognosis and course 70
  - psychoneurosis in 30 46
  - pyelophlebitis in 85
  - race incidence in 30 32
  - remissions in 70 81
  - resection primary 124 130
  - roentgenography of 88 98 100 102
    - ex distribution 29 143
  - skip lesions 39 71, 91 97 144
  - social status 30
  - spontaneous cure 71 114 154
  - stricture in 155
  - string sign 6 88 91 96 98 101 145
  - submucosal type 40 81
  - surgical treatment of 193
  - vascular changes in 24
  - virus infection 14
  - vitamin deficiency in 100 108
- Religion in regional ileitis 30
- Remission in regional ileitis 10 81
- Resection
  - surgical in ileitis 123 144
  - massive in ileitis 192
  - primary in ileitis 124, 130
- Resolution in ileitis
  - complete 150
  - spontaneous 113 151

- Rheumatic heart disease 83  
 Roentgenography  
   acute ileitis 145  
   barium enema in 98  
   lead-saline solution in 100  
   ileo-jejunitis 178  
   negative finding 104  
   regional ileitis 88 98 100 102  
 Saline solution lead in roentgenography 100  
 Sarcoidosis 16 112  
 Secretory changes gastric 52  
 Sedimentation rate 51 176  
 Serum agglutinations 13 14  
 Serum lipids 184  
 Sex distribution  
   in ileo-colitis 204  
   in ileitis 29 143  
   in ileo-jejunitis 166  
 Sexual maturation 52 172  
 Skip lesions intestinal 39 71 91  
   91 144 161  
 Social status in regional ileitis 39  
 Spleen palpable 57 159 177 207  
 Sprue differential diagnosis 100  
   109 176 182  
 Stenosis intestinal cicatricial 46  
   113  
 Stomach possible involvement in  
   ileo-jejunitis 163  
 Streptococcus  
   aerobic 11  
   viridans 11 12  
 Streptomycin therapy 1 2 189  
 String sign in ileitis 6 88 91  
   96 98 101 145  
 Submucosal type of regional ileitis  
   40 81  
 Sulfasalazine therapy 1 - 189  
 Sulfathiazine 1 2 189  
 Sulfonamides insoluble 190 191  
 Surgical treatment of  
   end-to-end anastomosis 131  
   failures in 114  
   ileitis 123  
   ileo-jejunitis 191  
   ileo-colitis 207  
   two-stage procedure 123  
 Syphilis differential diagnosis 10  
 Tetany 84 174  
 Therapy  
   acute ileitis 1 2  
   ileo-colitis 209  
   ileo-jejunitis 188  
   indications for 115  
   medical of regional ileitis 113  
   radiation 122  
 Transfusions of blood 120  
 Trauma 17 169 172  
 Tuberculosis primary intestinal  
   109  
 Tuberculosis  
   hypertrophic ileo-cecal 10 101  
   intestinal 110  
 Tumor argentaffine 111  
 Two-stage procedure surgical 12  
 Ulcer  
   in ileitis 83  
   isolated jejunal 169  
 Typhoid fever *see* Brucellosis  
 Urinary bladder fistula 59 64  
 Valve of Pouch 89 90  
 Vascular changes in ileitis 24  
 Viruses possible etiologic agent  
   14  
 Vitamin  
   P complex therapeutic use 1 0  
   190  
   deficiency in ileitis 100 108  
   in ileo-jejunitis 173 174  
   K synthetic 120  
 Vitamins  
   fat soluble 1 4  
   replacement of 1 0  
 Vomiting 50 58  
 Weight loss 52 171

- in regional ileitis, 34
- microscopic, in regional ileitis 41
- Penicillin therapy, 121, 189
- Perforation, intestinal 46, 82
- Peritoneal fluid 82
- Peritonitis, 82
- Personality type in regional ileitis, 30
- Physical examination
  - in regional ileitis, 54
  - in ileo jejunitis 176
- Polyps 102 167
- Pregnancy relation to ileitis 86
- Prodromal of regional ileitis, 47
- Prognosis
  - in acute ileitis 147
  - in combined ileo colitis 208
  - in ileo jejunitis 186
  - in regional ileitis 70
- Protein hydrolysate 115 120 209
- Psychic manifestation of ileitis 33
- Psychoneurosis in ileitis 30 46 53
- Purpura hemorrhagica 145
- Pyclophlebitis in ileitis 83
- Race incidence in regional ileitis 30 32
- Radiation therapy of ileitis 122
- Rectal complications 47 178 207
- Recto vaginal fistulas *see* Fistula recto vaginal
- Recurrences in ileitis 124 128 130 131 134 137
  - treatment of 139
- Regional ileitis 1, 9
  - acute 142
    - clinical manifestations 144
    - prognosis 147
    - pathological involvement 144
    - roentgenography of 143
    - therapy of 152
  - anemia in 51 103
  - blood calcium 84
  - blood changes 51
  - clinical features 46
  - complications of 82, 107
  - differential diagnosis 100 103 108
  - familial incidence in 21
  - intestinal obstruction in, 46
  - life cycle 113
  - medical therapy of, 113
  - mortality rates 82, 123 127 109
  - nutrition in 70
  - onset 46
  - pain abdominal, 48, 49
  - pathology microscopic 41
  - pathology gross 34
  - perforation intestinal 46 82
  - personality type 30
  - physical examination 54
  - pregnancy 86
  - prodromal 47
  - prognosis and course 70
  - psychoneurosis in 30 46
  - pyclophebitis in 83
  - race incidence in 30 32
  - remissions in 70 81
  - resection primary 124 130
  - roentgenography of 88 98 100 102
  - sex distribution 29 143
  - skip lesions 39 71, 91 97 144
  - social status 30
  - spontaneous cure 71 114 134
  - stricture in 15
  - string sign 6 88 91 96 98 101 143
  - ulcerous type 40 81
  - surgical treatment of 123
  - vascular changes in 24
  - virus infection 14
  - vitamin deficiency in 100 108
- Religion in regional ileitis 30
- Remission in regional ileitis 10 81
- Resection
  - surgical in ileitis 123 134
  - massive in ileitis 192
  - primary in ileitis 124 130
- Resolution in ileitis
  - complete 150
  - spontaneous 113, 151

